

# Supporting Information

# Posterior Predictive Check




Figure S1: (Caption next page.)

Figure S1: (Previous page.) Posterior predictive check of germination rate from discretized log-normal model for the South cohort (left facets) and North cohort (right facets). Each histogram in gray bars is for one of five *M. cardinalis* populations arranged from northern (top) to southern (bottom). The black points are median values from randomly simulated data sets drawn from each iteration of the posterior distributions. Lines indicate the 95% confidence interval but are usually too narrow to be seen behind the point estimates. A close match between observations and posterior predictions indicate that the model adequately describes the distribution of the data.




Figure S2: The average temperature during the 2015-16 experiment (lines) was warmer than the historical normal from 1961-1990 (points). The  $y$ -axis is the average temperature for a season in the southern (solid line) and northern (dashed line) garden. Points are the average seasonal temperature for the 1961-1990 normal for each population source location. Population color is arranged from orange to blue by latitude of origin from south to north. Season is along the  $x$ -axis. Autumn encompasses September-November; winter encompasses December-February; spring encompasses March-May. All down-scaled climate are from ClimateNA (see Materials and Methods).

Table S1: We decomposed variance in germination rate among populations ( $V_{\text{pop}}$ ), additive genetic differences among individuals in the base population ( $V_G$ ), experimental blocks within the greenhouse (Block), and residual environmental variance ( $V_E$ ). The narrow-sense heritability  $h^2$  is  $V_G/(V_G + V_E)$ . Point estimates and 95% confidence intervals (CI) are estimated from the median and quantiles of the posterior distribution.

Parameter	Median (95% CI)
$V_{\text{pop}}$	0.139 (0.0877–0.203)
$V_G$	0.116 (0.0839–0.168)
Block	0.0354 (0.024–0.0556)
$V_E$	0.266 (0.258–0.275)
$H^2$	0.278 (0.215–0.358)

Table S2: We decomposed variance in winter survival ( $\text{logit}(p_{\text{surv}})$ ) among populations ( $V_{\text{pop}}$ ), additive genetic differences among individuals in the base population ( $V_G$ ), experimental blocks in the field (Block), and residual environmental variance ( $V_E$ ) for the North and South garden separately. The narrow-sense heritability  $h^2$  is  $V_G/(V_G + V_E)$ . Point estimates and 95% confidence intervals (CI) are estimated from the median and quantiles of the posterior distribution.

Population	Garden	Parameter	Median (95% CI)
Sweetwater River	North	$H^2$	0.00815 (0.00263–0.0205)
Sweetwater River	North	$V_E$	0.181 (0.15–0.217)
Sweetwater River	North	$V_G$	0.0015 (0.000512–0.00424)
Sweetwater River	North	$V_{\text{pop}}$	0.00743 (0.0045–0.0115)
Sweetwater River	North	Block	0.0015 (0.000512–0.00424)
West Fork Mojave River	North	$H^2$	0.0087 (0.00274–0.0209)
West Fork Mojave River	North	$V_E$	0.197 (0.159–0.226)
West Fork Mojave River	North	$V_G$	0.00175 (0.000563–0.00463)
West Fork Mojave River	North	$V_{\text{pop}}$	0.00743 (0.0045–0.0115)
West Fork Mojave River	North	Block	0.00175 (0.000563–0.00463)
North Fork Middle Tule River	North	$H^2$	0.00794 (0.00259–0.0202)
North Fork Middle Tule River	North	$V_E$	0.174 (0.134–0.213)
North Fork Middle Tule River	North	$V_G$	0.00141 (0.000464–0.00422)
North Fork Middle Tule River	North	$V_{\text{pop}}$	0.00743 (0.0045–0.0115)
North Fork Middle Tule River	North	Block	0.00141 (0.000464–0.00422)
Little Jamison Creek	North	$H^2$	0.00963 (0.00296–0.0226)
Little Jamison Creek	North	$V_E$	0.221 (0.191–0.24)

Population	Garden	Parameter	Median (95% CI)
Little Jamison Creek	North	$V_G$	0.0021 (0.000701–0.00546)
Little Jamison Creek	North	$V_{\text{pop}}$	0.00743 (0.0045–0.0115)
Little Jamison Creek	North	Block	0.0021 (0.000701–0.00546)
Rock Creek	North	$H^2$	0.0101 (0.00307–0.0236)
Rock Creek	North	$V_E$	0.241 (0.227–0.248)
Rock Creek	North	$V_G$	0.00247 (0.000765–0.00586)
Rock Creek	North	$V_{\text{pop}}$	0.00743 (0.0045–0.0115)
Rock Creek	North	Block	0.00247 (0.000765–0.00586)
Sweetwater River	South	$H^2$	0.000285 (5.3e-05–0.00326)
Sweetwater River	South	$V_E$	0.0644 (0.0393–0.101)
Sweetwater River	South	$V_G$	$2.07 \times 10^{-5}$ ( $2.95 \times 10^{-6}$ – $0.000305$ )
Sweetwater River	South	$V_{\text{pop}}$	0.00143 (0.000626–0.0028)
Sweetwater River	South	Block	0.00528 (0.00155–0.0205)
West Fork Mojave River	South	$H^2$	0.000316 ( $5.88 \times 10^{-5}$ – $0.00328$ )
West Fork Mojave River	South	$V_E$	0.0714 (0.0483–0.113)
West Fork Mojave River	South	$V_G$	$2.41 \times 10^{-5}$ ( $4.54 \times 10^{-6}$ – $0.000321$ )
West Fork Mojave River	South	$V_{\text{pop}}$	0.00143 (0.000626–0.0028)
West Fork Mojave River	South	Block	0.00645 (0.00201–0.0266)
North Fork Middle Tule River	South	$H^2$	0.000356 ( $6.54 \times 10^{-5}$ – $0.00334$ )
North Fork Middle Tule River	South	$V_E$	0.0844 (0.0534–0.122)

Population	Garden	Parameter	Median (95% CI)
North Fork Middle Tule River	South	$V_G$	$3.42 \times 10^{-5}$ ( $5.84 \times 10^{-6}$ – $0.000425$ )
North Fork Middle Tule River	South	$V_{\text{pop}}$	0.00143 (0.000626–0.0028)
North Fork Middle Tule River	South	Block	0.00894 (0.00264–0.031)
Little Jamison Creek	South	$H^2$	0.000381 ( $7.29 \times 10^{-5}$ – $0.00424$ )
Little Jamison Creek	South	$V_E$	0.0934 (0.0606–0.13)
Little Jamison Creek	South	$V_G$	$4.07 \times 10^{-5}$ ( $8.42 \times 10^{-6}$ – $0.000513$ )
Little Jamison Creek	South	$V_{\text{pop}}$	0.00143 (0.000626–0.0028)
Little Jamison Creek	South	Block	0.0108 (0.00354–0.0353)
Rock Creek	South	$H^2$	0.000525 (0.000104–0.00627)
Rock Creek	South	$V_E$	0.131 (0.1–0.159)
Rock Creek	South	$V_G$	$8.36 \times 10^{-5}$ ( $1.57 \times 10^{-5}$ – $0.000972$ )
Rock Creek	South	$V_{\text{pop}}$	0.00143 (0.000626–0.0028)
Rock Creek	South	Block	0.0225 (0.00926–0.0511)

Table S3: Between-population directional selection coefficient estimates and confidence intervals. In both North and South gardens, we estimated selection on increased germination rate (log-mean scale) caused by increased winter survival ( $\text{logit}(p_{\text{surv}})$ ). The parameters are the linear regression slope and intercept. Point estimates and 95% confidence intervals (CI) are estimated from the median and quantiles of the posterior distribution.

Garden	Parameter	Median (95% CI)
North	intercept	4.06 (3.09 – 5.26)
North	slope	-0.456 (-0.629 – -0.324)
South	intercept	4.9 (3.84 – 6.08)
South	slope	-0.355 (-0.53 – -0.225)

Table S4: Genotypic directional selection coefficient estimates and confidence intervals. In both North and South gardens, we estimated selection on increased germination rate (log-mean scale) caused by increased winter survival ( $\text{logit}(p_{\text{surv}})$ ). The parameters are the linear regression slope and intercept. Point estimates and 95% confidence intervals (CI) are estimated from the median and quantiles of the posterior distribution.

Garden	Parameter	Median (95% CI)
North	intercept	-0.0022 (-0.034 – 0.0273)
North	slope	-0.131 (-0.371 – 0.0122)
South	intercept	0.000691 (-0.0176 – 0.0266)
South	slope	-0.00177 (-0.14 – 0.13)