

Guard cell size and ~~pore aperture~~ initial conductance influence stomatal closure kinetics

Christopher D. Muir^{1,2*}

Wei Shen Lim¹

¹School of Life Sciences, University of Hawai‘i, Mānoa, Hawai‘i, USA

²Department of Botany, University of Wisconsin, Madison; Wisconsin, USA

*Corresponding author: cdmuir@wisc.edu

ORCID: Christopher D. Muir — [0000-0003-2555-3878](https://orcid.org/0000-0003-2555-3878)

ORCID: Wei Shen Lim — [0009-0001-1406-7957](https://orcid.org/0009-0001-1406-7957)

Keywords: ~~amphistomy~~ guard cell size, leaf gas exchange, phylogenetic heritability, *Solanum*, stomatal kinetics, ~~stomatal size~~ vapor pressure deficit

Summary

- In fluctuating environments, the kinetics of stomatal opening and closing influence the balance between carbon gain and water loss. ~~The rate of stomatal aperture change varies widely among plants, but the underlying causes of this variation are not well understood. One hypothesis is that smaller guard cells~~ Smaller guard cells may respond faster to fluctuating environmental conditions because of their greater surface area for osmolyte flux relative to cell volume. A related hypothesis is that operational stomatal conductance (g_{op}) is often well below its theoretical maximum (g_{max}) because at this stomatal aperture, guard cell volume is poised to change rapidly with small changes in turgor pressure. ~~The support for both hypotheses is limited and mixed, in part because they have not been tested together.~~
- We analyzed 2,124-125 estimates of stomatal closure kinetics in response to an abrupt increase in vapor pressure deficit (VPD) among 29 diverse wild tomato populations in the genus *Solanum*.
- ~~Both smaller guard cell size~~ Leaves with small guard cells and a lower g_{op} to g_{max} ratio (f_{gmax}) ~~were associated with faster stomatal closure kinetics, but~~ initial stomatal conductance (g_i) closed faster, but each explained variation in kinetic parameters at different levels of biological organization. Guard cell size had high phylogenetic heritability and varied relatively little within populations, whereas f_{gmax} g_i varied mostly among individuals and between light intensity treatments.
- ~~Together, these results indicate that smaller~~ Smaller stomata can be speedier, but ~~only if stomata are held at an aperture where they are responsive to changing turgor pressure. We predict that selection~~ the current physiological state of the leaf also affects the responsiveness of stomatal conductance to fluctuations. Selection on stomatal speed ~~will~~ may influence not only anatomical traits like guard cell size, but also physiological controls on g_{op} .

Introduction

25 In nature, change is the only constant. One way that vascular plants (tracheophytes) cope with change is by adjusting stomatal pore aperture to optimize the trade-off between carbon gain and water loss (Cowan & Farquhar, 1977; Raven, 2002; Hetherington & Woodward, 2003). In the absence of physical limits and energetic tradeoffs, natural selection would favor plants that could instantaneously adjust stomatal conductance (g_{sw}) to perfectly track dynamic environmental conditions (Lawson *et al.*, 2010).
30 In reality, stomatal responses take time, creating a lag between actual and optimal stomatal aperture. For example, many species use hydroactive feedback mechanisms to precisely control stomatal responses to changing vapor pressure deficit (VPD), the driving force for evaporation (Buckley, 2019). ~~Hydroactive stomatal control requires sensing, but hydroactive control takes time. Leaves must sense~~ a change in VPD through as-yet-unknown mechanisms ~~, signal transduction, and and then transduce a signal that~~
35 ~~causes~~ efflux of osmotic solutes to alter guard cell turgor more than the surrounding epidermal cells. Abscisic acid (ABA) dependent responses require several minutes to accumulate or synthesize hormone pools *de novo* (Xie *et al.*, 2006; Bauer *et al.*, 2013; McAdam *et al.*, 2016; Desai & Stroock, 2025; Pichaco *et al.*, 2026); ABA-independent stomatal responses require changes in gene expression and protein synthesis that take time (Jalakas *et al.*, 2021). Once anion channels are activated, the rate
40 of change in stomatal aperture depends on the rate of change in guard cell turgor (dP_g/dt) and the sensitivity of aperture to change in turgor (da_s/dP_g). As we discuss below, these two factors (dP_g/dt and da_s/dP_g) may be influenced by guard cell size and the ratio of operational to maximum stomatal conductance (f_{gmax}), respectively.

Over a century ago Darwin (1898) observed that after an initial increase, leaf transpiration decreases in
45 response to decreasing humidity. He correctly attributed changes in transpiration to changes in stomatal aperture and subsequent studies have confirmed that stomatal aperture responds to humidity (Lange *et al.*, 1971) and, more specifically, transpiration rate (Mott & Parkhurst, 1991; Monteith, 1995), which is the product of VPD and g_{sw} . The VPD is related to relative humidity (RH) but also temperature. In angiosperms and some pteridophytes, a sudden increase in VPD, usually induced by lower RH at a
50 constant temperature, causes a transient increase in g_{sw} , as Darwin observed (Cowan, 1972; Buckley *et al.*, 2011; Westbrook & McAdam, 2021). After a period of several minutes, stomata close until the leaf reaches a new steady-state g_{sw} . These phases of stomatal response to VPD are often referred to as the “wrong-way response” (WWR) and “right-way response” (RWR), respectively. The normative ‘wrong’ and ‘right’ appellations refer to the presumed adaptive significance. At steady state, g_{sw} is
55 negatively correlated with VPD, all else being equal, which is interpreted as an adaptive response to the marginal carbon cost of water (Givnish & Vermeij, 1976; Cowan & Farquhar, 1977; Buckley *et al.*, 2017b). Models built around this assumption generally predict steady state g_{sw} well, although the functional form and presumed fitness costs are areas of active research (Damour *et al.*, 2010; Medlyn *et*

al., 2011; Prentice *et al.*, 2014; Sperry *et al.*, 2017; Potkay *et al.*, 2025).

60 Stomatal responses are as much about the journey (kinetics) as they are about the destination (steady state). If stomatal response kinetics are too slow relative to the temporal grain size of environmental variability, then it would be futile for g_{sw} to track environmental variation since the leaf will constantly lag the environment, perpetually fighting the last war. This may explain why g_{sw} can be insensitive to short sun/shade flecks (Knapp & Smith, 1990) and shaded leaves are likely primed to cope with the
65 hydraulic demands of sudden increases in light, temperature, and VPD (Schymanski *et al.*, 2013). If stomatal responses can track environmental variation, then minimizing the lag time can greatly improve resource-use efficiency (Lawson *et al.*, 2010). For example, optogenetic control of stomatal aperture in response to fluctuating light increases water-use efficiency in *Arabidopsis* compared to wild-type responses (Papanatsiou *et al.*, 2019). Therefore the rate of stomatal opening and closing is predicted
70 to have a major effect on how natural selection optimizes stomatal control in variable environments (Buckley *et al.*, 2023).

Much of the research on stomatal kinetics focuses on light responses because light intensity is usually the most important, rapidly fluctuating environmental variable affecting photosynthesis. In nature, fluctuating light intensity ~~simultaneously~~ simultaneously alters leaf temperature and VPD (e.g. Ishida
75 *et al.*, 1992). We focus on stomatal closure kinetics in response to an abrupt ~~increased-in-VPD~~ increase in VPD, rather than light for four reasons. First of all, the primary aims and *a priori* hypotheses of this study were not to study stomatal kinetic parameters and are described in Muir *et al.* (2025). The idea to opportunistically use g_{sw} -response curves ~~generated during this study to test hypotheses about which traits influence to study~~ stomatal kinetics came after the data had been collected, but not yet analyzed
80 for this purpose. Second, VPD responses exhibit a similar functional form as light-responses, suggesting similar underlying mechanisms (Buckley, 2019). Factors that affect stomatal response kinetics to VPD likely generalize to other environmental variables. Third, humidity responses tend to dominate over and are faster than other environmental responses (Aasamaa & Söber, 2011a,b), ~~but~~ although there is variation among species (Merilo *et al.*, 2014). Thus, factors that constrain maximum rates of stomatal
85 closure are most likely to be observed in response to VPD. Finally, the ecological salience of VPD responses is increasing as climate change warms the atmosphere (Grossiord *et al.*, 2020).

Fully describing stomatal responses to the environment is complex (Buckley *et al.*, 2003; Jezek *et al.*, 2021; Desai & Stroock, 2025), but the dynamics of g_{sw} in response to an abrupt environmental change is well described by relatively simple phenomenological functions (Vico *et al.*, 2011; McAusland *et al.*,
90 2016). A four-parameter equation describes stomatal response kinetics to a step change in light intensity across dozens of plant species (Woning *et al.*, 2026):

$$g_{sw,t} = g_f + (g_i - g_f)e^{-(\frac{t}{\tau})^\lambda}. \quad (1)$$

In this equation, $g_{sw,t}$ is stomatal conductance at time t , g_i is the initial g_{sw} before the step change, g_f is the final g_{sw} after the step change, τ is a time constant that describes how quickly stomata respond to the step change, and λ is a shape parameter that describes how the response rate changes over time. When $\lambda = 1$, the response rate is constant over time and τ represents the time required for g_{sw} to reach 36.8% ($1/e$) of the way between initial and final values. When $\lambda > 1$, the response rate increases over time, and when $\lambda < 1$, the response rate decreases over time. Woning *et al.* (2026) refer to λ as a lag-time parameter because empirically $\lambda > 1$. We refer to stomatal kinetic parameters τ (time-constant) and λ (lag-time) henceforth. The first goal of this study is to evaluate whether kinetics of the RWR after a step change in VPD are adequately described by Equation 1. If so, it suggests that some fundamentally similar mechanisms are shared between light and hydroactive stomatal responses (Grantz & Zeiger, 1986).

The main goal of this study is to test hypotheses about which stomatal anatomical traits affect kinetic parameters τ and λ . We consider two interconnected hypotheses that have thus far been treated in isolation. The first hypothesis is that smaller guard cells open and close faster because of their intrinsically greater surface area to volume ratio (Franks & Farquhar, 2007; Drake *et al.*, 2013; Lawson & Vialet-Chabrand, 2019; Woning *et al.*, 2026). For approximately cylindrical cell geometries, like that found in guard cells, surface area increases linearly with radius, whereas volume increases in proportion to the radius squared. Consequently, larger guard cells will require more time, all else being equal, to achieve a given change in turgor pressure and, hence, stomatal pore aperture. A plant with large guard cells can partially compensate for this geometric constraint by increasing the activity and/or density of ion and solute transporters in the guard cell plasma membrane (Homann & Thiel, 2002; Lawson & Blatt, 2014; Raven, 2014). However, compensation may be limited by the available membrane area and/or energetic costs of maintaining high transporter densities (Vico *et al.*, 2011). Empirical support for the hypothesis that smaller guard cells are faster is mixed and has primarily come from step changes in light (Drake *et al.*, 2013; Elliott-Kingston *et al.*, 2016; McAusland *et al.*, 2016; Kardiman & Røsbild, 2018; Haworth *et al.*, 2023; Yoshiyama *et al.*, 2024; Brench *et al.*, 2026; Tang *et al.*, 2026). Meta-analysis of dozens of species suggests that stomatal size alone is weakly related to kinetics and modulated by guard cell shape (Woning *et al.*, 2026). The relationship between guard cell size and stomatal response kinetics to leaf water potential (Aasamaa *et al.*, 2001) and VPD (Kübarsepp *et al.*, 2020; Qie *et al.*, 2024) is less well studied, but similarly equivocal.

One factor that might complicate the relationship between size and speed is aperture. Stomatal aperture can vary from near 0 when guard cell turgor pressure is low and asymptotically approach g_{max} as guard cell turgor pressure approaches ∞ (Figure 1D). The relationship between guard cell turgor pressure and stomatal aperture is nonlinear (Franks & Farquhar, 2001) as cell walls become less elastic at greater turgor (Franks *et al.*, 2001). The nonlinear relationship between guard cell turgor

and pore aperture implies that if the rate of turgor change is constant ($\frac{dP}{dt} = C$), the rate of change in aperture, and hence stomatal conductance, will vary depending on the initial aperture (a_0) relative to the maximum aperture ($a_{\max}$). Hence, $\frac{da}{dt} = f(a_0/a_{\max})$. When initial aperture is high, g_{sw} will respond more slowly than when initial aperture is low. Since we generally do not observe individual stomatal aperture, relating this idea to leaf-level g_{sw} requires scaling by stomatal density, biophysical constants, and morphological constants (e.g. Sack & Buckley, 2016). We will use the term “fraction of anatomical maximum stomatal conductance”, symbolized as $f_{g_{\max}}$ (this is denoted as g_{ratio} by Xie *et al.* (2022)). This value should be proportional to the average aperture divided by its maximum aperture for any arbitrary stomatal density. Leaves operating at $f_{g_{\max}}$ closer to unity will, all else being equal, be slower to respond than leaves operating with a $f_{g_{\max}}$ close to zero, independent of stomatal density. Franks *et al.* (2012) hypothesized that selection on stomatal density would maintain typical operational stomatal conductance (g_{op}) relative to $g_{\max}$ at a sufficiently low value that g_{sw} would be sensitive to relatively small changes in guard cell turgor pressure. Consistent with this hypothesis, the $g_{op}:g_{\max}$ ratio is often near 0.25 (McElwain *et al.*, 2016; Murray *et al.*, 2020), well below $f_{g_{\max}} = 1$, and in a range where g_{sw} would be responsive to small changes in guard cell turgor pressure. There is likely variation within and among species and growth forms in the $g_{op}:g_{\max}$ ratio (Xie *et al.*, 2022; Ochoa *et al.*, 2024).

Putting these two hypotheses together, we predict that both guard cell size and $f_{g_{\max}}$ will influence stomatal kinetics, but possibly at different scales of biological organization. Guard cell size tends to vary less than stomatal density or aperture at many biological scales. For example, on a mature leaf, guard cell length is nearly constant even as cell volume and pore aperture change with turgor (Willmer & Fricker, 1996; Franks *et al.*, 2001). Compared to stomatal density, guard cell size can be less variable (Lammertsma *et al.*, 2011; Bucher *et al.*, 2016) and evolve more slowly between species (Muir *et al.*, 2023). If there is relatively little plastic or genetic variation among individuals within a species, then most of the variance in stomatal kinetics within species cannot be explained by variation in guard cell size. In contrast, $f_{g_{\max}}$ can vary immensely within and among individuals because stomatal aperture responds dynamically over the day and in response to environmental variation. Because guard cell size and aperture typically vary at different levels of biological organization, we predict that guard cell size likely explains more variation in stomatal kinetics among species, whereas $f_{g_{\max}}$ will explain more variation within species. Within species variation can arise from either genetic differences among individuals or environmental plasticity. We will also test if adaxial stomata respond faster than abaxial stomata (Haworth *et al.*, 2018; Wall *et al.*, 2022; Woning *et al.*, 2026) and, if so, whether this difference can be explained by variation in guard cell size and/or $f_{g_{\max}}$ between surfaces.

One challenge in determining whether traits covary at higher levels of organization is that statistical procedures to account for phylogenetic nonindependence can obscure ecologically relevant covariation caused by natural selection (Westoby *et al.*, 1995; Uyeda *et al.*, 2018). For example, in a meta-analysis of




Figure 1: **Conceptual overview of the two hypotheses linking stomatal anatomy to closure kinetics.**

(A) **a.** Stomatal conductance (g_{sw}) in response to a step increase in vapor pressure deficit (VPD) typically exhibits a transient wrong-way response (WWR), in which g_{sw} briefly rises, followed by the right-way response (RWR), in which g_{sw} declines to a lower steady state. The RWR is well described by a Weibull function (dashed line; Equation 1) with initial conductance g_i , final conductance g_f , and kinetic parameters τ and λ . (B) **b.** The time constant τ sets the overall speed of closure, and the lag-time λ determines whether the response rate is roughly constant ($\lambda \approx 1$; solid) or accelerates over time ($\lambda > 1$; dashed). (C) **c.** Smaller guard cells have a greater surface area to volume ratio (SA:V; schematic cross-sections shown at right) and are predicted to exchange osmolytes more quickly across the plasma membrane, yielding a faster change in guard cell turgor pressure and faster closure. (D) **d.** The relationship between guard cell turgor pressure (P_g) and g_{sw} is nonlinear and saturating because guard cell walls become less elastic at higher turgor (Franks & Farquhar, 2001; Franks *et al.*, 2001). Consequently, the same change in turgor pressure (ΔP_g , double-headed arrows) produces a larger change in g_{sw} when stomata are at low f_{gmax} (blue/orange; steep tangent slope) than at high f_{gmax} (orange/blue; shallow tangent slope), predicting slower closure when f_{gmax} is high.

stomatal kinetic responses to light, the explanatory power of traits like guard cell size and stomatal ratio declined in phylogenetic compared to nonphylogenetic regression (Woning *et al.*, 2026). This finding may be caused in large part because grasses have small stomata, high adaxial:abaxial stomatal ratio, and rapid stomatal kinetics. Are these associations a coincidence caused by shared ancestry or ecologically important covariation that helps explain how grasses adapt to their environment? Recent advances in phylogenetic comparative methods use hierarchical, multiresponse phylogenetic models (Muir *et al.*, 2023; Westoby *et al.*, 2023; Halliwell *et al.*, 2025) to tease apart “conservative trait correlations” that act at higher levels of organization (e.g. among species) from trait correlations that explain variation at lower levels of organization (e.g. among individuals within species and plastic responses). We use this approach to test whether guard cell size and f_{gmax} explain variation in stomatal kinetics at different levels of biological organization. Throughout, we use guard cell length (l_{gc}) as a proxy for size (Sack & Buckley, 2016).

The four main questions we address in this study are:

1. Are VPD kinetics adequately characterized by the same functional form in Equation 1 as light responses?
2. Are l_{gc} and f_{gmax} positively associated with τ ?
3. At what biological levels of organization do l_{gc} and f_{gmax} vary and influence τ ?
4. Are abaxial-only stomatal responses slower than whole leaf stomatal responses?

As noted above, we are opportunistically analyzing data from an experiment designed to answer different questions. Before analyzing the kinetic data, we planned to test whether 1) guard cell size would influence τ and 2) that τ would be greater on the abaxial-only leaves based on a recent meta-analysis of stomatal responses to light (Woning *et al.*, 2026). We made the decision to test for an association between f_{gmax} and τ after exploratory analyses revealed a potential association. Adding hypotheses after some results have been analyzed, also known as “Hypothesizing After Results Known” (HARKing), can lead to reports of spurious associations that are not repeatable in later studies (Kerr, 1998; Smaldino & McElreath, 2016). Therefore, our conclusions about the effect of f_{gmax} on stomatal kinetics should be interpreted with caution until they are replicated.

Materials and Methods

Populations and phylogeny

We grew 29 populations of wild tomato (Table S1), including representatives of all described species of *Solanum* sect. *Lycopersicon* and sect. *Lycopersicoides* (Peralta *et al.*, 2008), as described in Muir *et al.* (2025). These populations were selected because they encompass the breadth of climatic variation

in the wild tomato clade (Pease *et al.*, 2016). ~~In one case, we substituted a congeneric population of the species (*S. galapagense*) and added a population of *S. pennellii*.~~ Due to constraints on growth space and time, we spread out measurements over 61.1 weeks ~~from August 29, 2022 to October 31, 2023.~~ Replicates within population were evenly spread out over this period to prevent confounding of temporal variation in growth conditions with variation among populations. ~~To estimate phylogenetic effects~~ For phylogenetic comparative analyses, we used the RAXML whole-transcriptome concatenated phylogeny based on 2,745 100-kb genomic windows from Pease *et al.* (2016). Two of our populations were not in this tree. We used *S. galapagense* accession LA1044 in place of LA3909. *S. pennellii* accession LA0750 was added as sister to LA0716. The node separating LA0750 from LA0716 was placed ~~half-way between the~~ halfway along the branch to the next deepest node.

Plant growth conditions and light treatments

A thorough description of plant growth conditions can be found in Muir *et al.* (2025), therefore we summarize the most important information here. Seeds provided by the Tomato Genetics Resource Center germinated on moist paper in plastic boxes after soaking for 30-60 minutes in a 50% (volume per volume) solution of household bleach and water, followed by a thorough rinse. We transferred seedlings to cell-pack flats containing Pro-Mix BX potting mix (Premier Tech, Rivière-du-Loup, Quebec, Canada) once cotyledons fully emerged, typically within 1-2 weeks of sowing. We grew seeds and seedlings for both sun and shade treatments under the same environmental conditions (12:12 h, 24.3:21.7 °C, 49.6:58.4 RH day:night cycle). LED light provided PPFD = $267 \mu\text{mol m}^{-2} \text{s}^{-1}$ (Fluence RAZRx, Austin, Texas, USA) during the germination and seedling stages.

Immediately prior to starting growth light intensity treatments (sun and shade), we transplanted seedlings to 3.78 L plastic pots containing 60% Pro-Mix BX potting mix, 20% coral sand (Pro-Pak, Honolulu, Hawai'i, USA), and 20% cinders (Niu Nursery, Honolulu, Hawai'i, USA) with slow release NPK fertilizer following manufacturer instructions (Osmocote Smart-Release Plant Food Flower & Vegetable, The Scotts Company, Marysville, Ohio, USA). Percentage composition is on a volume basis. We watered to field capacity three times per week to prevent drought stress. Seedlings were randomly assigned in alternating order within population to the sun or shade treatment during transplanting. The average daytime PPFD was $761 \mu\text{mol m}^{-2} \text{s}^{-1}$ and $115 \mu\text{mol m}^{-2} \text{s}^{-1}$ for sun and shade treatments, respectively.

Humidity response curves

We selected a fully expanded, unshaded leaf at least six leaves above the cotyledons during early vegetative growth. This typically meant that plants had grown in light treatments for ≈ 4 weeks,

ensuring they had time to sense and respond developmentally to the light intensity of the treatment rather than the seedling conditions (Schoch *et al.*, 1980).

We measured stomatal conductance over time after a step change in humidity at a constant leaf temperature, which we refer to this as a humidity-response curve. We measured humidity-response curves using a portable infrared gas analyzer (LI-6800PF, LI-COR Biosciences, Lincoln, Nebraska, USA). ~~Light-acclimated~~ During gas exchange measurements, light-acclimated plants were placed under LEDs dimmed to match their light treatment ~~during gas exchange measurements~~. We collected four humidity-response curves per leaf, an amphistomatous (untreated) curve and a pseudohypostomatous (treated) curve at high light-intensity (PPFD = 2000 $\mu\text{mol m}^{-2} \text{s}^{-1}$; 97.8:2.24 red:blue) and low light-intensity (PPFD = 150 $\mu\text{mol m}^{-2} \text{s}^{-1}$; 87.0:13.0 red:blue). We always measured high light-intensity curves first because photosynthetic downregulation is faster than upregulation in these species. As a consequence, measurement light intensity is confounded with measurement order in our design: low light-intensity curves were always measured after high light-intensity curves on the same leaf, following a low-humidity closure event, so we cannot fully separate an effect of light intensity *per se* from order, recovery state, or prior VPD exposure (see Discussion). Pseudohypostomatous leaves are the same as the amphistomatous leaves except that gas exchange through the upper (adaxial) surface is blocked by a neutral density plastic (propafilm). For brevity, we hereafter refer to these as “pseudohypo” and “amphi” leaf types. Comparing amphi and pseudohypo leaf types enabled us to test whether stomatal kinetics are different for ab- and adaxial stomata on the same leaf. To compensate for reduced transmission, we increased incident PPFD for pseudohypo leaves by a factor 1/0.91, the inverse of the measured transmissivity of the propafilm. We also set the stomatal conductance ratio, for purposes of calculating boundary layer conductance, to 0 for pseudohypo leaves following manufacturer directions. To control for order effects, we alternated between starting with amphi or pseudohypo measurements. We made measurements over two days. On the first day, we measured high and low light-intensity curves for either amphi or pseudohypo leaves; on the second day, we measured high and low light-intensity curves on the other leaf type at approximately the same time of day.

In all humidity-response curves, we acclimated the focal leaf ambient CO_2 ($C_a = 415 \mu\text{mol mol}^{-1}$), $T_{\text{leaf}} = 25.0^\circ\text{C}$, high light (PPFD = 2000 $\mu\text{mol m}^{-2} \text{s}^{-1}$), and high relative humidity (RH = 70%; VPD $\approx 0.82 \text{ kPa}$) until g_{sw} reached its maximum. After that, we scrubbed water vapor from the incoming air using silica desiccant as quickly as possible, resulting in a water vapor concentration in the incoming air to be on average 0.36 mmol mol^{-1} . This treatment induced rapid stomatal closure, but it did not standardize chamber air humidity because ~~water vapor was introduced by leaf transpiration~~ leaf transpiration introduced water vapor. Consequently, the RH was systematically higher at high light intensity, in sun plants, and amphi leaf types (Table S2) because the g_{sw} was typically greater in these conditions. In all treatment conditions, the final RH (~~7.62%—20.4%~~ 5.37%—13.8%) and VPD

(~~2.5 kPa—2.92 kPa~~2.72 kPa — 3 kPa) elicited rapid stomatal response and is comparable to treatments used in similar experiments (Merilo *et al.*, 2014; Kübarsepp *et al.*, 2020; Hsu *et al.*, 2021; Qie *et al.*, 2024), but was not intended to mimic natural environmental variability. ~~We started logging data after~~Because the realized VPD stimulus varied among treatments, we checked whether this could
 265 confound treatment, f_{gmax} components, and l_{gc} effects on stomatal closure kinetics using curve-level VPD covariates (Pichaco *et al.*, 2024). We found that our conclusions were robust to this potential confound (Notes S1; Figs. S1–S4). Following the transient WWR, we started logging data once g_{sw} declined back to its pre-step value (g_i) and continued logging data until g_{sw} reached its nadir (Figure 1a). We then acclimated the leaf to low light (PPFD = 150 $\mu\text{mol m}^{-2} \text{s}^{-1}$) and RH = 70% before inducing
 270 stomatal closure with low RH and logging data again, on average 48.9 minutes after the start of the high-light curve.

Prior to analysis, we removed unreliable and unusable data points, removed points not at instrument steady state, removed the hysteretic portion of each ~~curves~~curve at low g_{sw} , removed outliers within each curve, removed replicates with no overlap between amphi and pseudohypo curves from the same
 275 leaf, and thinned redundant data points within each curve. The rationale and procedure for each cleaning step is described in Muir *et al.* (2025). Additionally, we excluded stomatal kinetic parameter estimates from ~~four~~five curves where τ was estimated to be greater than 1097 s, far exceeding the typical estimates (see Results). After these steps, the total number of humidity response curves analyzed is given in Table S3. The median number of curves per population per treatment was 9 and the median number of
 280 points per curve was 26. We extracted the posterior distribution for τ and λ from each fitted curve to calculate the parameter estimate (median) and uncertainty (standard deviation) for subsequent analyses.

~~Total number of humidity response curves analyzed in this study broken down by treatments.~~

Stomatal anatomical traits l_{gc} and f_{gmax}

We estimated the stomatal density and l_{gc} on ab- and adaxial surfaces from all leaves used for humidity
 285 response curves. ~~We used~~using standard methods described fully in Muir *et al.* (2025). We excluded three individuals with extreme outlying guard cell length values. Preliminary analysis of variation in guard cell length among growth treatments, leaf types, and accessions identified these outliers based on a residual value greater than 0.45, which visually appeared as a clear break in the residual distribution. When the leaflet used for the LI-6800 humidity-response curve did not have a matching leaflet-level
 290 anatomical measurement, we used the first available anatomical measurement from a different leaflet on the same leaf as a proxy; this fallback was used for 9 of 561 individual-by-leaf-type combinations (1.6%). For each surface we calculated the anatomical maximum stomatal conductance (g_{max}) at a reference leaf temperature of 25 °C following Sack & Buckley (2016) as:

$$g_{\max} = bmdl_{\text{gc}}/\sqrt{2}.$$

The biophysical and morphological constants b and m are:

$$b = \frac{D_{\text{wv}}}{v}, \text{ and}$$

$$m = \frac{\pi c^2}{j^{0.5}(4hj + \pi c)},$$

295

where D_{wv} is the diffusion coefficient of water vapor in air and v is by the kinematic viscosity of dry air. We assumed $D_{\text{wv}} = 2.49 \times 10^{-5} \text{ m}^2 \text{ s}^{-1}$ and $v = 2.24 \times 10^{-2} \text{ m}^3 \text{ mol}^{-1}$ (Monteith & Unsworth, 2013). For kidney-shaped guard cells like those in wild tomatoes, $c = h = j = 0.5$. ~~The is the $g_{\max}$ of the adaxial surface divided by the sum total $g_{\max}$ of both surfaces.~~ To estimate f_{gmax} , we divided the initial g_{sw} estimated from the parameter $g_{\text{I}}-g_{\text{L}}$ in Equation 1 from each humidity response curve by the anatomical $g_{\max}$ for that leaf. The estimated $g_{\text{I}}-g_{\text{L}}$ was very close to the maximum g_{sw} observed near the beginning of each curve (Figure S5), suggesting that the estimated $g_{\text{I}}-g_{\text{L}}$ was a good proxy for stomatal conductance at the time of the step change in humidity. Because g_{I} is estimated from the same nonlinear fit as τ and λ , we used a null simulation to confirm that this does not, by itself, induce a spurious $f_{\text{gmax}}-\tau$ association (Notes S2). We used the anatomical $g_{\max}$ for the surface(s) measured in each curve to calculate f_{gmax} . For amphi curves, we used the sum of adaxial and abaxial $g_{\max}$; for pseudohypo curves, we used only the abaxial $g_{\max}$. Although $g_{\max}$ calculated this way is a theoretical upper bound based on stomatal geometry rather than a directly measured quantity, it corresponds well to empirically measured maximum g_{sw} in experimentally controlled *Arabidopsis* (Dow *et al.*, 2014).

300
305
310

Estimating kinetic parameters

We estimated τ and λ for each response curve by fitting Equation 1 to a time course of g_{sw} values using Bayesian nonlinear regression. All models were fit using Bayesian HMC sampling in the probabilistic programming language *Stan* (Stan Development Team, 2026) using the *R* package **brms** version 2.23.0 (Bürkner, 2017). We used *CmdStan* version 2.38.0 and **cmdstanr** version 0.9.0 (Gabry *et al.*, 2025) to interface with *R* version 4.5.3-4.6.1 (R Core Team, 2026). We sampled ~~1000~~ 4,000 post-warmup draws from the posterior distribution ~~on a single chain~~ across four chains. We adjusted the number of sampling iterations and thinning interval for each curve so that the Gelman-Rubin convergence statistic ($\hat{R}$) (Gelman & Rubin, 1992) was less than 1.05 and the bulk effective sample size (ESS) was greater than 400 for all model parameters, and there were fewer than 10 divergent transitions. To answer

315
320

Question 1, we assessed model fit using Bayesian R^2 (Gelman *et al.*, 2019) implemented in **brms**.

Estimating effects of stomatal anatomy on kinetic parameters

We first used model selection to determine whether anatomy (l_{gc} and f_{gmax}) affected kinetic parameters (λ and τ , Question 2) and at what level of biological organization (Question 3). To infer the effect of f_{gmax} , we modeled effects of component traits $\log(q_i)$ and $\log(q_{max})$ directly because $\log(f_{gmax}) = \log(q_i) - \log(q_{max})$. If f_{gmax} *per se* affects kinetics then both $\log(q_i)$ and $\log(q_{max})$ should be in the selected model. Alternatively, q_i or q_{max} could effect kinetic parameters independent of f_{gmax} . From the selected model, we used parameter estimates and 95% confidence intervals to test the predictions of our hypotheses. We interpreted nonzero fixed effects of l_{gc} ~~and f_{gmax}~~ , q_i , and q_{max} on λ and τ as evidence that these anatomical traits influence variation in stomatal kinetics among individuals. We interpreted nonzero phylogenetic correlations among traits as evidence that these traits covary among populations. We predicted that l_{gc} and q_i should be positively associated with λ and τ , whereas the effect of q_{max} should be negative. Finally, we used ~~mediation-path~~ analysis to test whether treatment-associated plasticity in τ was statistically consistent with an indirect path through l_{gc} and/or f_{gmax} ~~contribute to plasticity in stomatal kinetics components~~. Table 1 summarizes the statistical criteria for determining whether and at what level of biological organization stomatal anatomical traits affected kinetic parameters. The subsections below describe specific procedures for deciding whether those are met.

Bayesian multiresponse phylogenetic models

Multiresponse models ~~enable us to~~ can disentangle whether traits are associated among populations, individuals, and/or treatments that induce plasticity. The response variables in our models are τ , λ , l_{gc} , ~~and f_{gmax}~~ , q_i , and q_{max} . All models included a subset of explanatory variables that were integral to the experimental design or necessary to account for statistical nonindependence. As explanatory variables, we included fixed effects of manipulative treatments: growth light intensity (sun and shade); measurement light intensity (high and low); and leaf type (amphi and pseudohypo). All models also included population as both a phylogenetically structured random effect and a non-phylogenetic random effect following Halliwell *et al.* (2025). Prior to fitting, the phylogenetic (co)variance matrix was rescaled as a correlation matrix (De Villemereuil & Nakagawa, 2014). For traits ~~measured multiple times estimated from multiple curves~~ within the same leaf (~~f_{gmax}~~ , λ , τ , q_i) we included a random effect of individual to account for repeated measures. We accounted for uncertainty in τ and λ using the standard deviation of the posterior distribution of each estimate (see ‘Humidity response curves’ for details). We modeled residual (co)variance, a combination of among-individual (co)variance and measurement error, as a multivariate Student t -distribution, which is more robust to extreme values than

Table 1: Criteria used to determine whether and at what level of biological organization stomatal anatomical traits affected kinetic parameters.

Level of organization	Criteria supporting an association between stomatal anatomy and kinetics
Among individuals	- Anatomical parameters as fixed effects improve model fit (lower LOOIC) - Anatomical effects on kinetics are in the predicted direction - Confidence intervals of anatomical effects do not overlap zero
Plasticity among individuals	Plasticity in anatomy is statistically consistent with an indirect path underlying plasticity in stomatal kinetics
Among populations	- Confidence intervals of phylogenetic correlation among traits do not overlap zero - Phylogenetic covariance is in the predicted direction

the multivariate normal distribution (Gelman *et al.*, 2014). To account for heteroskedasticity, bounded
 355 measurements, and to linearize relationships l_{gc} , λ , and τ all response variables were log-transformed
 prior to analysis whereas f_{gmax} was logit-transformed. All models were fit with Stan following on three
 chains using the same methods and convergence criteria described above for fitting Equation 1 to the
 humidity response curves.

Phylogenetic correlation among traits

360 We inferred that traits covary among populations if the 95% confidence intervals of the phylogenetic
 correlation between traits did not overlap zero and the correlation was in the predicted direction. For
 example, a positive phylogenetic correlation between l_{gc} and τ would be consistent with the prediction
 that species with larger guard cells have slower stomatal kinetics. Phylogenetic covariance specifically
 indicates long-term evolutionary covariation among traits because our model simultaneously accounts
 365 for trait associations among individuals, shared plastic responses, and residual covariance. To test
 for possible causation, we used the inferred phylogenetic trait covariance matrix to estimate partial
 correlations between anatomical and kinetic variables (Halliwell *et al.*, 2025). We interpret a significant
 partial correlation between variables as a plausible causal hypothesis.

Model selection and parameter estimation of individual-level effects

To infer whether individual-level variation in f_{gmax} and l_{gc} influence kinetic parameters, we used the leave-one-out cross-validation information criterion (LOOIC) to compare the fit of models using the *R* package **loo** version 2.9.2.10.0 (Vehtari *et al.*, 2017) to calculate LOOIC values. We compared models with all permutations of f_{gmax} and l_{gc} , g_i , and g_{max} influencing λ and τ . We considered all models within two ΔLOOIC standard errors of the minimum LOOIC (Table S4) to be in the set of plausible models to generate posterior predictions for hypothesis testing. Within each plausible model, we assessed whether an a fixed effect was significantly different from than zero based on the 95% confidence intervals. Among the set of plausible models, we selected the model with the highest proportion of significant focal parameters for further analysis. Focal parameters are the fixed and phylogenetic effects of l_{gc} , g_i , and g_{max} on λ and τ . This approach aids interpretability because it favors simpler models that exclude additional nonsignificant parameters. Because this choice is somewhat arbitrary and not based on rigorous statistical principles, we discuss alternative interpretations based on the other plausible models. We report Pareto $\hat{k}$, PSIS effective sample size, and Monte Carlo standard error diagnostics for the plausible models in Table S5. Nearly every curve had good Pareto $\hat{k}$ (≤ 0.7), but the minimum PSIS effective sample size per curve was sometimes low (as low as 94 draws for some models), so the PSIS-LOO estimates for a small number of individual curves, and by extension the model-level Monte Carlo standard error, should be interpreted with some caution.

Mediation-Path analysis to test for plastic effects

We used causal mediation path analysis (Pearl, 2022) to test whether treatment-associated plasticity in τ was mediated by plasticity in f_{gmax} or l_{gc} statistically consistent with an indirect path through plasticity in stomatal traits, in response to measurement light intensity, growth light intensity, and curve-type treatments. We focused on g_i because it was the only individual-level trait significantly associated with τ in the selected model (see Results). We do not treat this as a randomized-treatment causal mediation analysis because g_i is not randomized, is estimated from the same curve as τ , and could share unmeasured confounders (e.g., hydraulic status, measurement order, realized VPD trajectory) with τ , violating the sequential ignorability assumption required for a causal interpretation. We assessed the plausibility of this assumption directly with a sensitivity analysis (Notes S3). We inferred the direct effect of treatments on τ from model coefficients. We estimated the mediated indirect effect of treatments using as the product of the effect of treatment on mediator (f_{gmax} or l_{gc}) g_i and the effect of the mediator that trait on τ (Imai *et al.*, 2010). We determined significant direct and mediator indirect effects based on whether the 95% confidence intervals of the effects were different from than zero. We calculated effect sizes as the point estimate (median of the posterior) divided by the standard error (standard deviation of the posterior).

Variance decomposition and phylogenetic heritability

One explanation for why traits have effects at different levels of organization is that variance in those traits is concentrated at those different levels of organization. To evaluate this, we decomposed the variance in response variables (λ , τ , f_{gmax} , l_{gc} , g_{gc} , g_{max}) into phylogenetic, population (nonphylogenetic), and individual levels. The phylogenetic variance component quantifies trait evolution between populations congruent with the phylogenetic relationships, whereas the population (nonphylogenetic) component quantifies variance among populations independent of phylogenetic relationships (Lynch, 1991; De Villemereuil & Nakagawa, 2014). We estimated these components from the associated random-effect standard deviations for each response variable. The among-individual variance components quantify variance among individuals within the same population and treatment. We estimated the among-individual component as the sum of the individual random-effect and residual standard deviations. It was not possible to estimate individual random-effect in l_{gc} random-effects in l_{gc} or g_{max} with our study design because we only measured average guard cell length and g_{max} once per leaf. The among-individual component is biased upwards because it also includes measurement error, which cannot be separately estimated with our experimental design. The phylogenetic heritability is equivalent to the phylogenetic variance component and is estimated following Lynch (1991) and De Villemereuil & Nakagawa (2014) as:

$$h_{phy}^2 = \frac{\sigma_{phy}^2}{\sigma_{phy}^2 + \sigma_{pop}^2 + \sigma_{ind}^2},$$

where σ_{phy}^2 , σ_{pop}^2 , σ_{ind}^2 are the phylogenetic, population (nonphylogenetic), and among-individual variance components, respectively. Because phylogenetic heritability was estimated after accounting for fixed treatment effects, estimates obtained in controlled environments likely overestimate phylogenetic heritability in natural environments, where plasticity is a larger component of trait variation.

Effect of adaxial stomata on kinetics

If adaxial stomata respond faster than abaxial stomata (Question 4), we predicted that τ should be greater in pseudohypo leaves compared to amphi. We evaluated this prediction by estimating the fixed effect of leaf type (amphi vs pseudohypo) on τ in the multiresponse model. We did not estimate adaxial-only kinetics because it was not part of the original experimental design (Muir *et al.*, 2025).

Results

The time constant τ explains most variation in stomatal closure kinetics

Stomatal conductance (g_{sw}) decreased rapidly in response to a step change in humidity. The model in Equation 1 fit the data extremely well (Figure S6). The mean and minimum Bayesian R^2 across all curves was 0.9994 and 0.98020.9803, respectively. Consider a typical leaf in the experiment characterized by the median time constant (τ) and lag time (λ) among wild tomato ~~accessions~~ populations in all treatments. After humidity decreased ~~and~~, the transient WWR elapsed, ~~and g_{sw} returned to the initial value g_i~~ , it took on average 127-128 s for g_{sw} to decrease half-way (t_{50}) from its initial (~~g_i~~) to final (~~g_i~~) steady state value. In terms of the kinetic model parameters, most variation among leaves was due to difference in the time constant rather than the lag time. In the previous example, increasing λ from its median to its maximum estimated value among ~~accessions~~ populations increased the t_{50} by only 7.2 s ~~7.1 s~~. In contrast, increasing τ from its median to maximum value increased t_{50} by 88 s ~~86 s~~. Hence, we primarily focus on treatments and anatomical parameters affecting τ .

~~Variation in guard cell length and aperture~~ Variation in guard cell length and f_{gmax} components

Guard cell length (l_{gc}) differed among populations and varied to a lesser extent between growth light intensity treatments and leaf surfaces. In contrast, stomatal conductance as a fraction of maximum conductance (f_{gmax}) was most strongly determined by measurement light intensity. Because g_{max} is unaltered by measurement light intensity, this treatment effect was entirely caused by changes in g_i . Excluding treatment effects, phylogenetically structured differences among populations (i.e., phylogenetic heritability) accounted for 63.3% ~~61.9%~~ [95% CI: 21.4% ~~to 83.9%~~ 18.9% to 82.8%] of the variance in l_{gc} (Figure 2, Table S6). There was also developmental plasticity in l_{gc} . On average, l_{gc} increased by 12.1% [95% CI: 11.1% ~~to 13.0%~~ to 13.1%] in the sun treatment (Figure 3a; Table S7). The average l_{gc} of pseudohypo leaves was shorter by -5.8% [95% CI: -6.6% to -5.0%] ~~shorter than compared to~~ amphi leaves (Figure 3a; Table S7). This was not because of plasticity, as we measured amphi and pseudohypo curves on the same leaf. Rather, this was a composition effect. Adaxial stomata are larger than abaxial stomata in these species (Figure S7), so the average l_{gc} in pseudohypo leaves was lower than in amphi leaves because the former only included abaxial stomata.

The phylogenetic heritability of f_{gmax} ~~was low~~ components was low (g_i) to moderate (g_{max}), accounting for 16.9% [95% CI: 1.7% ~~to 40.1%~~ 1.1% to 37.9%] ~~and 59.1%~~ [95% CI: 34.4% to 75.5%] of the variance, respectively (Figure 2, Table S6). High measurement light intensity increased ~~f_{gmax}~~ g_i by an average of 0.85 logit units ~~113.2%~~ [95% CI: 0.82 ~~to 0.87~~ 107.7% to 118.5%] (Figure 3b; Table S7). ~~For example, if $f_{gmax} = 0.1$ in low light, the typical f_{gmax} for the same leaf acclimated to high light was 0.21.~~ Sun leaves had on average -0.28 logit units ~~41.6%~~ [95% CI: -0.35 ~~to -0.21~~ 34.4% to 49.5%] ~~lower f_{gmax}~~

greater $g_{\max}$ than shade-grown plants (Figure 3b; Table S7). The pseudohypo leaf type had on average 0.21 logit units -19.5% [95% CI: 0.18 to 0.24 -21.5% to -17.3%] higher $f_{\max}$ lower g_i than untreated, amphistomatous leaves (Figure 3b; Table S7). This indicates that stomatal aperture was more open in leaves where gas exchange through adaxial stomata was blocked is presumably because a large fraction of stomata were blocked by the pseudohypo treatment and increased aperture of the abaxial stomata did not fully compensate. This observation differs from wheat, where g_{sw} on one surface was not affected by blocking gas exchange in the other (Wall *et al.*, 2022).




Figure 2: **Traits vary at different levels of biological organization.** A preponderance of the variation in guard cell length (l_{gc}), anatomical maximum stomatal conductance ($g_{\max}$), and the time-constant time constant τ is phylogenetically structured among populations. In contrast, most of the variance in initial stomatal conductance as a fraction of anatomical maximum stomatal conductance ($f_{\max} g_i$) and the lag-time lag time (λ) is among individuals. Nonphylogenetic variation among populations was low for all traits.

Variation in stomatal closure kinetic parameters

Estimates of stomatal closure kinetic parameters for each curve are in Table S8 and estimates for each population are in Table S9. These will be deposited on Dryad upon publication. The time-constant The time constant τ exhibited moderate phylogenetic heritability, accounting for $42.239.8\%$ [95% CI: 18.6 to $61.218.7\%$ to 60.4%] of the variance, whereas λ varied more within populations (Figure 2, Table S6). Some light treatments also affected kinetic parameters. In sun plants, τ was on average 5.56 The τ in sun plants was -10.7% [95% CI: 1.05 to $-10.415.1\%$ to -6.06%] higher than in lower than shade plants after accounting for other explanatory variables (Figure 4, Table S7). Within the same leaf, increasing the measurement light intensity prior to the humidity step change from $150 \mu\text{mol m}^{-2} \text{s}^{-1}$ to $2000 \mu\text{mol m}^{-2} \text{s}^{-1}$ significantly increased τ by $17.98.6\%$ [95% CI: 11.6 to $24.81.4\%$ to 16.4%] (Figure 4, Table S7). The time-lag lag time λ responded to growth, but not measurement light intensity




Figure 3: Caption on following page.

Figure 3: (previous page) **Guard cell length (l_{gc}) and stomatal conductance as a fraction of maximum conductance (f_{gmax}) vary among populations and treatments.** In both panels, black points are mean trait value per population within a particular treatment. Gray lines connect values within populations across treatments. (a) Most of the variation in l_{gc} is among populations, but developmental plasticity to growth light intensity tends to increase l_{gc} in sun (right facet) compared to shade (left facet) grown plants. There is no plasticity to leaf type (amphi vs. pseudohypo), but the average l_{gc} is slightly lower in pseudohypo leaves because adaxial stomata tended to be larger than abaxial stomata on the same leaf. (b) Measurement light intensity has the most direct effect on f_{gmax} because initial stomatal aperture-conductance (g_i) increases under high light intensity. To a lesser extent, f_{gmax} varies among populations, is slightly lower in sun-grown plants, and slightly higher in pseudohypo leaf types.

480 after accounting for other explanatory variables (Figure 4, Table S7). In sun plants, λ was on average 12.23.43% [95% CI: 10.1 to 14.21.4% to 5.67%] greater than in shade plants. High measurement light had little effect on λ , with an estimated increase of 1.17change of -1.55% [95% CI: -1.1 to 3.54-4.35% to 1.39%].

Guard cell length and g_i influence stomatal kinetics

485 Guard cell length and $f_{gmax} g_i$, a component of f_{gmax} , affected stomatal closure kinetics in the directions predicted by our hypotheses, but they operated at different levels of biological organization. Among the models we tested, six were plausible in that their predictive accuracy was similar based on Δ LOOIC (Table S4). All models in the plausible set indicated among-individual level effects of $f_{gmax} g_i$ on τ and λ , as we discuss in more detail below. Models that included both individual and phylogenetic effects of l_{gc} had similar LOOIC values but we selected a model with the selected model retained only phylogenetic effects for two reasons because our selection criteria favored models that excluded nonsignificant parameters (Table S4). First, in models that included both individual and phylogenetic effects, neither the phylogenetic correlation nor the fixed effect regression coefficient were significantly different from zero (results not shown) than zero (Figure S8; Figure S9). These parameters are highly colinear-collinear in the posterior (Figure ??S10) indicating difficulty in estimating both terms simultaneously and “smearing” out their confidence intervals (McElreath, 2020). Hence, the model with only phylogenetic effects is easier to interpret. The second reason we favor this model is that it is This model is also consistent with the high phylogenetic heritability in both l_{gc} and τ (Figure 2). Visually, we observed positive relationships among population mean trait values (Figure 5) but little relationship between l_{gc} and τ within populations (results not shown). When comparing many models simultaneously, differences in LOOIC performance can occur by chance (McLachrie & Vehtari, 2024), so judgement calls among

490

500




Figure 4: **Stomatal kinetics parameters λ (lag time; top facets) and τ (time constant; lower facets) vary among wild tomato populations.** Stomatal conductance decreases faster (lower τ) in response to a step change in vapor pressure deficit (VPD) in leaves when measured under low light intensity. The pattern was consistent for both shade (left facets) and sun (right facets) grown plants. The pattern for λ was qualitatively similar to that for τ . Each point is the average parameter value for one accession in that treatment combination. The growth and measurement light intensity treatments are described in the Materials and Methods section.

models with similar performance can be necessary. The preponderance of evidence indicates that l_{gc} explains variation primarily among rather than within our populations. The results in the main text are based on estimates from the selected model. Individual-level fixed effects (Figure S8) and phylogenetic correlations (Figure S9) of l_{gc} , g_i , and g_{max} on τ and λ were qualitatively consistent across all plausible models.

~~Model comparison results. Models with a Δ LOOIC less than two standard errors (SE) from the minimum LOOIC are considered to have substantial support and are used for posterior predictions to test hypotheses about the effects of guard cell length (l_{gc}) and the stomatal conductance as a fraction of anatomical maximum stomatal conductance (f_{gmax}) on stomatal kinetic parameter λ (time lag) and τ (time constant). Unsupported models are in gray italic font. The model selected for subsequent inference is in bold. LOOIC stands for leave-one-out cross-validation information criterion.~~

~~Leaves with stomatal apertures closer to their maximum aperture~~ Leaves with greater g_i closed more slowly than those which were less open (Figure ??6; Table S7). For example, increasing f_{gmax} from 20% to 30% ~~g_i from 0.2 to 0.4 mol m⁻² s⁻¹~~ resulted in a ~~10.725.1%~~ [95% CI: ~~7.2 to 14.4~~17.6% to 32.6%] increase in τ for a typical leaf in the reference treatment set (shade, low, amphi). ~~The f_{gmax}~~ We found little evidence that g_{max} affected τ (Figure S11). Leaves with greater g_i also had a ~~significant, positive, individual-level effect on λ~~ (Table S7) significantly greater lag time, λ (Figure S12), but there was little effect of g_{max} (Figure S13).

Populations that tended to have greater l_{gc} also tended to have greater τ as evidenced by a positive phylogenetic correlation (Table S10, Figure 5). However, we found little evidence that variation in l_{gc} explained τ or λ within populations. There were no other significant phylogenetic correlations (Table S10). The only marginally significant partial correlation was between l_{gc} and τ ($\rho = 0.74$ $\rho = 0.71$ [95% CI: ~~-0.18 -0.17~~ to 0.98]), indicating a plausible causal effect of l_{gc} on τ among populations.

Plasticity in τ is an indirect effect of plasticity in g_i

We imposed three treatments that influenced stomatal kinetics: growth light intensity, measurement light intensity, and leaf type. All three treatments directly affected τ and had indirect effects mediated by $f_{gmax} g_i$ (Figure 7). We did not estimate effects mediated by l_{gc} because the selected model did not retain direct effects of l_{gc} on τ (Table S4). The ~~direct effect of the sun growth treatment~~ indirect effect of greater g_i in sun leaves increased τ by ~~5.58~~11.78% [95% CI: ~~1.05 to 10.44~~8.27% to 15.99%], but the indirect effect of lower f_{gmax} counteracted it by ~~-5.06%~~ 95% CI: ~~-7.30 to -3.18%~~ counteracting the direct negative effect. The direct effect of the pseudohypo treatment ~~decreased~~ increased τ slightly by ~~-2.64~~ 8.60% [95% CI: ~~-4.98 to -0.15~~5.62% to 11.56%], but the indirect effect mediated by greater f_{gmax} increased lower g_i changed τ by ~~4.02 -6.72%~~ [95% CI: ~~2.64 to 5.49 -8.64%~~ to -4.83%].




Figure 5: **Phylogenetic covariance between guard cell length (l_{gc} , x -axis) and the stomatal closure time-constant (τ , y -axis) indicates that populations with larger stomata tend to close more slowly in response to humidity.** Phylogenetic covariance between guard cell length (l_{gc} , x -axis) and the stomatal closure time constant (τ , y -axis) indicates that populations with larger stomata tend to close more slowly in response to humidity. The overall pattern was consistent across growth light intensity treatments (left and right facets), measurement light intensity treatments (top and bottom facets), and leaf types (point colors). The direction and magnitude of the ellipses are such that they should encompass 95% of the phylogenetically structured (co)variance in l_{gc} and τ on log-transformed scales. The location of ellipses are set to the median trait values in each treatment combination for visual comparison. Each point is the average parameter value for one ~~accession~~population in that treatment combination. The growth and measurement light intensity treatments are described in the Materials and Methods section.




Figure 6: **Individual-level variation in stomatal apertures influences stomatal closures kinetics.** Individual-level variation in initial stomatal conductance (g_i) influences stomatal closure kinetics. As ~~the stomatal conductance as a fraction of anatomical maximum stomatal conductance~~ g_i (f_{gmax} , x -axis) ~~increase~~increases, the stomatal closure ~~time-constant~~time constant (τ , y -axis) increases. The overall pattern was consistent across grow light intensity treatments (left and right facets), measurement light intensity treatments (top and bottom facets), and leaf types (point colors). Lines are estimated using linear regression along with 95% confidence ribbons. The growth and measurement light intensity treatments are described in the Materials and Methods section.

In contrast, direct and indirect effects of high measurement light intensity increased τ . The direct effect increased τ by ~~17.90~~8.56% [95% CI: ~~11.63 to 24.82~~1.40% to 16.40%] and the indirect effect mediated by ~~f_{gmax}~~ g_i increased τ by ~~17.31~~27.62% [95% CI: ~~11.34 to 23.48~~19.18% to 36.24%].

~~No evidence that adaxial stomata close faster than abaxial stomata~~ Adaxial stomata close faster than abaxial stomata, but the effect is masked by plasticity in g_i

~~The time constant~~ Although a simple comparison of τ was slightly greater in pseudohypo leaves compared to the paired amphi leaf between paired amphi and pseudohypo leaf types shows little net difference (results not shown), ~~which would suggest the selected model reveals a significant direct effect of leaf type on τ once g_i is accounted for: pseudohypo leaves, which have only abaxial stomata, closed more slowly than amphi leaves (Figure 7; Table S7). This direct effect is consistent with the hypothesis that adaxial stomata close faster than abaxial stomata. However, pseudohypo leaves also had greater f_{gmax} , which slows stomatal closure, as discussed in the previous section. Once, the indirect effect of f_{gmax} on τ is accounted for, lower g_i and lower g_i is itself associated with faster closure (Figure 6). This indirect path through g_i counteracts most of the direct effect of pseudohypo leaf type on τ disappears, indicating that the observed difference in τ between pseudohypo and amphi leaves is mediated by f_{gmax} (Figure 7). There is actually a tendency for τ to be slightly lower when only abaxial stomata are measured in pseudohypo leaves, as evidenced by the negative effect of this leaf type on τ (Table S7), which is why the raw comparison of paired leaves does not reveal much difference in kinetics between leaf types even though adaxial stomata appear to close faster once g_i is controlled for.~~

Discussion

The ability for stomata to adjust to fluctuating environmental conditions likely evolved in early land plants because of selection to optimize water use relative to carbon gain (Raven, 2002). Hydroactive control of stomatal aperture in angiosperms increases flexibility to modulate carbon gain and water loss depending on leaf water status (Buckley, 2019), perhaps enabling a competitive advantage in environments where light and evaporative demand fluctuate at high frequency (McAdam & Brodribb, 2012, 2015; Pichaco *et al.*, 2026). We now understand the steady-state response of stomatal conductance (g_{sw}) to global increases in vapor pressure deficit (VPD) better but there is greater uncertainty surrounding kinetic responses to VPD (Buckley *et al.*, 2023). Here we considered two nonmutually exclusive mechanistic hypotheses for how stomatal anatomy influences stomatal closure kinetics. First, smaller stomata may close faster because of greater surface area to volume ratio (Drake *et al.*, 2013). Second, guard cells under higher turgor pressure respond more slowly because of reduced cell wall elasticity (Franks *et al.*, 2012). Based on these hypotheses, we predicted that leaves with smaller guard cells and operating further




Figure 7: **Stomatal aperture as a fraction of anatomical maximum stomatal conductance (f_{gmax}) mediates plasticity in the time constant τ . Treatment-associated plasticity in the time constant τ is statistically consistent with an indirect path through initial stomatal conductance (g_i).** The three treatments are growth light intensity (top facet; shade vs. sun), leaf type (middle facet; amphi vs. pseudohypo), and measurement light intensity (bottom facet; low vs. high). Horizontal arrows at the bottom of each graph indicate the direct effect of the treatment level on τ , slanted arrows leading to or from $f_{gmax} \sim g_i$ indicate mediated-indirect effects of the treatment on τ through g_i . Line thickness is proportional to the effect size and line color indicates whether the effects are negative (red) or positive (blue). Parameter estimates of the effect and 95% confidence intervals are above each line. The selected model did not retain direct effects of l_{gc} on τ and hence this variable could not be considered as included in this analysis. This path analysis assumes sequential ignorability (no unmeasured confounder of the g_i - τ relationship); see Notes S3 for a mediatorsensitivity analysis of this assumption.

from their anatomical maximum g_{sw} (i.e., lower f_{gmax}) would have faster stomatal closure kinetics (lower τ). The parameter f_{gmax} is here defined as the initial stomatal conductance (g_i) divided by g_{max} .

570 Analysis of 2,124-125 time courses of stomatal closure from 29 populations in multiple environments demonstrates that both guard cell size and ~~stomatal aperture g_i , a component of f_{gmax}~~ influence kinetics at different levels of biological organization. This new understanding suggests unexplored explanations for the relationships between stomatal size, density, speed, and operational g_{sw} (g_{op}) among ~~land~~ plants/angiosperms.

575 The kinetics of stomatal closure in response to VPD closely match response to light, suggesting similar underlying mechanisms. The lag-time model proposed by Woning *et al.* (2026) for light responses also describes humidity responses extremely well following the passive wrong-way response (WWR). The fitted models explained on average 99.94% of the variation in g_{sw} during closure (Figure S6). The formal mathematical similarity of light and humidity response might indicate that certain factors, ~~like such as~~ guard cell size, influence stomatal kinetics responses in general. Alternatively, disparate underlying processes may be described by the same macroscale formalism. For example, humidity responses may be controlled by ABA accumulation or loss (McAdam *et al.*, 2016), whereas C_i -independent red-light responses can be influenced by accumulation or loss of photosynthetic products (Matthews *et al.*, 2020). These disparate processes may both have similar kinetic properties, but would not necessarily imply that
585 VPD and light responses are strongly correlated. Future research comparing the correlation between light, VPD, CO_2 , and other stomatal kinetic responses can determine the extent to which generic or specific factors determine stomatal opening and closing rates (Creese *et al.*, 2014; Kübarsepp *et al.*, 2020).

The positive association between l_{gc} and τ (Figure 5) is consistent with the hypothesis that smaller
590 cells respond faster because they have a greater surface area to volume ratio for osmolyte flux (Franks & Farquhar, 2007; Drake *et al.*, 2013; Lawson & Vialet-Chabrand, 2019; Woning *et al.*, 2026). ~~If correct, this~~ Among several of the same *Solanum* species, larger guard cells are also associated with slower opening kinetics in response to light (Yoshiyama *et al.*, 2024), suggesting a general mechanism. Specifically, this association implies that the maximum rate of stomatal closure is limited by membrane
595 surface area. We do not have the data to evaluate the hypothesis mechanistically, but this should be tested in the future. In addition to guard cell size, it will be useful to test whether the ratio of guard cell to adjacent epidermal size affects the strength of the WWR, which may be necessary to induce more rapid stomatal closure (Buckley *et al.*, 2003; Pichaco *et al.*, 2025). An alternative explanation is that the correlation between l_{gc} and τ is not causal. The fact that l_{gc} did not explain much of the variation in
600 τ among individuals within populations suggests that l_{gc} may not directly affect τ . However, the high phylogenetic heritability and, hence, low variability in l_{gc} within populations may also explain why we were unable to detect a within-population effect of l_{gc} on τ .

A potential consequence of greater stomatal aperture is that ~~they close the pore closes~~ more slowly when the environment changes. Much of the variation in τ among individuals within populations was explained by variation in $f_{g_{\max}} g_i$ (Figure ??6). This association is not simply an artifact of the magnitude of the difference between g_i and g_s , because τ captures the proportional rate of change, not the absolute change, in g_{sw} . This factor was positively associated with τ among individuals ~~in the same combination of within~~ light treatments, ~~but also mediated much of the plastic effects of these treatments on~~ and treatment-associated plasticity in τ was statistically consistent with an indirect path through g_i (Figure 7). ~~To our knowledge, this relationship has not been previously documented. However, the~~ The rate of stomatal opening in response to light is associated with g_{op} in a sample of rainforest trees (Kardiman & Ræbild, 2018), which may indicate this is a more general phenomenon.

The association between $f_{g_{\max}} g_i$ and τ could be explained by the hypothesis that guard cell wall elasticity declines at greater turgor pressure (Franks *et al.*, 2012), like a rubber band that is stretched under tension. However, if this were the case, we would also expect a negative association between $g_{\max}$ and τ because leaves with greater $g_{\max}$ could have the same g_i with a smaller stomatal aperture. However, this effect was weakly supported (Figure S11, Figure S13). The selected model did not include an individual-level effect of $g_{\max}$ on kinetic parameters. Although the second-best model included $g_{\max}$ and the effect was in the predicted direction (Table S4), the effect was not significantly different than 0 (Figure S8). The weak effect of $g_{\max}$ compared to g_i casts doubt on whether a nonlinear relationship between turgor pressure and pore aperture is the best explanation. An alternative, albeit not mutually exclusive explanation, is a nonlinear relationship between pore aperture and g_{sw} that weakens the association between these variables at high g_{sw} . Airspace resistance within the substomatal cavity and stomatal interference are plausible mechanisms that would have this effect (Ting & Loomis, 1965; Kaiser, 2009; Lehmann & Or, 2015), although it is not clear that they are sufficient to produce the observed relationship between g_i and τ in wild tomatoes. Our study is limited by the fact that we did not directly measure guard cell turgor or stomatal aperture, but future studies should use newly developed methods (Brodersen *et al.*, 2025) to directly evaluate the association between these traits.

The impact of $f_{g_{\max}} g_i$ was most pronounced when comparing measurement light intensity treatments. Leaves acclimated to high light intensity were on average ~~144~~113% more open than the same leaves at low light intensity in the reference treatment (shade, amphi). As a consequence, stomata closed more slowly in leaves acclimated to high light intensity, with a ~~17.3% greater time constant~~27.6% greater time constant on average in the reference treatment, based on parameter estimates from the selected model (Table S7). We did not assess whether $f_{g_{\max}} g_i$ also affects stomatal opening and this should be measured to further test this hypothesis. The implication is that natural selection should favor g_{op} values that not only optimize instantaneous carbon gain and water loss in the present environment, but also carbon gain and water loss integrated through time in a fluctuating environment. We predict

that rapid, temporally-uncorrelated fluctuating environments will favor leaves with greater $f_{\text{gmax}} g_{\text{op}}$ to slow kinetic responses and prevent overreacting to transitory changes that are likely to revert; slower, temporally-correlated changes will favor lower $f_{\text{gmax}} g_{\text{op}}$ to enable close tracking of the environment.

The ~~absence of a~~ comparison between measurement light intensity treatments should be interpreted cautiously, however, because it was confounded with measurement order: high light curves always preceded low light curves on the same leaf, following a prior low-humidity closure event (see Materials and Methods). We cannot rule out that some of the apparent effect of measurement light intensity on g_i and τ , and the corresponding indirect path through g_i in our mediation analysis (Figure 7), reflects order, incomplete recovery, or prior VPD exposure rather than measurement light intensity itself. This concern applies specifically to the measurement light comparison and not to growth light intensity or leaf type, which were randomized or alternated, respectively.

The direct effect of leaf type on τ ~~suggests that adaxial and abaxial stomata close at similar rates when operating at similar apertures. The association between species with more adaxial~~ indicates that adaxial stomata close faster than abaxial stomata after accounting for differences in g_i . This result is consistent with the broader association between greater adaxial stomatal density and faster stomatal closure across land plants (Haworth *et al.*, 2018; Woning *et al.*, 2026) ~~may not be caused by adaxial stomata responding faster per se. Instead, we hypothesize that selection may favor faster closure in leaves with adaxial stomata to prevent desiccation, causing increased abaxial stomatal speed as a byproduct. When there is a sudden increase in evaporative demand, such as from a sunfleck, open adaxial stomata could cause.~~ One plausible reason adaxial stomata may have evolved to close intrinsically faster is because adaxial stomata directly overlie the photosynthetically active palisade mesophyll ~~to desiccate (Buckley et al., 2017a). If stomata do not close quickly, this will lower water potential and failure to close quickly could rapidly desiccate this tissue and impair photosynthesis (Tezara et al., 1999; Buckley et al., 2017a).~~ Hypostomatous leaves ~~are less impacted~~ may experience less dehydration risk because the palisade ~~are~~ is hydraulically buffered from changing VPD ~~when stomata are confined to the abaxial surface~~ (Buckley *et al.*, 2017a). If traits, such as guard cell size, ~~that affect adaxial stomata kinetics similarly affect abaxial stomata, this could~~ that influence kinetics act similarly on both surfaces, this shared basis, together with the intrinsic speed difference between surfaces documented here, could jointly explain why leaves with greater stomatal ratio also have faster kinetics despite independent environmental responses on each surface (Mott, 2007; Wall *et al.*, 2022). ~~and species with a greater proportion of adaxial stomata tend to have faster overall stomatal kinetics.~~

Our findings also suggest a novel explanation for the well-documented negative relationship between stomatal size and density across land plants (recently reviewed in Liu *et al.* (2025)). Conventional explanations for this relationship focus on optimizing g_{op} and minimizing area allocated to stomata, but not on kinetics. Our results point to an additional, previously unrecognized constraint. Smaller

stomata do appear to close faster, consistent with the surface area to volume hypothesis, but this kinetic advantage is contingent on operating at a moderate or low f_{gmax} . If a leaf reduced guard cell size while maintaining the same g_{op} , the reduced g_{max} would push f_{gmax} higher, partially offsetting the speed benefit of smaller cells. The only way to simultaneously reduce guard cell size, maintain g_{op} , and keep f_{gmax} from rising is to increase stomatal density—raising g_{max} and thereby limiting f_{gmax} . This logic predicts a coupled evolution of small guard cells and high stomatal density, a pattern observed broadly across vascular plants (Franks & Beerling, 2009; de Boer *et al.*, 2016; Haworth *et al.*, 2023; Liu *et al.*, 2025). Although physical constraints on pore geometry, such as increased resistance to diffusion at very small apertures (Leuning, 1983; Hodgson *et al.*, 2010), likely impose a lower limit on guard cell size, selection to prevent operating at high f_{gmax} (Franks *et al.*, 2012) may also explain why optimal guard cell size varies.

Future research is needed to determine how stomatal kinetic responses measured in controlled conditions impact performance in nature. Our experimental protocol induced stomatal closure by abruptly switching incoming air to near-zero humidity, a perturbation far more severe than the gradual or partial VPD fluctuations leaves experience in the field. This approach is well suited for estimating the maximum rate of stomatal closure and comparing it among individuals and treatments, but it need not follow that the same anatomical and physiological factors that govern τ in our experiment also govern kinetic variation under more naturalistic conditions, where closure VPD may fluctuate more gradually. Future studies should also control the VPD stimulus throughout the entire response curve, rather than allowing it to vary as transpiration itself alters chamber humidity, as it did here (Table S2). It will also be important to consider whether the substomatal airspace remains saturated during the humidity step, because g_{sw} will be underestimated if the airspace is subsaturated (Cernusak *et al.*, 2018; Buckley & Sack, 2019; Wong *et al.*, 2022; Rockwell *et al.*, 2022). We hypothesize that ecological conditions which favor close tracking of fluctuating environments will select for leaves with traits like smaller guard cell size and low $f_{gmax} g_i$ that enable rapid kinetics.

Acknowledgements

Comments from two anonymous reviewers greatly improved the manuscript. Sam McKlin and Tom Buckley helped with protocol development. Justin Alter, Max Gatlin, Joana Kim, Jenna Matsuyama, Brandon Najarian, Dachuan Wang, and Kai Yasuda contributed to data collection. We used Copilot, ChatGPT, and Posit Assistant (Claude) to assist with coding and proofreading text.

Competing interests

None declared.

705 **Funding**

US National Science Foundation OIA-1929167 (C.D.M.)

Author contributions

Conceptualization: C.D.M.; Data acquisition: C.D.M., W.S.L.; Data analysis and visualization: C.D.M.; Writing – original draft: C.D.M.; Writing – review & editing: C.D.M., W.S.L.

710 **Data availability**

All data and code are available on GitHub (<https://github.com/cdmuir/solanum-kinetics>). Data will be archived on Dryad upon publication; code will be archived on Zenodo upon publication.

References

715 **Aasamaa K, Söber A. 2011a.** Stomatal sensitivities to changes in leaf water potential, air humidity, CO₂ concentration and light intensity, and the effect of abscisic acid on the sensitivities in six temperate deciduous tree species. *Environmental and Experimental Botany* **71**: 72–78.

Aasamaa K, Söber A. 2011b. Responses of stomatal conductance to simultaneous changes in two environmental factors. *Tree Physiology* **31**: 855–864.

720 **Aasamaa K, Söber A, Rahi M. 2001.** Leaf anatomical characteristics associated with shoot hydraulic conductance, stomatal conductance and stomatal sensitivity to changes of leaf water status in temperate deciduous trees. *Australian Journal of Plant Physiology* **28**: 765–774.

Bauer H, Ache P, Lautner S, Fromm J, Hartung W, Al-Rasheid KAS, Sonnewald S, Sonnewald U, Kneitz S, Lachmann N, et al. 2013. The Stomatal Response to Reduced Relative Humidity Requires Guard Cell-Autonomous ABA Synthesis. *Current Biology* **23**: 53–57.

725 **Brench RA, Wilson MJ, Thorne SJ, Fleming AJ, Gray JE. 2026.** Was the evolution of faster stomata driven by increased gas exchange rates rather than increasing water use efficiency? *New Phytologist* **249**: 2355–2371.

730 **Brodersen CR, Brodribb TJ, Hochberg U, Holbrook NM, McAdam SAM, Zailaa J, Huggett BA, Marmottant P. 2025.** In situ cavitation bubble manometry reveals a lack of light-activated guard cell turgor modulation in bryophytes. *Proceedings of the National Academy of Sciences* **122**: e2419887122.

Bucher SF, Auerswald K, Tautenhahn S, Geiger A, Otto J, Müller A, Römermann C. 2016. Inter-

and intraspecific variation in stomatal pore area index along elevational gradients and its relation to leaf functional traits. *Plant Ecology* **217**: 229–240.

735 **Buckley TN. 2019.** How do stomata respond to water status? *New Phytologist* **224**: 21–36.

Buckley TN, Frehner EH, Bailey BN. 2023. Kinetic factors of physiology and the dynamic light environment influence the economic landscape of short-term hydraulic risk. *New Phytologist* **238**: 529–548.

740 **Buckley TN, John GP, Scoffoni C, Sack L. 2017a.** The sites of evaporation within leaves. *Plant Physiology* **173**: 1763–1782.

Buckley TN, Mott KA, Farquhar GD. 2003. A hydromechanical and biochemical model of stomatal conductance. *Plant, Cell & Environment* **26**: 1767–1785.

Buckley TN, Sack L. 2019. The humidity inside leaves and why you should care: Implications of unsaturation of leaf intercellular airspaces. *American Journal of Botany* **106**: 618–621.

745 **Buckley TN, Sack L, Farquhar GD. 2017b.** Optimal plant water economy: Optimal plant water economy. *Plant, Cell & Environment* **40**: 881–896.

Buckley TN, Sack L, Gilbert ME. 2011. The Role of Bundle Sheath Extensions and Life Form in Stomatal Responses to Leaf Water Status. *Plant Physiology* **156**: 962–973.

750 **Bürkner P-C. 2017.** ~~{brms}~~brms: An R package for Bayesian multilevel models using Stan. *Journal of Statistical Software* **80**.

Cernusak LA, Ubierna N, Jenkins MW, Garrity SR, Rahn T, Powers HH, Hanson DT, Sevanto S, Wong SC, McDowell NG, et al. 2018. Unsaturation of vapour pressure inside leaves of two conifer species. *Scientific Reports* **8**: 7667.

755 **Cowan IR. 1972.** Oscillations in stomatal conductance and plant functioning associated with stomatal conductance: Observations and a model. *Planta* **106**: 185–219.

Cowan IR, Farquhar GD. 1977. Stomatal function in relation to leaf metabolism and environment. *Symposia of the Society for Experimental Biology* **31**: 471–505.

760 **Creese C, Oberbauer S, Rundel P, Sack L. 2014.** Are fern stomatal responses to different stimuli coordinated? Testing responses to light, vapor pressure deficit, and $\epsilon\alpha\text{CO}_2$ for diverse species grown under contrasting irradiances. *New Phytologist* **204**: 92–104.

Damour G, Simonneau T, Cochard H, Urban L. 2010. An overview of models of stomatal conductance at the leaf level: Models of stomatal conductance. *Plant, Cell & Environment* **33**: 1419–1438.

Darwin F. 1898. [Observations on stomata](#). *Proceedings of the Royal Society of London* **63**: 413–417.

765 ~~de Boer HJ, Price CA, Wagner-Cremer F, Dekker SC, Franks PJ, Veneklaas EJ. 2016. Optimal allocation of leaf epidermal area for gas exchange. *New Phytologist* **210**: 1219–1228.~~

De Villemereuil P, Nakagawa S. 2014. [General Quantitative Genetic Methods for Comparative Biology](#). In: Garamszegi LZ, ed. *Modern Phylogenetic Comparative Methods and Their Application in Evolutionary Biology*. Berlin, Heidelberg: Springer Berlin Heidelberg, 287–303.

770 Desai S, Stroock A. 2025. [Absciscic acid-mediated water stress regulation can mechanistically explain oscillations and water stress memory in stomatal conductance](#).

[Dow GJ, Bergmann DC, Berry JA. 2014. An integrated model of stomatal development and leaf physiology. *New Phytologist* **201**: 1218–1226.](#)

Drake PL, Froend RH, Franks PJ. 2013. [Smaller, faster stomata: Scaling of stomatal size, rate of response, and stomatal conductance](#). *Journal of Experimental Botany* **64**: 495–505.

775 Elliott-Kingston C, Haworth M, Yearsley JM, Batke SP, Lawson T, McElwain JC. 2016. [Does Size Matter? Atmospheric CO₂ May Be a Stronger Driver of Stomatal Closing Rate Than Stomatal Size in Taxa That Diversified under Low CO₂](#). *Frontiers in Plant Science* **7**.

780 ~~Franks PJ, Beerling DJ. 2009. Maximum leaf conductance driven by CO₂ effects on stomatal size and density over geologic time. *Proceedings of the National Academy of Sciences* **106**: 10343–10347.~~

Franks PJ, Buckley TN, Shope JC, Mott KA. 2001. [Guard cell volume and pressure measured concurrently by confocal microscopy and the cell pressure probe](#). *Plant Physiology* **125**: 1577–1584.

785 Franks PJ, Farquhar GD. 2001. [The Effect of Exogenous Absciscic Acid on Stomatal Development, Stomatal Mechanics, and Leaf Gas Exchange in *Tradescantia Virginiana*](#). *Plant Physiology* **125**: 935–942.

Franks PJ, Farquhar GD. 2007. [The Mechanical Diversity of Stomata and Its Significance in Gas-Exchange Control](#). *Plant Physiology* **143**: 78–87.

790 Franks PJ, Leitch IJ, Ruszala EM, Hetherington AM, Beerling DJ. 2012. [Physiological framework for adaptation of stomata to CO₂ from glacial to future concentrations](#). *Philosophical Transactions of the Royal Society B: Biological Sciences* **367**: 537–546.

Gabry J, Češnovar R, Johnson A, Bröder S. 2025. *Cmdstanr: R Interface to ‘CmdStan’*.

Gelman A, Carlin JB, Stern HS, Dunson DB, Vehtari A, Rubin DB. 2014. *Bayesian data analysis*.

Boca Raton London New York: CRC Press, Taylor & Francis Group.

795 **Gelman A, Goodrich B, Gabry J, Vehtari A. 2019.** [R-squared for Bayesian Regression Models](#). *The American Statistician* **73**: 307–309.

Gelman A, Rubin DB. 1992. Inference from iterative simulation using multiple sequences. *Statistical Science* **7**: 457–472.

Givnish TJ, Vermeij GJ. 1976. Sizes and shapes of liane leaves. *American Naturalist* **110**: 743–778.

800 **Grantz DA, Zeiger E. 1986.** [Stomatal Responses to Light and Leaf-Air Water Vapor Pressure Difference Show Similar Kinetics in Sugarcane and Soybean](#). *Plant Physiology* **81**: 865–868.

Grossiord C, Buckley TN, Cernusak LA, Novick KA, Poulter B, Siegwolf RTW, Sperry JS, McDowell NG. 2020. [Plant responses to rising vapor pressure deficit](#). *New Phytologist* **226**: 1550–1566.

Halliwel B, Holland BR, Yates LA. 2025. [Multi-response phylogenetic mixed models: Concepts and application](#). *Biological Reviews* **100**: 1294–1316.

805 **Haworth M, Marino G, Materassi A, Raschi A, Scutt CP, Centritto M. 2023.** [The functional significance of the stomatal size to density relationship: Interaction with atmospheric \[CO₂\] and role in plant physiological behaviour](#). *Science of The Total Environment* **863**: 160908.

810 **Haworth M, Scutt CP, Douthe C, Marino G, Gomes MTG, Loreto F, Flexas J, Centritto M. 2018.** [Allocation of the epidermis to stomata relates to stomatal physiological control: Stomatal factors involved in the evolutionary diversification of the angiosperms and development of amphistomaty](#). *Environmental and Experimental Botany* **151**: 55–63.

Hetherington AM, Woodward FI. 2003. [The role of stomata in sensing and driving environmental change](#). *Nature* **424**: 901–908.

815 ~~**Hodgson JG, Sharafi M, Jalili A, Díaz S, Montserrat-Martí G, Palmer C, Cerabolini B, Pierce S, Hamzehee B, Asri Y, et al. 2010.** [Stomatal vs. Genome size in angiosperms: The somatic tail wagging the genomic dog?](#) *Annals of Botany* **105**: 573–584.~~

Homann U, Thiel G. 2002. [The number of K⁺ channels in the plasma membrane of guard cell protoplasts changes in parallel with the surface area](#). *Proceedings of the National Academy of Sciences* **99**: 10215–10220.

820 **Hsu P-K, Takahashi Y, Merilo E, Costa A, Zhang L, Kernig K, Lee KH, Schroeder JI. 2021.** [Raf-like kinases and receptor-like \(pseudo\)kinase GHR1 are required for stomatal vapor pressure difference response](#). *Proceedings of the National Academy of Sciences* **118**: e2107280118.

Imai K, Keele L, Tingley D. 2010. A general approach to causal mediation analysis. *Psychological Methods* **15**: 309–334.

825 Ishida A, Yamamura Y, Hori Y. 1992. Roles of leaf water potential and soil-to-leaf hydraulic conductance in water use by understorey woody plants. *Ecological Research* **7**: 213–223.

Jalakas P, Takahashi Y, Waadt R, Schroeder JI, Merilo E. 2021. Molecular mechanisms of stomatal closure in response to rising vapour pressure deficit. *New Phytologist* **232**: 468–475.

830 Jezek M, Silva-Alvim FAL, Hills A, Donald N, Ishka MR, Shadbolt J, He B, Lawson T, Harper JF, Wang Y, *et al.* 2021. Guard cell endomembrane Ca²⁺-ATPases underpin a ‘carbon memory’ of photosynthetic assimilation that impacts on water-use efficiency. *Nature Plants* **7**: 1301–1313.

Kaiser H. 2009. The relation between stomatal aperture and gas exchange under consideration of pore geometry and diffusional resistance in the mesophyll. *Plant, Cell & Environment* **32**: 1091–1098.

835 Kardiman R, Ræbild A. 2018. Relationship between stomatal density, size and speed of opening in Sumatran rainforest species. *Tree Physiology* **38**: 696–705.

Kerr NL. 1998. HARKing: Hypothesizing After the Results are Known. *Personality and Social Psychology Review* **2**: 196–217.

Knapp AK, Smith WK. 1990. Stomatal and photosynthetic responses to variable sunlight. *Physiologia Plantarum* **78**: 160–165.

840 Kübarsepp L, Laanisto L, Niinemets Ü, Talts E, Tosens T. 2020. Are stomata in ferns and allies sluggish? Stomatal responses to CO₂, humidity and light and their scaling with size and density. *New Phytologist* **225**: 183–195.

845 Lammertsma EI, Boer HJD, Dekker SC, Dilcher DL, Lotter AF, Wagner-Cremer F. 2011. Global CO₂ rise leads to reduced maximum stomatal conductance in Florida vegetation. *Proceedings of the National Academy of Sciences* **108**: 4035–4040.

Lange OL, Lösch R, Schulze E-D, Kappen L. 1971. Responses of stomata to changes in humidity. *Planta* **100**: 76–86.

Lawson T, Blatt MR. 2014. Stomatal Size, Speed, and Responsiveness Impact on Photosynthesis and Water Use Efficiency. *Plant Physiology* **164**: 1556–1570.

850 Lawson T, Violet-Chabrand S. 2019. Speedy stomata, photosynthesis and plant water use efficiency. *New Phytologist* **221**: 93–98.

Lawson T, Von Caemmerer S, Baroli I. 2010. Photosynthesis and Stomatal Behaviour. In: Lüttge

UE, Beyschlag W, Büdel B, Francis D, eds. Progress in Botany 72. Berlin, Heidelberg: Springer Berlin Heidelberg, 265–304.

855

~~Leuning R, Lehmann P, Or D. 1983~~2015. Transport Effects of gases into leaves stomata clustering on leaf gas exchange. *Plant, Cell & Environment* 6: 181–194.

~~Liu C, Muir CD, Sack L, Li Y, Xu L, Li M, Zhang J, De Boer HJ, Han X, Yu G, et al. 2025.~~
860 ~~Bounds on stomatal size can explain sealing with stomatal density in forest plants.~~*New Phytologist*
~~248~~207: 2910–29261015–1025.

Lynch M. 1991. Methods for the analysis of comparative data in evolutionary biology. *Evolution* 45: 1065–1080.

Matthews JSA, Violet-Chabrand S, Lawson T. 2020. Role of blue and red light in stomatal dynamic behaviour(J Evans, Ed.). *Journal of Experimental Botany* 71: 2253–2269.

865 McAdam SAM, Brodribb TJ. 2012. Stomatal innovation and the rise of seed plants. *Ecology Letters* 15: 1–8.

McAdam SAM, Brodribb TJ. 2015. The Evolution of Mechanisms Driving the Stomatal Response to Vapor Pressure Deficit. *Plant Physiology* 167: 833–843.

870 McAdam SAM, Sussmilch FC, Brodribb TJ. 2016. Stomatal responses to vapour pressure deficit are regulated by high speed gene expression in angiosperms. *Plant, Cell & Environment* 39: 485–491.

McAusland L, Violet-Chabrand S, Davey P, Baker NR, Brendel O, Lawson T. 2016. Effects of kinetics of light-induced stomatal responses on photosynthesis and water-use efficiency. *New Phytologist* 211: 1209–1220.

875 McElreath R. 2020. *Statistical Rethinking: A Bayesian course with examples in R and Stan*. Boca Raton: Taylor and Francis, CRC Press.

McElwain JC, Yiotis C, Lawson T. 2016. Using modern plant trait relationships between observed and theoretical maximum stomatal conductance and vein density to examine patterns of plant macroevolution. *New Phytologist* 209: 94–103.

880 McLatchie Y, Vehtari A. 2024. Efficient estimation and correction of selection-induced bias with order statistics. *Statistics and Computing* 34: 132.

Medlyn BE, Duursma RA, Eamus D, Ellsworth DS, Prentice IC, Barton CVM, Crous KY, De Angelis P, Freeman M, Wingate L. 2011. Reconciling the optimal and empirical approaches to modelling stomatal conductance. *Global Change Biology* 17: 2134–2144.

- 885 **Merilo E, Jõesaar I, Brosché M, Kollist H. 2014.** To open or to close: Species-specific stomatal responses to simultaneously applied opposing environmental factors. *New Phytologist* **202**: 499–508.
- Monteith JL. 1995.** A reinterpretation of stomatal responses to humidity. *Plant, Cell & Environment* **18**: 357–364.
- Monteith JL, Unsworth MH. 2013.** *Principles of environmental physics: Plants, animals, and the atmosphere*. Amsterdam ; Boston: Elsevier/Academic Press.
- 890 ~~**Mott KA. 2007.** Leaf hydraulic conductivity and stomatal responses to humidity in amphistomatous leaves. *Plant, Cell & Environment* **30**: 1444–1449.~~
- Mott KA, Parkhurst DF. 1991.** Stomatal responses to humidity in air and helox. *Plant, Cell & Environment* **14**: 509–515.
- Muir CD, Conesa MÀ, Galmés J, Pathare VS, Rivera P, López Rodríguez R, Terrazas T, Xiong D. 2023.** How important are functional and developmental constraints on phenotypic evolution? An empirical test with the stomatal anatomy of flowering plants. *The American Naturalist* **201**: 794–812.
- Muir CD, Lim WS, Wang D. 2025.** Plasticity and adaptation to high light intensity amplify the advantage of amphistomatous leaves.
- 900 **Murray M, Soh WK, Yiotis C, Spicer RA, Lawson T, McElwain JC. 2020.** Consistent relationship between field-measured stomatal conductance and theoretical maximum stomatal conductance in C_3 woody angiosperms in four major biomes. *International Journal of Plant Sciences* **181**: 142–154.
- Ochoa ME, Henry C, John GP, Medeiros CD, Pan R, Scoffoni C, Buckley TN, Sack L. 2024.** Pinpointing the causal influences of stomatal anatomy and behavior on minimum, operational, and maximum leaf surface conductance. *Plant Physiology* **196**: 51–66.
- 905 **Papanatsiou M, Petersen J, Henderson L, Wang Y, Christie JM, Blatt MR. 2019.** Optogenetic manipulation of stomatal kinetics improves carbon assimilation, water use, and growth. *Science* **363**: 1456–1459.
- Pearl J. 2022.** *Causality: Models, reasoning, and inference*. Cambridge New York, NY Port Melbourne New Delhi Singapore: Cambridge University Press.
- 910 **Pease JB, Haak DC, Hahn MW, Moyle LC. 2016.** Phylogenomics reveals three sources of adaptive variation during a rapid radiation(D-Penny, Ed.). *PLOS Biology* **14**: e1002379.
- Peralta IE, Spooner DM, Knapp S. 2008.** Taxonomy of wild tomatoes and their relatives (*Solanum* sect. *Lycopersicoides*, sect. *Juglandifolia*, sect. *Lycopersicon*; Solanaceae). **84**.

- 915 [Pichaco J, Diaz-Espejo A, Rodriguez-Dominguez CM. 2025. Stomatal-To-Epidermal Cell Size Ratio and Physiological Coordination Influence Stomatal Response Speed to Leaf Water Disruption in Tomato Plants. *Physiologia Plantarum* **177**: e70543.](#)
- [Pichaco J, Manandhar A, McAdam SAM. 2024. Mechanical advantage makes stomatal opening speed a function of evaporative demand. *Plant Physiology* **195**: 370–377.](#)
- 920 [Pichaco J, Rodriguez-Dominguez CM, Diaz-Espejo A. 2026. Hydroactive stomata: A key innovation underpinning angiosperm ecological dominance. *Trends in Plant Science*: S1360138526002438.](#)
- 925 **Potkay A, Cabon A, Peters RL, Fonti P, Sapes G, Sala A, Stefanski A, Butler E, Bermudez R, Montgomery R, et al. 2025. Generalized Stomatal Optimization of Evolutionary Fitness Proxies for Predicting Plant Gas Exchange Under Drought, Heatwaves, and Elevated CO₂. *Global Change Biology* **31**: e70049.**
- [Prentice IC, Dong N, Gleason SM, Maire V, Wright IJ. 2014. Balancing the costs of carbon gain and water transport: Testing a new theoretical framework for plant functional ecology\(J-Penuelas, Ed.\)](#). *Ecology Letters* **17**: 82–91.
- 930 [Qie Y-D, Zhang Q-W, McAdam SAM, Cao K-F. 2024. Stomatal dynamics are regulated by leaf hydraulic traits and guard cell anatomy in nine true mangrove species. *Plant Diversity* **46**: 395–405.](#)
- R Core Team. 2026. *R: A Language and Environment for Statistical Computing*. Vienna, Austria: R Foundation for Statistical Computing.**
- [Raven JA. 2002. Selection pressures on stomatal evolution. *New Phytologist* **153**: 371–386.](#)
- [Raven JA. 2014. Speedy small stomata? *Journal of Experimental Botany* **65**: 1415–1424.](#)
- 935 [Rockwell FE, Holbrook NM, Jain P, Huber AE, Sen S, Stroock AD. 2022. Extreme undersaturation in the intercellular airspace of leaves: A failure of Gaastra or Ohm? *Annals of Botany* **130**: 301–316.](#)
- [Sack L, Buckley TN. 2016. The developmental basis of stomatal density and flux. *Plant Physiology* **171**: 2358–2363.](#)
- 940 [Schoch P-G, Zinsou C, Sibi M. 1980. Dependence of the stomatal index on environmental factors during stomatal differentiation in leaves of *Vigna Sinensis* L.: 1. Effect of light intensity. *Journal of Experimental Botany* **31**: 1211–1216.](#)
- [Schymanski SJ, Or D, Zwieniecki M. 2013. Stomatal control and leaf thermal and hydraulic capacitances under rapid environmental fluctuations\(W-Bauerle, Ed.\)](#). *PLoS ONE* **8**: e54231.

- 945 Smaldino PE, McElreath R. 2016. [The natural selection of bad science](#). *Royal Society Open Science* **3**: 160384.
- Sperry JS, Venturas MD, Anderegg WRL, Mencuccini M, Mackay DS, Wang Y, Love DM. 2017. [Predicting stomatal responses to the environment from the optimization of photosynthetic gain and hydraulic cost: A stomatal optimization model](#). *Plant, Cell & Environment* **40**: 816–830.
- Stan Development Team. 2026. *Stan Modeling Language Users Guide and Reference Manual*.
- 950 Tang X, Huang W, Yuan X, Liu X, Wan J. 2026. [Variation in Stomatal Traits and Photosynthetic Induction Between Subtropical Evergreen and Deciduous Broadleaved Tree Species](#). *Physiologia Plantarum* **178**: e70824.
- Tezara W, Mitchell VJ, Driscoll SD, Lawlor DW. 1999. [Water stress inhibits plant photosynthesis by decreasing coupling factor and ATP](#). *Nature* **401**: 914–917.
- 955 [Ting IP, Loomis WE. 1965. Further Studies Concerning Stomatal Diffusion. *Plant Physiology* **40**: 220–228.](#)
- Uyeda JC, Zenil-Ferguson R, Pennell MW. 2018. [Rethinking phylogenetic comparative methods](#) (N Matzke, Ed.). *Systematic Biology* **67**: 1091–1109.
- 960 Vehtari A, Gelman A, Gabry J. 2017. [Practical Bayesian model evaluation using leave-one-out cross-validation and WAIC](#). *Statistics and Computing* **27**: 1413–1432.
- Vico G, Manzoni S, Palmroth S, Katul G. 2011. [Effects of stomatal delays on the economics of leaf gas exchange under intermittent light regimes](#). *New Phytologist* **192**: 640–652.
- 965 Wall S, Violet-Chabrand S, Davey P, Van Rie J, Galle A, Cockram J, Lawson T. 2022. [Stomata on the abaxial and adaxial leaf surfaces contribute differently to leaf gas exchange and photosynthesis in wheat](#). *New Phytologist* **235**: 1743–1756.
- Westbrook AS, McAdam SAM. 2021. [Stomatal density and mechanics are critical for high productivity: Insights from amphibious ferns](#). *New Phytologist* **229**: 877–889.
- Westoby M, Leishman MR, Lord JM. 1995. [On Misinterpreting the ‘Phylogenetic Correction’](#). *The Journal of Ecology* **83**: 531.
- 970 Westoby M, Yates L, Holland B, Halliwell B. 2023. [Phylogenetically conservative trait correlation: Quantification and interpretation](#). *Journal of Ecology* **111**: 2105–2117.
- Willmer C, Fricker M. 1996. *Stomata*. Dordrecht: Springer Netherlands.

Wong SC, Canny MJ, Holloway-Phillips M, Stuart-Williams H, Cernusak LA, Márquez DA, Farquhar GD. 2022. Humidity gradients in the air spaces of leaves. *Nature Plants* 8: 971–978.

975 **Woning N, Al-Salman Y, Kaiser E, Berman SR, Brendel O, Cano FJ, Carpentier S, Centritto M, Drake PL, Durand M, *et al.* 2026. Revisiting the relationship between stomatal size and speed across species – a meta-analysis. *New Phytologist* 249: 2338–2354.**

Xie J, Wang Z, Li Y. 2022. Stomatal opening ratio mediates trait coordinating network adaptation to environmental gradients. *New Phytologist* 235: 907–922.

980 **Xie X, Wang Y, Williamson L, Holroyd GH, Tagliavia C, Murchie E, Theobald J, Knight MR, Davies WJ, Leyser HMO, *et al.* 2006. The Identification of Genes Involved in the Stomatal Response to Reduced Atmospheric Relative Humidity. *Current Biology* 16: 882–887.**

Yoshiyama Y, Wakabayashi Y, Mercer KL, Kawabata S, Kobayashi T, Tabuchi T, Yamori W. 2024. Natural genetic variation in dynamic photosynthesis is correlated with stomatal anatomical traits in diverse tomato species across geographical habitats (T Lawson, Ed.). *Journal of Experimental Botany* 75: 6762–6777.

985

Supporting Information

Supporting tables

Table S1: Accession information of *Solanum* populations used in this study. The species name, accession number, collection latitude, longitude, and elevation. TGRC: Tomato Genetics Resource Center; mas: meters above sea level.

Species	TGRC accession	Latitude	Longitude	Elevation (mas)
<i>S. arcanum</i>	LA2172	-6.008	-78.858	662
<i>S. cheesmaniae</i>	LA0429	-0.644	-90.329	800
<i>S. cheesmaniae</i>	LA3124	-0.804	-90.042	1
<i>S. chilense</i>	LA1782	-15.267	-74.633	1000
<i>S. chilense</i>	LA4117A	-22.907	-67.941	3540
<i>S. chmielewskii</i>	LA1028	-13.883	-73.017	3000
<i>S. chmielewskii</i>	LA1316	-13.400	-73.906	2920
<i>S. corneliomulleri</i>	LA0107	-13.117	-76.383	60
<i>S. corneliomulleri</i>	LA0444	-13.433	-76.133	100
<i>S. galapagense</i>	LA0436	-0.953	-90.978	40
<i>S. galapagense</i>	LA1044	-0.284	-90.548	0
<i>S. habrochaites</i>	LA0407	-2.181	-79.884	70
<i>S. habrochaites</i>	LA1777	-9.550	-77.700	3216
<i>S. huaylasense</i>	LA1358	-9.533	-77.967	750
<i>S. huaylasense</i>	LA1360	-9.546	-77.929	1490
<i>S. huaylasense</i>	LA1364	-10.133	-77.383	2920
<i>S. lycopersicoides</i>	LA2951	-19.317	-69.450	2200
<i>S. lycopersicoides</i>	LA4126	-19.287	-69.396	3120
<i>S. neorickii</i>	LA1322	-13.483	-72.442	2380
<i>S. neorickii</i>	LA2133	-3.400	-79.183	1980
<i>S. pennellii</i>	LA0716	-16.225	-73.617	50
<i>S. pennellii</i>	LA0750	-14.775	-75.034	550
<i>S. pennellii</i>	LA3778	-14.775	-75.034	616
<i>S. peruvianum</i>	LA2744	-18.550	-70.150	400
<i>S. peruvianum</i>	LA2964	-18.028	-70.835	75
<i>S. pimpinellifolium</i>	LA1269	-11.483	-77.075	400
<i>S. pimpinellifolium</i>	LA1589	-8.433	-78.817	30
<i>S. pimpinellifolium</i>	LA2933	-1.442	-80.562	375
<i>S. sitiens</i>	LA4116	-22.159	-68.782	2960

Table S2: Relative humidity (RH, %) and leaf vapor pressure deficit (VPD, kPa) during stomatal kinetics measurements systematically differed because of differences in stomatal conductance. Stomatal conductance tended to be greatest in sun-grown plants, measured at high light intensity, and untreated (amphistomatous). ~~The~~ For each treatment combination, we report the mean values across replicate curves in each treatment conditions are given; within each curve we calculated of the RH and VPD at the start (Initial) and end (Final) of the fitted interval, and of the median RH and VPD across time points before calculating within the mean among curves fitted interval (Median).

Growth light intensity	Measurement light intensity	Leaf type	RH			VPD		
			Initial	Median	Final	Initial	Median	Final
shade	low	pseudohypo	12	7.62	5.37	2.78	2.92	3.00
shade	low	amphi	13.8	8.25	5.68	2.72	2.91	2.99
shade	high	pseudohypo	18.4	13.3	9.98	2.58	2.74	2.85
shade	high	amphi	21.2	14.7	10.6	2.48	2.69	2.83
sun	low	pseudohypo	13.8	9.63	7.39	2.72	2.85	2.93
sun	low	amphi	16.7	11.1	8.23	2.62	2.81	2.91
sun	high	pseudohypo	24.3	16.8	11.6	2.36	2.62	2.79
sun	high	amphi	29.8	20.4	13.8	2.17	2.50	2.72

Table S3: Total number of humidity response curves analyzed in this study broken down by treatments.

Growth light intensity	Measurement light intensity				Total
	Low		High		
	amphi	pseudohypo	amphi	pseudohypo	
shade	275	275	270	266	1,086
sun	245	246	274	274	1,039
Total	520	521	544	540	2,125

Table S4: Model comparison results. Models with a ΔLOOIC less than two standard errors (SE) from the minimum LOOIC are considered to have substantial support and are used for posterior predictions to test hypotheses about the effects of guard cell length (l_{gc}) and the components (g_i, g_{max}) of stomatal conductance as a fraction of anatomical maximum stomatal conductance (f_{gmax}) on stomatal kinetic parameter λ (lag time) and τ (time constant). Unsupported models are in gray italic font. The model selected for subsequent inference is in bold. LOOIC stands for leave-one-out cross-validation information criterion.

ΔLOOIC	SE	λ			τ		
		l_{gc}	g_i	g_{max}	l_{gc}	g_i	g_{max}
0.00	0.00		✓			✓	
3.58	2.95	✓	✓	✓	✓	✓	✓
4.80	2.96		✓		✓	✓	
4.91	2.92		✓			✓	✓
4.92	2.90	✓	✓			✓	
5.20	2.72		✓	✓		✓	
5.91	2.91	✓	✓	✓		✓	✓
6.06	2.85	✓	✓	✓		✓	
6.21	2.85	✓	✓		✓	✓	
6.80	3.04		✓	✓	✓	✓	✓
7.70	2.96		✓	✓	✓	✓	
8.20	2.90		✓		✓	✓	✓
9.04	2.95	✓	✓		✓	✓	✓
9.05	2.85	✓	✓			✓	✓
9.91	2.98	✓	✓	✓	✓	✓	
10.76	2.88		✓	✓		✓	✓
38.52	13.42	✓	✓				
38.61	13.29	✓	✓				✓
39.19	13.38		✓	✓	✓		✓
39.20	13.37	✓	✓	✓			
39.34	13.43	✓	✓	✓	✓		✓
39.62	13.26		✓	✓			
39.93	13.34		✓	✓			✓
40.18	13.39	✓	✓	✓			✓
40.33	13.40	✓	✓		✓		✓
40.55	13.31	✓	✓		✓		
40.87	13.30		✓				
41.02	13.53		✓		✓		✓
41.40	13.34		✓	✓	✓		

(continued)

ΔLOOIC	SE	λ			τ		
		l_{gc}	g_i	g_{max}	l_{gc}	g_i	g_{max}
43.99	13.30	✓	✓	✓	✓		
44.44	13.42		✓				✓
45.04	13.23		✓		✓		
53.96	15.47			✓	✓	✓	
55.21	15.36			✓		✓	
56.20	15.41	✓		✓		✓	
57.58	15.46			✓	✓	✓	✓
57.85	15.37	✓			✓	✓	✓
57.94	15.52	✓			✓	✓	
58.12	15.40	✓		✓	✓	✓	✓
59.08	15.46	✓		✓		✓	✓
59.45	15.41					✓	
59.87	15.42	✓				✓	
60.11	15.43			✓		✓	✓
60.21	15.45	✓				✓	✓
60.63	15.41					✓	✓
61.45	15.37				✓	✓	
61.86	15.49				✓	✓	✓
63.84	15.41	✓		✓	✓	✓	
92.74	20.59						
93.78	20.53			✓			✓
94.61	20.53	✓		✓			
94.93	20.57			✓	✓		✓
95.18	20.58	✓		✓	✓		
96.21	20.59			✓			
96.69	20.51	✓					
96.75	20.54	✓		✓			✓
96.87	20.64			✓	✓		
97.11	20.50				✓		
97.98	20.56	✓					✓
98.01	20.53	✓			✓		✓
98.27	20.45	✓			✓		
99.69	20.68	✓		✓	✓		✓
100.41	20.64				✓		✓
101.31	20.43						✓

Table S5: Pareto $\hat{k}$, PSIS effective sample size (ESS), and Monte Carlo standard error (MCSE) diagnostics for the six plausible models identified by LOOIC (Table S4). The selected model is in bold. Good/bad/very bad refer to the conventional Pareto $\hat{k}$ thresholds of 0.7 and 1. MCSE(elpd_{loo}) is left blank (‘–’) for models with at least one curve with $\hat{k} \geq 1$, for which the MCSE estimate is undefined.

Model	ΔLOOIC	SE	n curves	Good $\hat{k}$ (%)	Bad $\hat{k}$ (%)	Very bad $\hat{k}$ (%)	Min. PSIS ESS	MCSE(elpd _{loo})
model 46	0.00	0.00	2125	99.95	0.05	0.00	155	0.99
model 01	3.58	2.95	2125	99.95	0.05	0.00	149	0.99
model 14	5.20	2.72	2125	100.00	0.00	0.00	137	1.00
model 30	4.92	2.90	2125	99.95	0.05	0.00	158	1.00
model 42	4.91	2.92	2125	100.00	0.00	0.00	94	1.00
model 44	4.80	2.96	2125	100.00	0.00	0.00	172	0.99

Table S6: The relative contribution of phylogenetic, among population (nonphylogenetic), and among-individual variation differs among stomatal traits. The phylogenetic component is equivalent to the phylogenetic heritability. The population component is the non-phylogenetic variance among populations. The among-individual variance component is the variation among individuals within a population after accounting for treatment effect. Estimates and 95% confidence intervals (CI) are estimated as the median and quantile intervals of the posterior distribution. l_{gc} : guard cell length; g_i : initial stomatal conductance; g_{max} : anatomical maximum stomatal conductance; λ : lag time; τ : time constant.

component	% variance	95% CI
$\log(l_{gc})$		
phylogenetic	61.9%	[18.9%, 82.8%]
population (nonphylogenetic)	7.5%	[0.6%, 36.2%]
among-individual	29.9%	[15.7%, 49.2%]
$\log(g_i)$		
phylogenetic	16.9%	[1.1%, 37.9%]
population (nonphylogenetic)	5.6%	[0.9%, 17.2%]
among-individual	77.0%	[57.7%, 88.0%]
$\log(g_{max})$		
phylogenetic	59.1%	[34.4%, 75.5%]
population (nonphylogenetic)	1.3%	[0.0%, 13.6%]
among-individual	38.8%	[23.4%, 55.7%]
$\log(\lambda)$		
phylogenetic	16.5%	[0.2%, 41.8%]
population (nonphylogenetic)	4.7%	[0.2%, 15.8%]
among-individual	78.3%	[56.3%, 90.3%]
$\log(\tau)$		
phylogenetic	39.8%	[18.7%, 60.4%]
population (nonphylogenetic)	1.1%	[0.0%, 11.2%]
among-individual	58.5%	[38.7%, 74.1%]

Table S7: Fixed effect parameter estimates and 95% confidence intervals (CIs) from the posterior distribution of the selected model. For each response variable, we estimated effects of growth light intensity (sun vs. shade), measurement light intensity (high vs. low), and leaf type (amphi vs. pseudohypo). The selected model potentially includes effects of l_{gc} , g_i , and g_{max} on τ and λ .

Parameter	Estimate [95% CI]
$\log(l_{gc})$	
intercept (shade, low light, amphi)	2.96 [2.80, 3.11]
effect of sun growth treatment on $\log(l_{gc})$	0.11 [0.11, 0.12]
effect of pseudohypo leaf type on $\log(l_{gc})$	-0.06 [-0.07, -0.05]
$\log(g_i)$	
intercept (shade, low light, amphi)	-1.71 [-1.93, -1.47]
effect of sun growth treatment on $\log(g_i)$	0.35 [0.30, 0.40]
effect of high measurement light intensity on $\log(g_i)$	0.76 [0.73, 0.78]
effect of pseudohypo leaf type on $\log(g_i)$	-0.22 [-0.24, -0.19]
$\log(g_{max})$	
intercept (shade, low light, amphi)	0.86 [0.55, 1.17]
effect of sun growth treatment on $\log(g_{max})$	0.61 [0.59, 0.63]
effect of pseudohypo leaf type on $\log(g_{max})$	-0.40 [-0.42, -0.38]
$\log(\lambda)$	
intercept (shade, low light, amphi)	0.39 [0.28, 0.48]
effect of sun growth treatment on $\log(\lambda)$	0.03 [0.01, 0.06]
effect of high measurement light intensity on $\log(\lambda)$	-0.02 [-0.04, 0.01]
effect of pseudohypo leaf type on $\log(\lambda)$	0.01 [-0.00, 0.03]
effect of loggi on $\log(\lambda)$	0.15 [0.11, 0.19]
$\log(\tau)$	
intercept (shade, low light, amphi)	5.58 [5.24, 5.91]
effect of sun growth treatment on $\log(\tau)$	-0.11 [-0.16, -0.06]
effect of high measurement light intensity on $\log(\tau)$	0.08 [0.01, 0.15]
effect of pseudohypo leaf type on $\log(\tau)$	0.08 [0.05, 0.11]
effect of loggi on $\log(\tau)$	0.32 [0.23, 0.41]

Table S8: Raw data file in CSV format containing estimates of stomatal anatomy and kinetic variables associated with each curve. These are raw data for fitting multiresponse models. A final version will be deposited on Dryad after acceptance. This version is available for reviewers.

Table file:

Download from:

<https://github.com/cdmuir/solanum-kinetics/raw/refs/heads/main/tables/tbl-estimates-curve.csv>

Table S9: Raw data file in CSV format containing estimates of stomatal anatomy and kinetic variables associated with each population based on multiresponse model predictions. A final version will be deposited on Dryad after acceptance. This version is available for reviewers.

Table file:

Download from:

<https://github.com/cdmuir/solanum-kinetics/raw/refs/heads/main/tables/tbl-estimates-accession.csv>

Table S10: Random effect parameter estimates and 95% confidence intervals (CIs) from the posterior distribution of the selected model. For each response variable, we estimated among-individual random effects from repeated measures of the same individual, where possible, random effects of population (nonphylogenetic), and phylogenetically structured random effects of population. We report the random effect standard deviations (SD). For population-level random effects, we also estimated correlations between traits.

Parameter	Estimate [95% CI]
<i>among-individual random effects</i>	
SD in $\log(\lambda)$	0.07 [0.07, 0.08]
SD in $\log(\tau)$	0.21 [0.19, 0.22]
SD in $\log(g_i)$	0.27 [0.25, 0.30]
<i>phylogenetic random effects</i>	
SD in $\log(\lambda)$	0.06 [0.01, 0.12]
SD in $\log(\tau)$	0.25 [0.16, 0.38]
SD in $\log(g_i)$	0.18 [0.05, 0.32]
SD in $\log(g_{\max})$	0.27 [0.18, 0.40]
SD in $\log(l_{gc})$	0.13 [0.06, 0.21]
correlation between in $\log(\lambda)$ and $\log(\tau)$	0.42 [-0.26, 0.84]
correlation between in $\log(\lambda)$ and $\log(g_i)$	-0.06 [-0.65, 0.64]
correlation between in $\log(\lambda)$ and $\log(g_{\max})$	-0.27 [-0.71, 0.45]
correlation between in $\log(\lambda)$ and $\log(l_{gc})$	0.09 [-0.53, 0.70]
correlation between in $\log(\tau)$ and $\log(g_i)$	0.13 [-0.49, 0.64]
correlation between in $\log(\tau)$ and $\log(g_{\max})$	-0.07 [-0.52, 0.37]
correlation between in $\log(\tau)$ and $\log(l_{gc})$	0.62 [0.14, 0.88]
correlation between in $\log(g_i)$ and $\log(g_{\max})$	0.38 [-0.28, 0.78]
correlation between in $\log(l_{gc})$ and $\log(g_i)$	0.41 [-0.27, 0.80]
correlation between in $\log(l_{gc})$ and $\log(g_{\max})$	-0.00 [-0.57, 0.45]
<i>population (nonphylogenetic) random effects</i>	
SD in $\log(\lambda)$	0.03 [0.01, 0.06]
SD in $\log(\tau)$	0.04 [0.00, 0.13]
SD in $\log(g_i)$	0.11 [0.05, 0.19]
SD in $\log(g_{\max})$	0.04 [0.00, 0.12]
SD in $\log(l_{gc})$	0.05 [0.02, 0.09]
correlation between in $\log(\lambda)$ and $\log(\tau)$	0.17 [-0.63, 0.79]
correlation between in $\log(\lambda)$ and $\log(g_i)$	-0.35 [-0.83, 0.37]
correlation between in $\log(\lambda)$ and $\log(g_{\max})$	-0.13 [-0.76, 0.71]
correlation between in $\log(\lambda)$ and $\log(l_{gc})$	-0.01 [-0.65, 0.66]
correlation between in $\log(\tau)$ and $\log(g_i)$	0.09 [-0.65, 0.74]
correlation between in $\log(\tau)$ and $\log(g_{\max})$	-0.06 [-0.78, 0.71]
correlation between in $\log(\tau)$ and $\log(l_{gc})$	0.24 [-0.64, 0.84]

(continued)

Parameter	Estimate [95% CI]
correlation between $\ln(g_i)$ and $\ln(g_{\max})$	-0.17 [-0.78, 0.56]
correlation between $\ln(l_{\text{gc}})$ and $\ln(g_i)$	0.44 [-0.18, 0.84]
correlation between $\ln(l_{\text{gc}})$ and $\ln(g_{\max})$	0.03 [-0.72, 0.71]

Supporting figures




Figure S1: Humidity-response curves and fitted lines. Pairwise relationships among curve-level VPD covariates. **a.** Median vs. final VPD during the Tomato Genetics Resource Center accession number, replicate letter, and species names fitted interval; **b.** median VPD vs. VPD saturation rate k ; **c.** final VPD vs. VPD saturation rate k . Each point is one humidity response curve, measurement light intensity (low or high), and colored by leaf type (amphi or pseudohypo). Points are raw data and lines are fitted curves (ordinary least-squares fits within leaf type). The Bayesian correlation coefficient, Bayes R^2 , VPD saturation rate is shown to the right rate constant of each curve (a saturating exponential fit to the VPD trajectory during the fitted interval (see Notes S1)).




Figure S2: The stomatal closure time constant τ (log scale) is strongly correlated with the inverse of the VPD saturation rate k^{-1} ($r = 0.77$ on log-log scale). This is consistent with k being driven largely by the same g_{sw} -decline signal that determines τ , rather than an independent property of the VPD stimulus (see Notes S1). Each point is one humidity response curve from the untreated (amphi) leaves. The blue line is ordinary least-squares regression fit.




Figure S3: The VPD saturation rate is driven by the sample airstream, not the reference airstream. **a.** VPD saturation rate vs. the saturation rate of W_s (sample cell water vapor concentration); the dashed line is the 1:1 line. **b.** VPD saturation rate vs. the saturation rate of W_r (reference cell water vapor concentration, upstream of the leaf). Each point is one humidity response curve; rate constants come from the same saturating-exponential model used for VPD.




Figure S4: **The g_i effect on τ and the phylogenetic correlation between l_{gc} and τ are robust to accounting for final VPD.** Posterior means and 95% credible intervals for the fixed effect of f_{gmax} on log τ (left) and the phylogenetic correlation between log l_{gc} and log τ (right), comparing the original model (grey) to a model refit with final VPD added as a covariate of τ and λ (pink; see Notes S1).




Figure S5: **The estimated (*x*-axis) and observed (*y*-axis) maximum stomatal conductances (g_{sw}) are nearly identical.** Each point is the empirical maximum g_{sw} plotted against the estimated g_{sw} for every humidity response curve. The orange line is the 1:1 line for reference. Both axes are on a log-scale.

Figure file:

Download from: <https://github.com/cdmuir/solanum-kinetics/raw/refs/heads/main/figures/rh-curves.pdf>

Figure S6: **Humidity-response curves and fitted lines.** The title provides the Tomato Genetics Resource Center accession number, replicate letter, and species names. The subtitle indicates the growth light intensity (sun or shade), measurement light intensity (low or high), and leaf type (amphi or pseudohypo). Points are raw data and lines are fitted curves. The Bayesian correlation coefficient, Bayes R^2 , is shown to the right of each curve.




Figure S7: **Adaxial (upper) stomata are consistently larger than abaxial (lower) stomata in both sun-grown (grey points) and shade-grown (black points) plants.** Each point was calculated by taking the median guard cell length within each biological replicate and then calculating the mean among replicates within each population and growth light intensity treatment. The dashed line is the 1:1 line for reference. All points above this line indicate larger values on the adaxial surface.




Figure S8: **Posterior estimates of phylogenetic correlations (x -axis) and fixed effects (y -axis) of guard cell length (l_{gc}) on the time-constant (τ) are colinear.** Each point is the estimate **Individual-level (fixed-effect) estimates of initial stomatal conductance (g_i , top facets), anatomical maximum stomatal conductance (g_{max} , middle facets), and guard cell length (l_{gc} , bottom facets) on the stomatal closure lag time (λ , left facets) and time constant (τ , right facets) are broadly consistent across all plausible models.** Points and horizontal bars are posterior medians and 95% credible intervals from **one-draw** each of the **posterior distribution for both parameters from six plausible models (rows; Table S4), ordered by Δ LOOIC; the selected model is marked with the lowest LOOIC an asterisk (*).** **The negative relationship—Black indicates estimates whose 95% credible interval excludes zero; grey indicates intervals that there is tradeoff in fitting the relationship between variables as a higher level phylogenetic effect versus a lower level, individual effect overlap zero. This results in more uncertainty and larger confidence intervals in each individual parameter.**




Figure S9: **Individual-level variation in stomatal apertures influences stomatal closures kinetics.** **As** **Phylogenetic correlations between initial stomatal conductance (g_i , top facets), anatomical maximum stomatal conductance (g_{max} , middle facets), and guard cell length (l_{gc} , bottom facets) on the stomatal closure lag time (λ , left facets) and time constant (τ , right facets) are broadly consistent across all plausible models.** Points and horizontal bars are posterior medians and 95% credible intervals of the **stomatal conductance as a fraction phylogenetic correlation from each of anatomical maximum stomatal conductance the six plausible models (f_{gmax} , x-axis rows; Table S4) increases**, ordered by Δ LOOIC; the **stomatal closure lag time selected model is marked with an asterisk (λ , y-axis*) increases.** **The overall pattern was consistent across growth light intensity treatments (left Black indicates estimates whose 95% credible interval excludes zero; grey indicates intervals that overlap zero.**




Figure S10: Posterior estimates of phylogenetic correlations (x -axis) and fixed effects (y -axis) of guard cell length (l_{gc}) on the time constant (τ) are collinear. Each point is the estimate from one draw of the posterior distribution for both parameters from the model with the lowest LOOIC. The negative relationship indicates that there is tradeoff in fitting the relationship between variables as a higher level phylogenetic effect versus a lower level, individual effect. This results in more uncertainty and larger confidence intervals in each individual parameter.




Figure S11: Individual-level variation in anatomical maximum stomatal conductance ($g_{\max}$) does not influence the stomatal closure time constant τ . Individual-level variation in $g_{\max}$ (x -axis) did not significantly covary with τ (y -axis). The overall pattern was consistent across growth light intensity treatments (left and right facets), measurement light intensity treatments (top and bottom facets), and leaf types (point colors). The growth and measurement light intensity treatments are described in the Materials and Methods section.




Figure S12: **Individual-level variation in initial stomatal conductance (g_i) influences stomatal closure kinetics.** As g_i (x -axis) increases, the stomatal closure lag time (λ , y -axis) increases. The overall pattern was consistent across growth light intensity treatments (left and right facets), measurement light intensity treatments (top and bottom facets), and leaf types (point colors). Lines are estimated using linear regression along with 95% confidence ribbons. The growth and measurement light intensity treatments are described in the Materials and Methods section.




Figure S13: Individual-level variation in anatomical maximum stomatal conductance ($g_{\max}$) does not influence the stomatal closure lag time λ . Individual-level variation in $g_{\max}$ (x -axis) did not significantly covary with λ (y -axis). The overall pattern was consistent across growth light intensity treatments (left and right facets), measurement light intensity treatments (top and bottom facets), and leaf types (point colors). The growth and measurement light intensity treatments are described in the Materials and Methods section.

Supporting notes

Notes S1: Effects of g_i and l_{gc} are robust to variation in VPD stimulus

Table S2 shows that the mean VPD realized during measurements differed among treatment combinations (2.72 kPa–3 kPa), because leaf transpiration itself humidifies the chamber and g_{sw} varied systematically with measurement light, growth light, and leaf type. Because VPD is the physiological driver of stomatal closure, we checked whether this variation in the realized stimulus could confound the effects of g_i and l_{gc} on the closure time constant, τ .

For every curve we extracted the leaf-to-air VPD trajectory from the logged humidity data and derived two curve-level covariates: final VPD and VPD saturation rate (k). Final VPD is the VPD at the end of the fitted interval; k is the rate constant of a saturating exponential, $VPD(t) = B + (R_0 - B) \exp(-kt)$, fit separately to each curve (median $R^2 = 0.99$, $n = 2,156$ curves; k excluded for 15 curves with a poor saturating fit or an extreme outlier rate estimate). We also considered median VPD. Median and final VPD were highly correlated ($r = 0.93$; Figure S1a), so we retained only final VPD.

We did not include the VPD saturation rate as a covariate of τ , because k and τ are themselves strongly correlated ($r = 0.77$; Figure S2) for a mechanistic rather than confounding reason: the VPD trajectory during the analyzed interval is driven mainly by declining leaf transpiration as stomata close, so k largely tracks the same g_{sw} -decline signal that determines τ . We tested this by fitting the same saturating-exponential model to the sample-cell water vapor concentration (W_s) and the reference-cell water vapor concentration (W_r , upstream of the leaf and largely reflecting the controlled incoming airstream). The VPD saturation rate closely tracked the W_s saturation rate ($r = 0.95$) but not the W_r saturation rate ($r = 0.02$) (Figure S3). Including k as a covariate of τ would therefore be close to circular.

To directly test whether the realized VPD stimulus confounds the g_i and l_{gc} effects on τ , we refit the selected model with final VPD added as a covariate of τ and λ (Figure S4). The individual-level effect of g_i on $\log(\tau)$ was slightly weaker (original: 0.32 [0.23, 0.41]; measurement-light-intensity-treatments 0.41; VPD-adjusted: 0.27 [0.18, 0.36]), as was the phylogenetic correlation between l_{gc} and τ (original: 0.59 [0.14, 0.88]; VPD-adjusted: 0.54 [-0.04, 0.88]). This indicates that variation in VPD slightly amplified the apparent effects of g_i and l_{gc} on stomatal closure kinetics, but does not qualitatively change the main conclusions. Final VPD itself had a negative fixed effect on $\log(\tau)$ (top -0.41 [-0.53, -0.27]), consistent with higher realized VPD accelerating stomatal closure.

We lack data on the rate of VPD change during the wrong-way response and the earliest part of the right-way response, because humidity logging did not begin until g_{sw} had already declined back to its pre-step value (Figure 1a); this limits our ability to evaluate whether the initial rate of VPD change, as opposed to the final realized VPD, influenced closure kinetics. We recommend that future studies of stomatal kinetics log chamber humidity continuously from the onset of the humidity step to more fully characterize the realized VPD stimulus and adjust the incoming airstream to keep VPD constant as stomata close.

Notes S2: The g_i - τ association is not an artifact of the curve-fitting procedure

The initial conductance g_i used to calculate $f_{g_{\max}}$ is estimated from the same nonlinear fit (Equation 1) that also estimates τ and λ for each curve. This raises the possibility that parameter covariance within the curve fit itself, rather than a true biological relationship, could generate an association between g_i and τ (and hence $f_{g_{\max}}$ and τ , since the denominator $g_{\max}$ is estimated independently from stomatal anatomy and cannot itself introduce a fitting artifact). To test this, we ran a null simulation in which any true association between g_i/g_c and the kinetic parameters was removed by construction. For every real curve i , we simulated a synthetic g_{sw} trajectory using curve i 's own measurement times, residual noise magnitude, and fitted τ and λ , but with g_i and g_c taken from a different, randomly chosen curve j . Because i and j were chosen independently at random, the synthetic curves retain realistic variability in both τ/λ and g_i/g_c , but there is no true association between g_i and the kinetic parameters. We re-fit the same Weibull functional form to each synthetic curve using nonlinear least squares (`nls()`), which closely approximated the full Bayesian pipeline used for the real data when compared directly on the real curves ($\text{cor}(\tau) = 0.99$, $\text{cor}(\lambda) = 1.00$), and computed the correlation between the *estimated* g_i and $\log(\tau)$. We repeated this 1,000 times to build a null distribution of this correlation.

The null-simulation correlations ranged from -0.06 to 0.09 (median -3.1×10^{-3}) across the 1,000 replicates, indistinguishable from zero (Figure S14). In contrast, the real, observed correlation between g_i and $\log(\tau)$ is 0.39 [0.35, 0.42], far outside the null range (empirical $p = 0.000$). This indicates that the curve-fitting procedure alone cannot explain the observed g_i - τ association, supporting our interpretation that it reflects a real relationship in the data rather than a statistical artifact of estimating g_i , τ , and λ from the same nonlinear fit.




Figure S14: **The curve-fitting procedure alone does not induce a g_i - τ association.** Null distribution of the correlation between estimated g_i and estimated τ across 1,000 simulated datasets in which the true g_i and g_τ are fixed at their real median values for every curve (so g_i has no true relationship with τ by construction), compared to the observed correlation in the real data (red line).

1050 **Notes S3: Sensitivity analysis of the g_i - τ path analysis to unmeasured confounding**

1055 The path analysis linking treatment, g_i , and τ assumes the causal structure in Figure S15: each treatment has a direct effect on τ and an indirect effect acting through g_i . This structure is only identified as causal under sequential ignorability, i.e., that no unmeasured confounder U (e.g., hydraulic status, measurement order, realized VPD trajectory) affects both g_i and τ . If such a U exists (dashed arrows), the correlation it induces between the g_i and τ equations' residuals is exactly the sensitivity parameter ρ examined below.




Figure S15: **Assumed causal structure of the g_i - τ path analysis.** Treatment has a direct effect on τ (bottom arrow) and an indirect effect acting through g_i (upper path). An unmeasured confounder U (dashed arrows) affecting both g_i and τ – e.g., hydraulic status, measurement order, or realized VPD trajectory – would violate the sequential ignorability assumption required to interpret the indirect path causally.

The path analysis linking g_i to τ (see Materials and Methods, “Path analysis to test for plastic effects”) assumes sequential ignorability: that there is no unmeasured confounder of the g_i - τ relationship after conditioning on treatment and the random effects already in the model. Because g_i is not randomized,

is estimated from the same curve as τ , and could share unmeasured confounders (e.g., hydraulic status, measurement order, realized VPD trajectory) with τ , we assessed this assumption directly with a sensitivity analysis following the general framework of Imai *et al.* (point-colors)2010), parameterized by $\rho \equiv \text{Cor}(e_{g_i}, e_\tau)$, the residual correlation between the g_i and τ equations. Under a violation of sequential ignorability with residual correlation ρ , the bias-adjusted mediator-on-outcome coefficient is $\beta_2(\rho) = \hat{\beta}_2 - \rho \cdot \sigma_\tau / \sigma_{g_i}$, so the indirect effect of a treatment acting through g_i is $\gamma_1 \cdot \beta_2(\rho)$ (where γ_1 is the effect of treatment on g_i), and the “breakdown point” ρ^* at which this indirect effect is exactly zero is $\rho^* = \hat{\beta}_2 \cdot \sigma_{g_i} / \sigma_\tau$. Lines are estimated using linear regression along with 95% confidence ribbons

Our selected model already estimates a free residual correlation between the g_i and τ equations, because we fit all five responses ($l_{gc}, g_i, g_{max}, \tau, \lambda$) jointly. Because the g_i equation’s predictors are a strict subset of τ ’s, jointly estimating this correlation does not change the point estimate of $\hat{\beta}_2$ relative to a model that assumes $\rho = 0$ (a standard result for seemingly unrelated regression models), so we can treat $\hat{\beta}_2$ as the “naive” coefficient in the sensitivity framework above, and the model’s own estimated residual correlation as a data-driven estimate of ρ . The growth-

We found $\rho_{\text{estimated}} = 0.06 [-0.07, 0.18]$ and measurement-light intensity treatments are described in the Materials and Methods section. $\rho^* = 0.40 [0.29, 0.51]$ (Figure S16). These posterior distributions do not substantially overlap and the posterior probability that the estimated residual correlation is smaller in magnitude than the breakdown point is nearly 100%. In other words, the plastic mediation effects of g_i on τ are probably not explained by unmeasured confounders.




Figure S16: **Sensitivity of the indirect effect of each treatment on $\log \tau$, acting through g_i , to an assumed residual correlation ρ between the g_i and τ equations.** The vertical dotted line is the model's own estimated residual correlation ($\rho_{\text{estimated}}$, shaded band = 95% credible interval); the vertical solid line is the breakdown point ρ^* at which the indirect effect is zero.