

Anderson Higgs Mechanism and Meissner Effect

Yiqiu Han

5/07/2019

Spontaneous Symmetry Breaking




Figure: Mexican Hat potential

Spontaneous Symmetry Breaking

If under a symmetry transformation, the potential of a system is still invariant, but the ground state transforms to another ground state, this process of selecting a ground state from a degenerate set of ground states leads to a "Spontaneous symmetry breakdown".

Goldstone Theorem

Each generator of a spontaneous broken symmetry corresponds to a massless and spinless field, called the Goldstone boson.

Goldstone boson in superfluid

- Lagrangian of the superfluid:

$$\mathcal{L} = \phi^*(\vec{r}, \tau) \left(\partial_\tau - \frac{\nabla^2}{2m} - \mu \right) \phi(\vec{r}, \tau) + \frac{g}{2} \phi^*(\vec{r}, \tau) \phi^*(\vec{r}, \tau) \phi(\vec{r}, \tau) \phi(\vec{r}, \tau)$$

$$V = -\mu |\phi|^2 + \frac{g}{2} |\phi|^4$$

- $\left. \frac{\partial V}{\partial \phi} \right|_{\phi_0} = 0,$

$$\phi_0 = \begin{cases} 0 & \text{if } \mu < 0, \\ \sqrt{\frac{\mu}{g}} e^{i\theta} = \sqrt{\rho} e^{i\theta} & \text{if } \mu > 0. \end{cases}$$

- The Lagrangian is invariant under the $U(1)$ symmetry transformation $\phi \rightarrow \phi e^{i\theta}$. However, when $\mu > 0$, $U(1)$ symmetry is spontaneously broken.

Goldstone boson in superfluid

- Lagrangian of the superfluid:

$$\mathcal{L} = \phi^*(\vec{r}, \tau) \left(\partial_\tau - \frac{\nabla^2}{2m} - \mu \right) \phi(\vec{r}, \tau) + \frac{g}{2} \phi^*(\vec{r}, \tau) \phi^*(\vec{r}, \tau) \phi(\vec{r}, \tau) \phi(\vec{r}, \tau)$$

$$V = -\mu |\phi|^2 + \frac{g}{2} |\phi|^4$$

- $\left. \frac{\partial V}{\partial \phi} \right|_{\phi_0} = 0,$

$$\phi_0 = \begin{cases} 0 & \text{if } \mu < 0, \\ \sqrt{\frac{\mu}{g}} e^{i\theta} = \sqrt{\rho} e^{i\theta} & \text{if } \mu > 0. \end{cases}$$

- The Lagrangian is invariant under the $U(1)$ symmetry transformation $\phi \rightarrow \phi e^{i\theta}$. However, when $\mu > 0$, $U(1)$ symmetry is spontaneously broken.

Goldstone boson in superfluid

- Lagrangian of the superfluid:

$$\mathcal{L} = \phi^*(\vec{r}, \tau) \left(\partial_\tau - \frac{\nabla^2}{2m} - \mu \right) \phi(\vec{r}, \tau) + \frac{g}{2} \phi^*(\vec{r}, \tau) \phi^*(\vec{r}, \tau) \phi(\vec{r}, \tau) \phi(\vec{r}, \tau)$$

$$V = -\mu |\phi|^2 + \frac{g}{2} |\phi|^4$$

- $\left. \frac{\partial V}{\partial \phi} \right|_{\phi_0} = 0,$

$$\phi_0 = \begin{cases} 0 & \text{if } \mu < 0, \\ \sqrt{\frac{\mu}{g}} e^{i\theta} = \sqrt{\rho} e^{i\theta} & \text{if } \mu > 0. \end{cases}$$

- The Lagrangian is invariant under the $U(1)$ symmetry transformation $\phi \rightarrow \phi e^{i\theta}$. However, when $\mu > 0$, $U(1)$ symmetry is spontaneously broken.

Goldstone boson in superfluid

- Lagrangian of the superfluid:

$$\mathcal{L} = \phi^*(\vec{r}, \tau) \left(\partial_\tau - \frac{\nabla^2}{2m} - \mu \right) \phi(\vec{r}, \tau) + \frac{g}{2} \phi^*(\vec{r}, \tau) \phi^*(\vec{r}, \tau) \phi(\vec{r}, \tau) \phi(\vec{r}, \tau)$$

$$V = -\mu |\phi|^2 + \frac{g}{2} |\phi|^4$$

- $\left. \frac{\partial V}{\partial \phi} \right|_{\phi_0} = 0,$

$$\phi_0 = \begin{cases} 0 & \text{if } \mu < 0, \\ \sqrt{\frac{\mu}{g}} e^{i\theta} = \sqrt{\rho} e^{i\theta} & \text{if } \mu > 0. \end{cases}$$

- The Lagrangian is invariant under the $U(1)$ symmetry transformation $\phi \rightarrow \phi e^{i\theta}$. However, when $\mu > 0$, $U(1)$ symmetry is spontaneously broken.

Goldstone boson in superfluid

- Now we choose a ground state $\phi_0 = \sqrt{\rho_0}$ and add a fluctuation around ϕ_0 :

$$\phi(\vec{r}, \tau) = \sqrt{\rho_0 + \delta\rho(\vec{r}, \tau)} e^{i\theta(\vec{r}, \tau)}, \delta\rho \ll 1.$$
- $\mathcal{L} \rightarrow \sqrt{\rho_0 + \delta\rho} e^{-i\theta} (\partial_\tau - \frac{\nabla^2}{2m} - \mu) \sqrt{\rho_0 + \delta\rho} e^{i\theta} + \frac{g}{2}(\rho_0 + \delta\rho)^2.$
 Substitute into the partition function,

$$\mathcal{Z} = \int \mathcal{D}(\delta\rho, \theta) e^{-\int d\tau \int d\vec{r} \mathcal{L}}$$

Integrate out $\delta\rho$, we get the effective Lagrangian,

$$\mathcal{L}_{\text{eff}} = \frac{1}{2g} (\partial_\tau \theta)^2 + \frac{\rho_0}{2m} (\nabla \theta)^2$$

- The $U(1)$ symmetry breaking generates a massless θ mode, it is our Goldstone boson in superfluid, phonon.

Goldstone boson in superfluid

- Now we choose a ground state $\phi_0 = \sqrt{\rho_0}$ and add a fluctuation around ϕ_0 :

$$\phi(\vec{r}, \tau) = \sqrt{\rho_0 + \delta\rho(\vec{r}, \tau)} e^{i\theta(\vec{r}, \tau)}, \delta\rho \ll 1.$$
- $\mathcal{L} \rightarrow \sqrt{\rho_0 + \delta\rho} e^{-i\theta} (\partial_\tau - \frac{\nabla^2}{2m} - \mu) \sqrt{\rho_0 + \delta\rho} e^{i\theta} + \frac{g}{2}(\rho_0 + \delta\rho)^2.$
 Substitute into the partition function,

$$\mathcal{Z} = \int \mathcal{D}(\delta\rho, \theta) e^{-\int d\tau \int d\vec{r} \mathcal{L}}$$

Integrate out $\delta\rho$, we get the effective Lagrangian,

$$\mathcal{L}_{\text{eff}} = \frac{1}{2g} (\partial_\tau \theta)^2 + \frac{\rho_0}{2m} (\nabla \theta)^2$$

- The $U(1)$ symmetry breaking generates a massless θ mode, it is our Goldstone boson in superfluid, phonon.

Goldstone boson in superfluid

- Now we choose a ground state $\phi_0 = \sqrt{\rho_0}$ and add a fluctuation around ϕ_0 :

$$\phi(\vec{r}, \tau) = \sqrt{\rho_0 + \delta\rho(\vec{r}, \tau)} e^{i\theta(\vec{r}, \tau)}, \delta\rho \ll 1.$$
- $\mathcal{L} \rightarrow \sqrt{\rho_0 + \delta\rho} e^{-i\theta} (\partial_\tau - \frac{\nabla^2}{2m} - \mu) \sqrt{\rho_0 + \delta\rho} e^{i\theta} + \frac{g}{2}(\rho_0 + \delta\rho)^2.$
 Substitute into the partition function,

$$\mathcal{Z} = \int \mathcal{D}(\delta\rho, \theta) e^{-\int d\tau \int d\vec{r} \mathcal{L}}$$

Integrate out $\delta\rho$, we get the effective Lagrangian,

$$\mathcal{L}_{\text{eff}} = \frac{1}{2g} (\partial_\tau \theta)^2 + \frac{\rho_0}{2m} (\nabla \theta)^2$$

- The $U(1)$ symmetry breaking generates a massless θ mode, it is our Goldstone boson in superfluid, phonon.

Minimal Coupling

- Write the superfluid lagrangian in a more symmetric form:

$$\mathcal{L}(\phi) = i\frac{1}{2}(\phi\partial_t\phi^* - \phi^*\partial_t\phi) + \frac{1}{2m}\partial_x\phi^*\partial_x\phi - \mu|\phi|^2 + \frac{g}{2}|\phi|^4$$
- By minimal coupling between the bosons and the electromagnetic gauge field A_μ :

$$\begin{aligned}\mathcal{L}(\phi, A_\mu) = & i\frac{1}{2}(\phi(\partial_0 - iA_0)\phi^* - \phi^*(\partial_0 + iA_0)\phi) - \frac{1}{2m}|(\partial_i + iA_i)\phi|^2 \\ & + V(\phi) + \frac{1}{4}F_{\mu\nu}F^{\mu\nu}\end{aligned}$$

The lagrangian is therefore gauge invariant under

$$\begin{aligned}\phi & \rightarrow e^{if(\vec{x},t)}\phi \\ A_\mu & \rightarrow A_\mu - \partial_\mu f\end{aligned}$$

Minimal Coupling

- Write the superfluid lagrangian in a more symmetric form:

$$\mathcal{L}(\phi) = i\frac{1}{2}(\phi\partial_t\phi^* - \phi^*\partial_t\phi) + \frac{1}{2m}\partial_x\phi^*\partial_x\phi - \mu|\phi|^2 + \frac{g}{2}|\phi|^4$$
- By minimal coupling between the bosons and the electromagnetic gauge field A_μ :

$$\begin{aligned}\mathcal{L}(\phi, A_\mu) = & i\frac{1}{2}(\phi(\partial_0 - iA_0)\phi^* - \phi^*(\partial_0 + iA_0)\phi) - \frac{1}{2m}|(\partial_i + iA_i)\phi|^2 \\ & + V(\phi) + \frac{1}{4}F_{\mu\nu}F^{\mu\nu}\end{aligned}$$

The lagrangian is therefore gauge invariant under

$$\begin{aligned}\phi &\rightarrow e^{if(\vec{x},t)}\phi \\ A_\mu &\rightarrow A_\mu - \partial_\mu f\end{aligned}$$

Anderson-Higgs Mechanism

- Select the ground state and fix the gauge so that ϕ is real, then $\mathcal{L} = A_0\phi^2 + \frac{1}{2m}(\partial_i\phi)^2 + \frac{\phi^2}{2m}(A_i)^2 + V(\phi) + \frac{1}{4}F_{\mu\nu}F^{\mu\nu}$
- Integrating out the fluctuation $\phi = \phi_0 + \delta\phi$, we get the low energy effective theory,

$$\mathcal{L}_{eff} = -\frac{1}{2g}A_0^2 + \frac{\rho}{2m}A_i^2 + \frac{1}{2}(\mathbf{B}^2 - \mathbf{E}^2)$$

Since A_0 contain no time derivative terms, it's not dynamical and can be integrated out to give

$$\mathcal{L}_{eff} = -\frac{1}{2}\partial_0 A_j(\delta_{ij} + g\partial_j\partial_i)\partial_0 A_i + \frac{\rho}{2m}A_i^2 + \frac{1}{2}\mathbf{B}^2$$

- The electromagnetic vector potential \vec{A} "eats" the Goldstone boson to acquire mass!

Anderson-Higgs Mechanism

- Select the ground state and fix the gauge so that ϕ is real, then $\mathcal{L} = A_0\phi^2 + \frac{1}{2m}(\partial_i\phi)^2 + \frac{\phi^2}{2m}(A_i)^2 + V(\phi) + \frac{1}{4}F_{\mu\nu}F^{\mu\nu}$
- Integrating out the fluctuation $\phi = \phi_0 + \delta\phi$, we get the low energy effective theory,

$$\mathcal{L}_{eff} = -\frac{1}{2g}A_0^2 + \frac{\rho}{2m}A_i^2 + \frac{1}{2}(\mathbf{B}^2 - \mathbf{E}^2)$$

Since A_0 contain no time derivative terms, it's not dynamical and can be integrated out to give

$$\mathcal{L}_{eff} = -\frac{1}{2}\partial_0 A_j(\delta_{ij} + g\partial_j\partial_i)\partial_0 A_i + \frac{\rho}{2m}A_i^2 + \frac{1}{2}\mathbf{B}^2$$

- The electromagnetic vector potential \vec{A} "eats" the Goldstone boson to acquire mass!

Anderson-Higgs Mechanism

- Select the ground state and fix the gauge so that ϕ is real, then $\mathcal{L} = A_0\phi^2 + \frac{1}{2m}(\partial_i\phi)^2 + \frac{\phi^2}{2m}(A_i)^2 + V(\phi) + \frac{1}{4}F_{\mu\nu}F^{\mu\nu}$
- Integrating out the fluctuation $\phi = \phi_0 + \delta\phi$, we get the low energy effective theory,

$$\mathcal{L}_{eff} = -\frac{1}{2g}A_0^2 + \frac{\rho}{2m}A_i^2 + \frac{1}{2}(\mathbf{B}^2 - \mathbf{E}^2)$$

Since A_0 contain no time derivative terms, it's not dynamical and can be integrated out to give

$$\mathcal{L}_{eff} = -\frac{1}{2}\partial_0 A_j(\delta_{ij} + g\partial_j\partial_i)\partial_0 A_i + \frac{\rho}{2m}A_i^2 + \frac{1}{2}\mathbf{B}^2$$

- The electromagnetic vector potential \vec{A} "eats" the Goldstone boson to acquire mass!

Anderson-Higgs Mechanism

- Select the ground state and fix the gauge so that ϕ is real, then $\mathcal{L} = A_0\phi^2 + \frac{1}{2m}(\partial_i\phi)^2 + \frac{\phi^2}{2m}(A_i)^2 + V(\phi) + \frac{1}{4}F_{\mu\nu}F^{\mu\nu}$
- Integrating out the fluctuation $\phi = \phi_0 + \delta\phi$, we get the low energy effective theory,

$$\mathcal{L}_{eff} = -\frac{1}{2g}A_0^2 + \frac{\rho}{2m}A_i^2 + \frac{1}{2}(\mathbf{B}^2 - \mathbf{E}^2)$$

Since A_0 contain no time derivative terms, it's not dynamical and can be integrated out to give

$$\mathcal{L}_{eff} = -\frac{1}{2}\partial_0 A_j(\delta_{ij} + g\partial_j\partial_i)\partial_0 A_i + \frac{\rho}{2m}A_i^2 + \frac{1}{2}\mathbf{B}^2$$

- The electromagnetic vector potential \vec{A} "eats" the Goldstone boson to acquire mass!

Meissner Effect

In BCS superconductivity, Cooper pairs can be viewed as bosonic composite particles. So it has the similar Lagrangian as superfluid. Except it assumes that:

1. No electric field, i.e., $A_0 = 0, \partial_0 A_i = 0$.
2. $\nabla \cdot \vec{A} = 0 \Rightarrow \mathbf{q} \cdot \mathbf{A}_q = 0$ in momentum space.

$$\mathcal{S}[\mathbf{A}]_{\text{eff}} = \frac{\beta}{2} \sum_q \left(\frac{\rho}{m} + q^2 \right) \mathbf{A}_q \cdot \mathbf{A}_{-q}$$

Vary the action with respect to \mathbf{A} , we get $(\frac{\rho}{m} + q^2) \mathbf{A}_q = 0$, or

London equation

$$\left(\frac{\rho}{m} - \nabla^2 \right) \mathbf{B}(\mathbf{r}) = 0$$

Meissner Effect

In BCS superconductivity, Cooper pairs can be viewed as bosonic composite particles. So it has the similar Lagrangian as superfluid. Except it assumes that:

1. No electric field, i.e., $A_0 = 0, \partial_0 A_i = 0$.
2. $\nabla \cdot \vec{A} = 0 \Rightarrow \mathbf{q} \cdot \mathbf{A}_q = 0$ in momentum space.

$$\mathcal{S}[\mathbf{A}]_{\text{eff}} = \frac{\beta}{2} \sum_q \left(\frac{\rho}{m} + q^2 \right) \mathbf{A}_q \cdot \mathbf{A}_{-q}$$

Vary the action with respect to \mathbf{A} , we get $(\frac{\rho}{m} + q^2) \mathbf{A}_q = 0$, or

London equation

$$\left(\frac{\rho}{m} - \nabla^2 \right) \mathbf{B}(\mathbf{r}) = 0$$

Meissner Effect

In BCS superconductivity, Cooper pairs can be viewed as bosonic composite particles. So it has the similar Lagrangian as superfluid. Except it assumes that:

1. No electric field, i.e., $A_0 = 0, \partial_0 A_i = 0$.
2. $\nabla \cdot \vec{A} = 0 \Rightarrow \mathbf{q} \cdot \mathbf{A}_q = 0$ in momentum space.

$$\mathcal{S}[\mathbf{A}]_{\text{eff}} = \frac{\beta}{2} \sum_q \left(\frac{\rho}{m} + q^2 \right) \mathbf{A}_q \cdot \mathbf{A}_{-q}$$

Vary the action with respect to \mathbf{A} , we get $(\frac{\rho}{m} + q^2) \mathbf{A}_q = 0$, or

London equation

$$\left(\frac{\rho}{m} - \nabla^2 \right) \mathbf{B}(\mathbf{r}) = 0$$

Meissner effect


London equation

$$\left(\frac{\rho}{m} - \nabla^2\right)\mathbf{B}(\mathbf{r}) = 0$$

For $\rho \neq 0$, $\mathbf{B}(x) \propto \mathbf{B}_0 \exp(-x/\lambda)$, where

$$\lambda = \sqrt{\frac{m}{\rho}}$$

is the **penetration depth**.



Reference

- A. Altland and B. D. Simons, *Condensed matter field theory* (Cambridge University Press, 2010).
- X.-G. Wen, *Quantum field theory of many-body systems: from the origin of sound to an origin of light and electrons* (Oxford University Press on Demand, 2004).
- H. Bertle, *Goldstone boson and higgs mechanism*, URL https://www.ethz.ch/content/dam/ethz/special-interest/phys/theoretical-physics/itp-dam/documents/gaberdiel/proseminar_fs2018/21_Bertle.pdf.