

Modelling With Differential Equations

August 2019

Charles Driver

Max Planck Institute for Human Development





■ Aims:



- Aims:
 - Basic concepts



- Aims:
 - Basic concepts
 - Advanced possibilities



- Aims:
 - Basic concepts
 - Advanced possibilities
 - Software workflow



- Aims:
 - Basic concepts
 - Advanced possibilities
 - Software workflow
- Content:



- Aims:
 - Basic concepts
 - Advanced possibilities
 - Software workflow
- Content:
 - What is a differential equation?



■ Aims:

- Basic concepts
- Advanced possibilities
- Software workflow

■ Content:

- What is a differential equation?
- Why are they useful?



- Aims:

- Basic concepts
- Advanced possibilities
- Software workflow

- Content:

- What is a differential equation?
- Why are they useful?
- Measurement



- Aims:
 - Basic concepts
 - Advanced possibilities
 - Software workflow
- Content:
 - What is a differential equation?
 - Why are they useful?
 - Measurement
 - Known inputs



■ Aims:

- Basic concepts
- Advanced possibilities
- Software workflow

■ Content:

- What is a differential equation?
- Why are they useful?
- Measurement
- Known inputs
- Uncertainty / unknown inputs



■ Aims:

- Basic concepts
- Advanced possibilities
- Software workflow

■ Content:

- What is a differential equation?
- Why are they useful?
- Measurement
- Known inputs
- Uncertainty / unknown inputs
- Complexity – state dependent parameters, random effects, etc.



What is a differential equation?











```
y0 = .2  
t=4  
b=.5  
y= y0 + b * t  
print(y)  
[1] 2.2
```

- $\frac{dy}{dt} = b$
- $\dot{y} = b$
- $y(t) = y_{t0} + bt$



How can we use differential equations?





```
y0 = 0.2
t= 4
b= 0.5
A = -0.3
y= exp(A * t) * y0 + (exp(A*t)-1)*(1/A)*b
print(y)
[1] 1.224915
```

- $\frac{dy}{dt} = Ay + b$
- $\dot{y} = Ay + b$
- $y(t) = e^{At}y_{t0} + (e^{At} - 1)A^{-1}b$



$$\underbrace{d \begin{bmatrix} \eta_1 \\ \eta_2 \end{bmatrix}}_{d\boldsymbol{\eta}(t)}(t) = \left(\underbrace{\begin{bmatrix} -0.3 & 0 \\ 0.4 & -0.1 \end{bmatrix}}_{\mathbf{A}} \underbrace{\begin{bmatrix} \eta_1 \\ \eta_2 \end{bmatrix}}_{\boldsymbol{\eta}(t)}(t) + \underbrace{\begin{bmatrix} 1 \\ 0 \end{bmatrix}}_{\mathbf{b}} \right) dt$$

$$\underbrace{\begin{bmatrix} Y_1 \\ Y_2 \\ Y_3 \end{bmatrix}}_{\mathbf{Y}(t)}(t) = \underbrace{\begin{bmatrix} 1 & 0 \\ 0.8 & 0 \\ 0 & 1 \end{bmatrix}}_{\mathbf{\Lambda}} \underbrace{\begin{bmatrix} \eta_1 \\ \eta_2 \end{bmatrix}}_{\boldsymbol{\eta}(t)}(t) + \underbrace{\begin{bmatrix} 1.5 \\ 0 \\ -2 \end{bmatrix}}_{\boldsymbol{\tau}} + \underbrace{\begin{bmatrix} \epsilon_1 \\ \epsilon_2 \\ \epsilon_3 \end{bmatrix}}_{\boldsymbol{\epsilon}(t)}(t)$$



Why are differential equations useful?





- Direct model of the process:



- Direct model of the process:
 - Difference form: $G(t) = G_{t0} + .98G(t - 1)$



- Direct model of the process:
 - Difference form: $G(t) = G_{t0} + .98G(t - 1)$
 - Differential form: $\dot{G}(t) = -.02G(t)$



- Direct model of the process:
 - Difference form: $G(t) = G_{t0} + .98G(t - 1)$
 - Differential form: $\dot{G}(t) = -.02G(t)$
- Increment size for t doesn't matter – can vary between different data points.



- Direct model of the process:
 - Difference form: $G(t) = G_{t0} + .98G(t - 1)$
 - Differential form: $\dot{G}(t) = -.02G(t)$
- Increment size for t doesn't matter – can vary between different data points.
- In multivariate systems, zeroes only imply 'no effect' in differential equation formulation.



- Direct model of the process:
 - Difference form: $G(t) = G_{t0} + .98G(t - 1)$
 - Differential form: $\dot{G}(t) = -.02G(t)$
- Increment size for t doesn't matter – can vary between different data points.
- In multivariate systems, zeroes only imply 'no effect' in differential equation formulation.
- Coupled with appropriate estimation routines – allows relatively unconstrained theory specification and testing.



Not just time!






Not just time!



- Useful for modelling change over any continuous variable.






Not just time!



- Useful for modelling change over any continuous variable.
- Besides time, age is an obvious candidate for us.





- Multivariate, hierarchical, continuous time state space modelling.



- Multivariate, hierarchical, continuous time state space modelling.
- Frequentist, Bayesian, or somewhere in the middle.



- Multivariate, hierarchical, continuous time state space modelling.
- Frequentist, Bayesian, or somewhere in the middle.
- Good reasons to use: More flexible than typical regression / SEM approaches. Random / time varying effects on any parameter. Estimation works under problematic conditions, many parameters etc.



- Multivariate, hierarchical, continuous time state space modelling.
- Frequentist, Bayesian, or somewhere in the middle.
- Good reasons to use: More flexible than typical regression / SEM approaches. Random / time varying effects on any parameter. Estimation works under problematic conditions, many parameters etc.
- Good reasons to avoid: More flexible than typical regression / SEM approaches. Less developed workflow, doesn't cope with high dimensions in variable domain.