

Knot Surgery and Integer Characterizing Slopes

Gabriel Agostini, Sophia Chen, Christian Serio, Cecilia Wang, Anton Wu, and Kexin Wu

Advisors: Kyle Hayden and Aliakbar Daemi

Columbia University

August 1, 2019

Table of Contents

Knots and links in the 3-sphere

Definition

A *knot* K is the image of a smooth embedding of the circle S^1 into a 3-manifold, usually the 3-sphere S^3 . In particular, K is diffeomorphic to S^1 . A *link* L is a disjoint union of knots, which may be knotted together.

Definition

Let M, N be manifolds and $g, h: N \rightarrow M$ embeddings. An *ambient isotopy* of M carrying g to h is a continuous map $F: M \times [0, 1] \rightarrow M$, such that $F_t = F(\cdot, t)$ is a homeomorphism of M for each $t \in [0, 1]$, $F_0 = \mathbb{1}$, and $F_1 \circ g = h$.

- We regard two knots $K, K' \subset S^3$ to be equivalent if they differ by an ambient isotopy of S^3 . We write $K \simeq K'$.
- Equivalently, we can view knots as subsets of \mathbb{R}^3 rather than S^3 .

Knot diagrams

- We can study a knot $K \subset \mathbb{R}^3$ by projecting it onto a hyperplane \mathbb{R}^2 .
- If $\pi: \mathbb{R}^3 \rightarrow \mathbb{R}^2$ is a projection such that $\pi(K)$ is an embedded curve except at finitely many *crossing points*, then $\pi(K)$ is a *diagram* for K .
- The *crossing number* $c(K)$ is the minimum number of crossings in a diagram of K .




Figure: Knots with $c(K) \leq 7$





Figure: Hopf link

Reidemeister moves


Theorem (Reidemeister)

Two knots K, K' are isotopic if and only if they have diagrams that differ by a sequence of planar isotopies and Reidemeister moves.



Crossings

- If we orient a knot K , then we can define a *sign* for each crossing by the right-hand rule.
- For a two-component link $L = K \cup K'$, the *linking number* $\text{lk}(L)$ is one half the sum of the signs of the crossings between K and K' in a diagram of L .



Unknotting number and band move

Definition

The *unknotting number* $u(K)$ of a knot K is the minimal number of crossing changes that are required to change some diagram of K into a diagram of the unknot.

- If $u(K)=0$, then K is an unknot.


Definition

Let K be any knot(or link) in S^3 and let $b \subset S^3$ be any embedded band where one pair of sides lie along arcs in K and where b is otherwise disjoint from K . Join K to itself by deleting the arcs $K \cap b$ and adding the arcs forming the other side of b . The resulting knot(or link) K' is obtained from K by a *band move* along b .

Example of band move and band presentation

- Example of band move:

The Stevedore knot 6_1 is obtained from a band move on two unknots:




- If K can be obtained from another knot(or link) K' by a single band move, then we say K has a *band presentation*.
- If K can be obtained from a Hopf link by a single band move, then K has a *banded Hopf link presentation*.

Example of banded Hopf link presentation

Theorem

If K has $u(K) = 1$, then it is obtained from the Hopf link by a single band move.


- Example: banded Hopf link presentation of Trefoil:



Twist knots and Twisted whitehead double

Definition

A *twist knot* K is a knot obtained by repeatedly twisting an unknot and linking the ends together.



Definition

A knot K is a *twisted Whitehead double* if there exists a band presentation for K in which the band does not cross either component of the Hopf link.

- Note: twist knots are twisted Whitehead doubles, but not all twisted Whitehead doubles are twist knots.

Simple Closed Curves

Definition

A *simple closed curve* (s.c.c.) is a submanifold of a smooth manifold X , with itself diffeomorphic to S^1 .



a s.c.c



a non-simple closed curve

- By definition, a s.c.c. is embedded in some smooth manifold and is intuitively a 1-dimensional curve diffeomorphic to S^1 without any self-intersection.
- In particular, we can embed s.c.c.'s in a manifold such as a torus T^2 . An alternate definition for a knot $K \subset S^3$ is a simple closed curve in S^3 .

Embedded s.c.c.'s on T^2


Definition

A s.c.c. $\gamma \subset T^2$ is called *non-separating* if $T^2 - \gamma$ is connected.


- Every non-separating s.c.c. in T^2 bounds a disc.
- Any two oriented and non-separating s.c.c.'s $\gamma, \gamma' \subset T^2$ are isotopic if and only if their signed intersection, the sum of their local intersections, is 0.

Definition

Any two s.c.c.'s $\gamma, \gamma' \subset T^2$ form a *basis* if they are non-separating and intersecting in a single point.



γ_1 = separating




γ_2, γ_3 = non-separating

Preferred meridian and longitude pair for a knot $K \subset S^3$

Definition

Given a knot $K \subset S^3$, $N(K) \cong S^1 \times D^2 \cong$ a solid torus \mathbb{T} where $\partial N(K) \cong$ a torus T^2 , we define a *meridian* μ of K to be an unknot $\partial D^2 \times 1 \subset \partial N(K) \cong T^2$ that bounds a disc which K crosses exactly once. Then we choose a longitude of K as a circle $\lambda = 1 \times S^1 \subset \partial N(K) \cong T^2$, which is a parallel copy of K on $\partial N(K)$.



- Note that our choice of the longitude λ for K intersects the meridian μ non-tangentially in exactly one point, and thus (μ, λ) form a basis of T^2 .

Isotopy classes of s.c.c.'s on T^2

- If a preferred basis of a torus T^2 is (m, l) , then any s.c.c $\gamma \subset T^2$ can be determined (up to isotopy) by two numbers, $p := |n(\gamma \cap m)|$ and $q := |n(\gamma \cap l)|$ where $n(a \cap b)$ is the signed intersection between the two oriented simple closed curves a and b on T^2 .
- Further, isotopy classes s.c.c's on a torus are classified by the set of extended rational numbers $\mathbb{Q} \cup \{\infty\}$.
- Hence, given a knot $K \in S^3$ where $\partial N(K) \cong$ a torus T^2 , any $p/q \in \mathbb{Q} \cup \{\infty\}$ determines a s.c.c β on $\partial N(K)$ that is well-defined up to isotopy. Intuitively, p (and q resp.) is the number of of times that β goes around the meridian (and longitude resp.) that we choose for K .
- We use β to define a 3-manifold $S^3_{p/q}(K)$, called p/q -surgery on K .



Surgery on a knot: *drilling* then *filling*

Definition

Given a knot $K \subset S^3$, the *exterior* of K , denoted $X(K)$, is a manifold with boundary obtained by removing the interior of $N(K) = S^1 \times D^2 \cong$ a solid torus \mathbb{T} from S^3 . Then $\partial X(K) = \partial N(K) \cong$ a torus T^2 . Let β be the s.c.c on T^2 determined (up to isotopy) by some extended rational number $p/q \in \mathbb{Q} \cup \{\infty\}$. Then the p/q -surgery on K , a 3-manifold denoted $S^3_{p/q}(K)$, is obtained by gluing another solid torus \mathbb{T}' back to $X(K)$ so that the meridian of \mathbb{T}' is identified with the curve β .

Examples of p/q -surgery on a knot

- Example: $\pm 1/0$ -surgery on a knot K gives back S^3 trivially. Thus any nontrivial surgery on a knot K has rational slopes.
- Example: $S^3_{1/n}(u) \cong S^3$ where u is an unknot and $n \in \mathbb{Z}$.




Surgery on a link

Definition

Similarly, given a link L in S^3 s.t. $\exists p/q \in \mathbb{Q} \cup \{\infty\}$ associated to each component of L , we can do surgery on L by doing surgery along each of its component. Any such link is called a *framed link*.

- Example: for any integer n , $(n, 0)$ -surgery on a Hopf link K yields S^3 .




Theorem

Any oriented, connected 3-manifold can be obtained by doing surgery along a framed link. So we can think of any 3-manifold in terms of a diagram for a framed link. Any such diagram is called a surgery diagram.

Example of surgery on a link


Theorem

For any integer n and any knot K with an unknot c as its meridian, $(n, 0)$ -surgery on $K \cup c$ yields S^3 . Such a pair of K and c forms a cancelling pair that can be deleted without changing the 3-manifold described by the surgery diagram.



Surgery dual to a knot

- Note that K is given by $S^1 \times 0$ inside $N(K) = S^1 \times D^2 \cong$ a solid torus \mathbb{T} . i.e. K is the core curve of the solid torus $N(K)$.




Definition

When doing p/q -surgery on a knot $K \in S^3$, the new solid torus $\mathbb{T}' \cong S^1 \times D^2$ that we glue back in $X(K)$ to produce $S^3_{p/q}(K)$ also has a core curve $S^1 \times 0$, which specifies a knot γ in $S^3_{p/q}(K)$. Call γ the *surgery dual* of K .

Surgery dual to a knot (cont.)

- For any knot K and any integer n , the *surgery dual* to K in $S_n^3(K)$ can be represented as a meridian to K in the surgery diagram.



$Y = S_0^3(K)$, with $K = 11n38$

Lemma

Surgery duality for knots in S^3 is symmetric.

$$\begin{array}{ccccccc}
 S^3 & n \in \mathbb{Z} & S_n^3(K) & & S^3 \\
 U & n\text{-surgery} & U & 0\text{-surgery} & U \\
 K & \rightsquigarrow & \gamma & \rightsquigarrow & K
 \end{array}$$

Surgery dual link to a framed link

Definition

A *surgery dual link* to a framed link $L = \bigcup_{i=1}^N L_i \in S^3$ is a link $L' = \bigcup_{i=1}^N L'_i$, where each component L'_i is the surgery dual to L_i in the surgered manifold Y that we obtain after doing surgery on L .

- Suppose that Dehn surgery on two surgery diagrams D_1, D_2 gives two 3-manifolds M_1, M_2 .

- Suppose that Dehn surgery on two surgery diagrams D_1 , D_2 gives two 3-manifolds M_1 , M_2 .
- Then M_1 and M_2 are homeomorphic if and only if D_1 and D_2 are related by a sequence of *Kirby moves*.

- Suppose that Dehn surgery on two surgery diagrams D_1 , D_2 gives two 3-manifolds M_1 , M_2 .
- Then M_1 and M_2 are homeomorphic if and only if D_1 and D_2 are related by a sequence of *Kirby moves*.
 - Handle-slide

- Suppose that Dehn surgery on two surgery diagrams D_1 , D_2 gives two 3-manifolds M_1 , M_2 .
- Then M_1 and M_2 are homeomorphic if and only if D_1 and D_2 are related by a sequence of *Kirby moves*.
 - Handle-slide
 - Rolfsen twist


- Suppose that Dehn surgery on two surgery diagrams D_1 , D_2 gives two 3-manifolds M_1 , M_2 .
- Then M_1 and M_2 are homeomorphic if and only if D_1 and D_2 are related by a sequence of *Kirby moves*.
 - Handle-slide
 - Rolfsen twist
 - Slam dunk

Handle-slide

- The *handle-slide* allows us to change the linking number between two link components.

Handle-slide

- The *handle-slide* allows us to change the linking number between two link components.




Rolfsen twist

- The *Rolfsen twist* allows us to remove an n -framed unknot while n -twisting the strands passing through it.

Rolfsen twist

- The *Rolfsen twist* allows us to remove an n -framed unknot while n -twisting the strands passing through it.

Rolfsen twist:



Slam dunk

- A 0-framed link component K with a $(-1/n)$ -framed meridian is equivalent to K with n -framing.

Slam dunk

- A 0-framed link component K with a $(-1/n)$ -framed meridian is equivalent to K with n -framing.




Table of Contents

Piccirillo's construction: Knots with the same surgery

Theorem (Piccirillo 2018)

Let $L = R \cup G \cup B$ be a surgery diagram for some 3-manifold Y such that:

- ① R is a zero-framed unknot, B and G have integral framings.
- ② Ignoring B , R is isotopic to a meridian of G .
- ③ Ignoring G , R is isotopic to a meridian of B .
- ④ B and G have linking number 0.

Then, there exist knots K and K' such that $Y \cong S_n^3(K) \cong S_n^3(K')$.

- Piccirillo's construction comes from an older construction, the *dualizable patterns* construction, to produce knots with the same surgery.

Piccirillo's construction (cont.)





Figure: Diagram of a link L used by Piccirillo on her original paper. The colors (red, blue, and green) match the component letters R , B , G .

Baker-Motegi: Knots with the same surgery

- We adapted the following construction from Baker and Motegi (2018).
- Let K be a knot, and suppose we can take an unknot c linked with K such that $(0,0)$ -surgery on $K \cup c$ is S^3 .
- Baker-Motegi present a method for producing knots K'_n with $S_n^3(K) \cong S_n^3(K'_n)$ from this link.
- Define K' to be the surgery dual to c in $S^3 = S^3_{(0,0)}(K \cup c)$.
- Then K' has the same 0-surgery as K .

- Also define c' to be the surgery dual to K in S^3 .
- After some Kirby calculus, we find that in the surgered manifold $S^3 = S^3_{(0,0)}(K \cup c)$, c' is an unknot linked with K' .
- Let K'_n be the result of twisting K' through c' , n times.
- We say that $\{K'_n\}$ forms a *twist family*.
- Then K'_n has the same n -surgery as K for all n .
- Moreover, if c is not a meridian to K , then $K \simeq K'_n$ for at most finitely many n .

Baker-Motegi Illustration




Obtaining a link $K \cup c$ when $u(K) = 1$ (Piccirillo)

- In the special case where K has unknotting number one, we can use a band presentation for K .
- We start with a Hopf link $R \cup B$ and slide one component over the other according to the band presentation for K .
- R remains an unknot, which we rename c , and B becomes K .


Lemma

Let $K \cup c$ be the link obtained by the handle slide above. Then $(0, 0)$ -surgery on $K \cup c$ gives S^3 .



Obtaining $K \cup c$ when $u(K) > 1$

- We do not have a systematic way of finding a link $K \cup c$ for a given knot K with $u(K) > 1$.
- If we perform multiple slides on a Hopf link, then we obtain a link $K \cup c$ for some knot K with higher unknotting number, but we have no way of predicting what K will be.
- Example: 10_{125}



Applications of the construction

- We have produced a banded Hopf link presentation for all the 78 unknotting number one knots K with $c(K) \leq 10$.
- We followed the above procedure to manually produce a diagram for a knot K' that has the same zero-surgery as K , using a software called KLO which can perform Kirby calculus on link diagrams.
- Using a software called SnapPy, we were able to verify that whenever K was not a twist knot, K' was not isotopic to K . Thus, zero was not a characterizing slope.
- We can use a similar process to check whether any integer slope is characterizing.
- **Question:** Can we classify all integer slopes once we have the link $K \cup c$, with finitely many computations?

Characterizing slopes

- Let K be a knot. Then p/q is a *characterizing slope* for K if $S_{p/q}^3(K')$ is not homeomorphic to $S_{p/q}^3(K)$ for any knot $K' \neq K$.

Characterizing slopes

- Let K be a knot. Then p/q is a *characterizing slope* for K if $S^3_{p/q}(K')$ is not homeomorphic to $S^3_{p/q}(K)$ for any knot $K' \neq K$.
- The following theorem shows that for a certain type of knot, most rational slopes are characterizing.

Characterizing slopes

- Let K be a knot. Then p/q is a *characterizing slope* for K if $S^3_{p/q}(K')$ is not homeomorphic to $S^3_{p/q}(K)$ for any knot $K' \neq K$.
- The following theorem shows that for a certain type of knot, most rational slopes are characterizing.

Theorem (Ni–Zhang, 2014)

For the torus knot $T_{r,s}$, any slope $p/q > \frac{30}{67}(r^2 - 1)(s^2 - 1)$ is characterizing.

Characterizing slopes

- However, the next theorem shows that for a certain type of knot, there are infinitely many non-characterizing slopes.

Characterizing slopes

- However, the next theorem shows that for a certain type of knot, there are infinitely many non-characterizing slopes.

Theorem (Baker–Moteği, 2018)

Let $K \subset S^3$ be a knot, where there is an unknot c which is not a meridian to K such that $(0, 0)$ -surgery on $K \cup c$ yields S^3 . Then K has infinitely many non-characterizing slopes.

Characterizing slopes

- However, the next theorem shows that for a certain type of knot, there are infinitely many non-characterizing slopes.

Theorem (Baker–Motegi, 2018)

Let $K \subset S^3$ be a knot, where there is an unknot c which is not a meridian to K such that $(0, 0)$ -surgery on $K \cup c$ yields S^3 . Then K has infinitely many non-characterizing slopes.

- Our work shows that for a certain type of knot, most integer slopes are non-characterizing!

Theorem

- ① *If K has unknotting number $u(K) = 1$ and K is not a twist knot, then K has at most one integer characterizing slope, namely ± 2 .*
- ② *If K is one of the knots 8_4 , 8_6 , 8_{10} , or 8_{12} , then K has $u(K) > 1$ and has no integer characterizing slopes.*
- ③ *If K is the twist knot 8_1 , then K has at most one integer characterizing slope, namely 0 .*

Theorem

- ① *If K has unknotting number $u(K) = 1$ and K is not a twist knot, then K has at most one integer characterizing slope, namely ± 2 .*
- ② *If K is one of the knots 8_4 , 8_6 , 8_{10} , or 8_{12} , then K has $u(K) > 1$ and has no integer characterizing slopes.*
- ③ *If K is the twist knot 8_1 , then K has at most one integer characterizing slope, namely 0 .*

Theorem

If a knot K has unknotting number $u(K) = 1$ and is not a twisted Whitehead double, then K has at most finitely many integer characterizing slopes.

Hyperbolic Dehn surgery

- A *hyperbolic 3-manifold* is a compact, orientable 3-manifold which admits a complete hyperbolic metric; that is, a Riemannian metric with constant negative sectional curvature.

Hyperbolic Dehn surgery

- A *hyperbolic 3-manifold* is a compact, orientable 3-manifold which admits a complete hyperbolic metric; that is, a Riemannian metric with constant negative sectional curvature.
- Hyperbolic 3-manifolds, being compact, have finite hyperbolic volume.

Hyperbolic Dehn surgery

- A *hyperbolic 3-manifold* is a compact, orientable 3-manifold which admits a complete hyperbolic metric; that is, a Riemannian metric with constant negative sectional curvature.
- Hyperbolic 3-manifolds, being compact, have finite hyperbolic volume.
- A knot $K \subset S^3$ is a *hyperbolic knot* if $S^3 \setminus N(K)$ is a hyperbolic 3-manifold.

Hyperbolic Dehn surgery

- A *hyperbolic 3-manifold* is a compact, orientable 3-manifold which admits a complete hyperbolic metric; that is, a Riemannian metric with constant negative sectional curvature.
- Hyperbolic 3-manifolds, being compact, have finite hyperbolic volume.
- A knot $K \subset S^3$ is a *hyperbolic knot* if $S^3 \setminus N(K)$ is a hyperbolic 3-manifold.
- If K is a hyperbolic knot, then the torus $\partial N(K)$ inherits a Euclidean metric.

Hyperbolic Dehn surgery

- A *hyperbolic 3-manifold* is a compact, orientable 3-manifold which admits a complete hyperbolic metric; that is, a Riemannian metric with constant negative sectional curvature.
- Hyperbolic 3-manifolds, being compact, have finite hyperbolic volume.
- A knot $K \subset S^3$ is a *hyperbolic knot* if $S^3 \setminus N(K)$ is a hyperbolic 3-manifold.
- If K is a hyperbolic knot, then the torus $\partial N(K)$ inherits a Euclidean metric.
- According to this Euclidean metric, the *length* $\ell(p/q)$ of a slope p/q is the length of the shortest curve on the torus $\partial N(K)$ with slope p/q .

Hyperbolic surgery theorems

Theorem (Gromov-Thurston)

Let K be a hyperbolic knot. If $\ell(p/q) > 2\pi$, then the (p/q) -filling on $S^3 \setminus N(K)$ is hyperbolic.

Hyperbolic surgery theorems

Theorem (Gromov-Thurston)

Let K be a hyperbolic knot. If $\ell(p/q) > 2\pi$, then the (p/q) -filling on $S^3 \setminus N(K)$ is hyperbolic.

Theorem (Futer, et al.)

Let K be a hyperbolic knot; let $X := S^3 \setminus N(K)$. If $\ell(p/q) > 2\pi$, then the volume $\text{vol}(X_{p/q})$ of the (p/q) -filling on X satisfies

$$\text{vol}(X_{p/q}) \geq \left(1 - \left(\frac{2\pi}{\ell(p/q)}\right)^2\right)^{3/2} \text{vol}(X).$$

Theorem (Cooper-Lackenby)

Let X be a cusped hyperbolic 3-manifold, and suppose s_1, s_2 are two slopes on a torus $T \subset \partial X$. Then

$$\ell(s_1)\ell(s_2) \geq \sqrt{3} \Delta(s_1, s_2),$$

where $\Delta(s_1, s_2)$ is the minimum possible number of intersections between a curve with slope s_1 and a curve with slope s_2 on T .

Sketch of proof of Theorem

Theorem

If a knot K has unknotting number $u(K) = 1$ and is not a twisted Whitehead double, then K has at most finitely many integer characterizing slopes.

Sketch of proof of Theorem

- Following Baker–Moteği, we can produce a link $(K \cup c) \subset S^3$, where c is an unknot which is not a meridian to K such that $(0, 0)$ -surgery on $K \cup c$ yields S^3 .

Sketch of proof of Theorem

- Following Baker–Moteği, we can produce a link $(K \cup c) \subset S^3$, where c is an unknot which is not a meridian to K such that $(0, 0)$ -surgery on $K \cup c$ yields S^3 .
- Add a meridian m to c ; notice that m is K'_n , in the non-standard representation of S^3 .

Sketch of proof of Theorem

- Following Baker–Moteği, we can produce a link $(K \cup c) \subset S^3$, where c is an unknot which is not a meridian to K such that $(0, 0)$ -surgery on $K \cup c$ yields S^3 .
- Add a meridian m to c ; notice that m is K'_n , in the non-standard representation of S^3 .
- Hence removing a neighborhood of m and performing 0-surgery on c gives a hyperbolic manifold X on which n -filling yields $S^3 \setminus K'_n$.

Sketch of proof of Theorem

- Following Baker–Moteği, we can produce a link $(K \cup c) \subset S^3$, where c is an unknot which is not a meridian to K such that $(0, 0)$ -surgery on $K \cup c$ yields S^3 .
- Add a meridian m to c ; notice that m is K'_n , in the non-standard representation of S^3 .
- Hence removing a neighborhood of m and performing 0-surgery on c gives a hyperbolic manifold X on which n -filling yields $S^3 \setminus K'_n$.
- Define $v := \text{vol}(S^3 \setminus K)$, $v'_n := \text{vol}(S^3 \setminus K'_n)$ and $v_X := \text{vol}(X)$.

Sketch of proof of Theorem

- Following Baker–Moteği, we can produce a link $(K \cup c) \subset S^3$, where c is an unknot which is not a meridian to K such that $(0, 0)$ -surgery on $K \cup c$ yields S^3 .
- Add a meridian m to c ; notice that m is K'_n , in the non-standard representation of S^3 .
- Hence removing a neighborhood of m and performing 0-surgery on c gives a hyperbolic manifold X on which n -filling yields $S^3 \setminus K'_n$.
- Define $v := \text{vol}(S^3 \setminus K)$, $v'_n := \text{vol}(S^3 \setminus K'_n)$ and $v_X := \text{vol}(X)$.
- Now we use the above bounds to show that:

Sketch of proof of Theorem

- Following Baker–Moteği, we can produce a link $(K \cup c) \subset S^3$, where c is an unknot which is not a meridian to K such that $(0, 0)$ -surgery on $K \cup c$ yields S^3 .
- Add a meridian m to c ; notice that m is K'_n , in the non-standard representation of S^3 .
- Hence removing a neighborhood of m and performing 0-surgery on c gives a hyperbolic manifold X on which n -filling yields $S^3 \setminus K'_n$.
- Define $v := \text{vol}(S^3 \setminus K)$, $v'_n := \text{vol}(S^3 \setminus K'_n)$ and $v_X := \text{vol}(X)$.
- Now we use the above bounds to show that:

$$v'_n \geq \left(1 - \left(\frac{2\pi\ell}{|n|\sqrt{3}}\right)^2\right)^{3/2} v_X > v$$

$$\text{for } |n| > N := \frac{2\pi}{\sqrt{3}} \ell (1 - (v/v_X)^{2/3})^{-1/2}.$$

Sketch of proof of Theorem

- Following Baker–Moteği, we can produce a link $(K \cup c) \subset S^3$, where c is an unknot which is not a meridian to K such that $(0, 0)$ -surgery on $K \cup c$ yields S^3 .
- Add a meridian m to c ; notice that m is K'_n , in the non-standard representation of S^3 .
- Hence removing a neighborhood of m and performing 0-surgery on c gives a hyperbolic manifold X on which n -filling yields $S^3 \setminus K'_n$.
- Define $v := \text{vol}(S^3 \setminus K)$, $v'_n := \text{vol}(S^3 \setminus K'_n)$ and $v_X := \text{vol}(X)$.
- Now we use the above bounds to show that:

$$v'_n \geq \left(1 - \left(\frac{2\pi\ell}{|n|\sqrt{3}}\right)^2\right)^{3/2} v_X > v$$

for $|n| > N := \frac{2\pi}{\sqrt{3}} \ell (1 - (v/v_X)^{2/3})^{-1/2}$.

- Hence all integer slopes $|n| > N$ are non-characterizing.

Sketch of proof of Theorem

- Following Baker–Motegi, we can produce a link $(K \cup c) \subset S^3$, where c is an unknot which is not a meridian to K such that $(0, 0)$ -surgery on $K \cup c$ yields S^3 .
- Add a meridian m to c ; notice that m is K'_n , in the non-standard representation of S^3 .
- Hence removing a neighborhood of m and performing 0-surgery on c gives a hyperbolic manifold X on which n -filling yields $S^3 \setminus K'_n$.
- Define $v := \text{vol}(S^3 \setminus K)$, $v'_n := \text{vol}(S^3 \setminus K'_n)$ and $v_X := \text{vol}(X)$.
- Now we use the above bounds to show that:

$$v'_n \geq \left(1 - \left(\frac{2\pi\ell}{|n|\sqrt{3}}\right)^2\right)^{3/2} v_X > v$$

for $|n| > N := \frac{2\pi}{\sqrt{3}} \ell (1 - (v/v_X)^{2/3})^{-1/2}$.

- Hence all integer slopes $|n| > N$ are non-characterizing.
- Finally, we check the finitely many remaining cases by computer.

Ruling Out Integer Characterizing Slopes

- We developed an algorithm that can be carried out by a computer script to rule out the integer characterizing slopes of some knot K , once we have manually produced the link $L = K \cup c$.
- ① Given a link L , run SnapPy commands to find out the volume of the knot K , the volume of the manifold $Z = S_0^3(c)$, and the length of the Seifert longitude in Z .
- ② Using these values, apply the theorems of hyperbolic Dehn surgery to find the bound N such that any integer $|n| > N$ is a characterizing slope for K .
- ③ For the remaining $2N + 1$ cases, verify if the volume of the knot with the same n -surgery as K matches the volume of K .
- We use the *DT code* of the link, which uniquely describes all links that we deal with, up to isotopy.

Theorem (Low-Crossing Knots)

Regarding the integer slopes of knots K such that $c(K) \leq 10$:

- If K has unknotting number $u(K) = 1$ and K is not a twist knot, then K has at most one integer characterizing slope, namely ± 2 .*
- If K is the twist knot 8_1 , then K has at most one integer characterizing slope, namely 0 .*
- If K is one of the $u(K) = 2$ knots $8_4, 8_6, 8_{10}, 8_{12}, 8_{16}, 10_{148}, 10_{149}$, or 10_{150} , K has no possible integer characterizing slope.*
- If K is one of the $u(K) = 2$ knots $8_3, 10_{125}$, or 10_{126} , K has at most one integer characterizing slope.*

Table of Contents

Main theorem

Theorem

If a knot K has unknotting number $u(K) = 1$ and is not a twisted Whitehead double, then K has at most finitely many integer characterizing slopes.

Proposition

Let K be a knot in S^3 . Suppose we can take an unknot c linked with K so that $(0,0)$ -surgery on $K \cup c$ yields S^3 and c is not a meridian of K . Then K has at most finitely many integer characterizing slopes.

- This is a strengthened form of a result proven by Baker-Motegi, which we proved using symmetry of surgical duality.
- Part of the proof of this proposition guarantees that our adapted version of Baker-Motegi construction yields the desired K'_n that shares the same n -surgery as K from purely theoretical grounds of duality.
- We can apply this proposition to prove our main theorem, which admits more concrete conditions on K than the hypothesis in the proposition.

Proof of theorem (cont.)

- Recall that for K with $u(K) = 1$, we could obtain a link $K \cup c$ as in Baker-Motegi with $(0, 0)$ -surgery S^3 by sliding over a Hopf link according to the band presentation for K .
- It remains to show that if K is not a twisted Whitehead double, then c is not a meridian to K after the handle slide.
- In this case, in any band presentation for K , the band must cross the disc bounded by one of the components of the Hopf link.

Lemma

Let $R \cup B$ be a Hopf link, and consider a handle slide of R over B yielding a link L in which R remains a meridian to B . Then there exists a handle slide of R over B , yielding a link isotopic to L , along a band that does not cross either of the discs bounded by R or B .

Extension of theorem

- How can we drop the condition on twisted Whitehead doubles? And on unknotting number?
- Can we produce an algorithm to find links $L = K \cup c$ from handle slides on a Hopf link?

Conjecture

If K is a knot with unknotting number $u(K) = 1$ and K is not a twist knot, then K has at most one integer characterizing slope: ± 2 .

Next Steps

- Rigorously prove the lemma on band presentations.
- Keep experimenting with handle slides in order to expand the list on Theorem 1.2.
- Attempt to use other tools to prove a version of Theorem 1.1 for Twisted Whitehead Doubles.
- Look into our final conjecture.