

INTEGER CHARACTERIZING SLOPES AND UNKNOTTING NUMBERS

GABRIEL AGOSTINI, SOPHIA CHEN, KYLE HAYDEN, CHRISTIAN SERIO,
CECILIA WANG, ANTON WU, AND KEXIN WU

ABSTRACT. [Write one!]

1. INTRODUCTION

[Introduction with some motivation and introduction of terms like “characterizing slopes.” See the papers by Yi and Zhang and by McCoy and by Lackenby on characterizing slopes for ideas to steal. Mention Piccirillo’s results for unknotting number one and using this to show that the Conway knot isn’t slice. Mention that McCoy’s work [McC18], which shows that a hyperbolic knot has only finitely many non-characterizing slopes p/q with $|q| \geq 3$. In a sense, this implies that “most” slopes p/q are characterizing for any given hyperbolic knot K : The probability that a randomly chosen slope p/q is characterizing approaches 1 as $|p| + |q| \rightarrow \infty$.]

Theorem 1.1. *If a knot K has unknotting number $u(K) = 1$ and is not a twisted Whitehead double, then K has at most finitely many integer characterizing slopes.*

We worked our way towards Theorem 1.1 by first manually finding knots K'_0 with the same 0-surgery as a given knot K with $u(K) = 1$ inspired by Piccirillo’s construction and then devising a mechanism to produce all K'_n that shares the same n -surgery with K . Beginning with the former process, we found the following result for knots with low crossing number:

Theorem 1.2. *For knots K with crossing number $c(K) \leq 10$:*

- (a) *If K has unknotting number $u(K) = 1$ and K is not a twist knot, then K has at most one integer characterizing slope, namely ± 2 .*
- (b) *If K is one of the knots 8_4 , 8_6 , 8_{10} , or 8_{12} , then K has $u(K) > 1$ and has no integer characterizing slopes.*
- (c) *If K is the twist knot 8_1 , then K has at most one integer characterizing slope, namely 0.*

This theorem specifies the number of integer characterizing slopes for several knots with $c(K) \leq 10$. It is possible that the bound on $c(K)$ can be increased, especially for knots with unknotting number $u(K) = 1$. Similarly, the list in part (b) can probably be expanded, encompassing every knot that fits Piccirillo’s construction. Part (c) of this theorem also shows that the

assumption in Theorem 1.1 that K is not a twisted Whitehead double is not a necessary condition for the conclusion to hold. Moreover, it was proven by Ozsváth and Szabó 2006, maybe cite that every slope is characterizing for the trefoil and the figure-eight knot. Part (c) shows that this fact does not generalize to twist knots with higher crossing numbers.

Theorem 1.2 affirmatively answers a question posed by Baker and Motegi, who ask if there are knots with crossing number less than eight that have infinitely many non-characterizing slopes [BM18, Question 1.7]. In fact, Theorem 1.2 shows that all knots with crossing number less than eight and unknotting number one which are not twist knots have infinitely many non-characterizing slopes. [See if we can expand this result to be independent of $u(K)$.]

The proof of this theorem relies in part on a computer program to calculate volumes of finitely many surgeries on each knot. We use the software SnapPy, which can triangulate hyperbolic manifolds. The script used is included on maybe put on Appendix?, and the files produced for all knots encompassed can be found on Kyle's website.

1.1. Surgery duals. Let Y be a 3-manifold given by p/q -surgery on a knot $K \subset S^3$. Then Y is obtained by thickening K to a solid torus $N(K)$, removing the interior of $N(K)$ from S^3 , and gluing a solid torus back into $S^3 \setminus \text{int}(N(K))$ in a different manner. Note that K is the core of the solid torus $N(K)$, i.e., K is identified with $S^1 \times \{0\}$ under the diffeomorphism between $N(K)$ and $S^1 \times D^2$. The core of the new solid torus that is glued into $S^3 \setminus \text{int}(N(K))$ specifies a knot γ in Y , which we call the *surgery dual* of K . In a surgery diagram for Y , γ can be represented by a meridian to K . If $K' \subset S^3$ is another knot which yields the same p/q -surgery as K , then one can ask how the surgery dual γ' of K' in Y is related to γ .

Conjecture 1.3 (Baker). *If K and K' are non-isotopic knots in S^3 which yield the same 3-manifold Y under p/q -surgery, then their surgery duals γ and γ' in Y are not homotopic.*

We showed that Baker's conjecture holds for all knots K with unknotting number $u(K) = 1$ and crossing number $c(K) \leq 10$ in the case of zero surgery. The files corresponding to this verification can be found on Kyle's website.

2. KNOTS WITH UNKNOTTING NUMBER ONE

2.1. Banded Hopf link diagrams. A knot K with unknotting number one can be represented by a diagram that consists of a Hopf link with its two components connected by a band. We have the following theorem:

Theorem 2.1. *If K has unknotting number one, then it is obtained from the Hopf link by a single band move.*

To find the banded Hopf link diagram for a knot K with $u(K) = 1$, we first find the unknotting crossing of K and alternate it. We then draw a

meridian adjacent to it and slide K over the meridian. After simplification, we obtain the banded Hopf link diagram for K .

Twisted Whitehead double knots can be defined in various ways. For our purposes, we use a characterization in terms of band presentations.

Definition 2.2. A knot K is a *twisted Whitehead double* if there exists a band presentation for K in which the band does not cross either component of the Hopf link.

We note that twist knots are twisted Whitehead doubles, but not all twisted Whitehead doubles are twist knots.

2.2. Piccirillo's construction. We have the following theorem due to Piccirillo:

Theorem 2.3 (Piccirillo, 2018). *Let L be a three-component link consisting of disjoint components R , B , and G , giving a surgery diagram such that*

- (a) R is a 0-framed unknot,
- (b) B and G have integral framings,
- (c) *If we remove G (resp. B), then R is isotopic to a meridian to B (resp. G).*

Let Y be the 3-manifold given by surgery on L . Then for any $n \in \mathbb{Z}$, there are knots $K, K' \subset S^3$ such that $Y \cong S_n^3(K) \cong S_n^3(K')$.

Thus for K a knot with unknotting number $u(K) = 1$, there exists another knot K'_n (not necessarily distinct from K) with the same n -surgery as K for any integer n .

Inspired by Piccirillo's construction, we can produce some form of the link $K \cup c$ for K a knot with $u(K) = 1$ that would serve as an intermediary between n -surgery on K and explicit K'_n . We start with a banded Hopf link diagram for K . We take a Hopf link with components R and B , both with framings 0, and we handle slide B over R according to the band presentation for K . This produces a two-component link where B becomes K and R a 0-framed unknot c linked with K . We then adjust the framing of B to n to correspond with n -surgery on K . Now add a meridian G to $R = c$ and slide B back over R and we hereby obtain a diagram that fits into Piccirillo's construction, where R remains a 0-framed unknot while B and G are both with integral framings and each would have R as a meridian to it if removing the other.

Overall, we seek an explicit construction that would turn $K \cup c$ into a twist family of knots K'_n with its initial step inspired by Piccirillo's as above. The next step makes use of the following concepts developed by Kouno, Motegi, and Shibuya.

2.3. Twist families of knots. We follow Kouno, Motegi, and Shibuya's treatment of a twisting operation on a knot inside a solid torus. Let K be an unoriented smooth knot in the oriented 3-sphere S^3 and V a solid

torus with a preferred framing that contains K in its interior. Define an orientation-preserving homeomorphism f_n of V such that $f_n(m) = m$ and $f_n(l) = l + nm$ in $H_1(\partial V)$ where (m, l) is a preferred meridian-longitude pair of V . This twisting homeomorphism f_n defines a new knot $f_n(K)$ in S^3 , the result of twisting K inside V . Write $K_{V,n} := f_n(K)$. The wrapping number $w_V(K)$ of K in V is defined to be the minimal intersection number of K with a meridional disc in V .

From a two-component link $K \cup c$ with c an unknot linked with K , we can obtain a new knot $K_n := K_{V,n}$ by taking V to be a solid torus that has preferred meridian c and contains K in its interior. In other words, we obtain K_n from $K \cup c$ by twisting K for n times along c . Note that $w_V(K) \geq 1$ since c is linked with K , and $w_V(K) = 1$ if and only if c is a meridian to K . Moreover, $K_0 = K$. We define the *twist family* with base knot K and unknot c to be the set $\{K_n \mid n \in \mathbb{Z}\}$. Note that for all $m, n \in \mathbb{Z}$, K_m is obtained from K_n by an $(m - n)$ -twist along V [KMS91, Remark 2.2], so any knot K_n can be taken as the base of the twist family.

To construct the knot K'_n from the two component link $K \cup c$, we first get at the diagram $B \cup R \cup G$ by Piccirillo's as in the previous section. Note that the framing of B is changed to n to corresponds to the n -surgery on K and thus by a slam dunk, we can change the framing of B to 0 if we add a meridian P with framing $-1/n$ to B . We slide B over R , and we isotope the diagram until B and R are both unknots that cross each other twice (i.e., B and R form a Hopf link if G is removed). We then slide G over B until G is no longer linked with R . Now we can remove R using a “lightbulb trick” which leaves us a two-component link $G \cup P$, with G as the knot K'_0 that shares the same 0-surgery as K and P as an unknot c' linked with K'_0 . Observe that the framing of P remains unchanged throughout this process and we can therefore do a n -Rolfsen twist on $G \cup P$ to get rid of $P = c'$. As a result, we are left with a knot obtained from twisting K'_0 along c' for n times, which is the knot K'_n as desired.

The knots K'_n for $n \in \mathbb{Z}$ clearly form a twist family with base K'_0 (in fact, any K'_n works) and unknot c' , according to our definition above.

Kouno, Motegi, and Shibuya proved that for a twist family $\{K_{V,n}\}$ in a solid torus V with $w_V(K) \geq 2$, there are at most finitely many integers n_i for which $K_{V,n_i} \sim K$ [KMS91, Theorem 3.2]. Here, $K_{V,n_i} \sim K$ means that there exists a homeomorphism of S^3 carrying K_{V,n_i} to K , which is weaker than the existence of an orientation-preserving homeomorphism of S^3 sending K_{V,n_i} to K . Further, the latter existence is equivalent to the existence of an ambient isotopy between K_{V,n_i} and K [KMS91, Introduction]. Hence, we can translate [KMS91, Theorem 3.2] into the language of the link $K \cup c$ as follows:

Theorem 2.4 (Kouno-Motegi-Shibuya). *Let K be a knot in S^3 and c an unknot linked with K . Let $\{K_n\}$ be the twist family with base knot K and*

unknot c . If c is not a meridian to K , then there are at most finitely many integers n_i such that $K_{n_i} \simeq K$.

Here, the relation \simeq denotes ambient isotopy of knots in S^3 .

2.4. Proof of Theorem 1.1. We show that under the hypotheses on K , $K'_n \simeq K$ for at most finitely many n . If $K'_n \not\simeq K$ for all n , the conclusion holds. If $K'_N \simeq K$ for some N , then we regard K'_N as the base knot of the family $\{K'_n\}$. Note that c is not a meridian to K'_N as long as it is not a meridian to K'_0 . To see this, note that if c' is a meridian to K'_N , then twisting through c' does not change K'_N ; hence by twisting $-N$ times, we see that $K'_N \simeq K'_0$, and c' is a meridian to K'_0 . It remains to show that c' is not a meridian to K'_0 , for then Theorem 2.5 applies to the twist family $\{K'_n\}$ to show that $K'_n \simeq K'_N \simeq K$ for at most finitely many n .

Since K is not a twisted Whitehead double, in any band presentation for K , the band must cross one component of the Hopf link. The following lemma therefore proves that in the link $K \cup c$ appearing in Piccirillo's construction, the unknot c is not a meridian to K .

Lemma 2.5. *Let $R \cup B$ be a Hopf link, and consider a handle slide of R over B which leaves R a meridian to B . Then there is an equivalent handle slide of R over B along a band which does not cross either R or B .*

Proof. TBD □

We now appeal to a lemma of Baker and Motegi [BM18, Lemma 2.4], with notation adapted:

Lemma 2.6 (Baker-Motegi, 2018). *Let $K'_0 \cup c'$ be a two-component link in S^3 such that c' is a meridian of K'_0 . Then $(0,0)$ -surgery on $K'_0 \cup c'$ results in S^3 with its surgery dual link $c \cup K$, for which c is a meridian to K .*

Note that in our context, $(0,0)$ -surgery on $K'_0 \cup c'$ always results in S^3 with its surgery dual link $c \cup K$. We can thus simplify the lemma as: c' being a meridian of K'_0 implies that c is a meridian to K . Hence, we obtain the following corollary as the contrapositive of the simplified lemma.

Corollary 2.7. *If c is not a meridian to K , then c' is not a meridian to K'_0 , with K , c , K'_0 and c' as above.*

Proof. We know that c' and K'_0 are surgery duals to K and c respectively. Show that it follows that c and K are dual to K'_0 and c' respectively. □

This completes the proof of the theorem.

3. KNOTS WITH LOW CROSSING NUMBER

3.1. Hyperbolic Dehn surgery. [Recall and discuss the relevant theorems used in our approach.]

3.2. Proof of Theorem 1.2. To prove Theorem 1.2, we need to rule out infinitely many characterizing slopes for each of the knots K we are interested in. In general, this is a two-part algorithm:

- (1) We use the theorems of hyperbolic Dehn surgery to find a bound $N \in \mathbb{Z}_{>0}$ such that any slope $n \in \mathbb{Z}$ is not characterizing if $|n| > N$.
- (2) For the $2N + 1$ remaining cases, we directly examine a knot K'_n with the same n -surgery as K . If $K'_n \neq K$, we know that n is noncharacterizing.

[First part, based on hyperbolic Dehn surgery theorems. Explain how we found N .]

Finding this bound and examining the volumes of each of $2N + 1$ knots in the twist family of each knot K is a tedious computational problem. It was convenient for us to let a computer program run this proof given minimal input.

3.2.1. Code and Data. We produced a framed link $L = K \cup R \cup c$ for each of the knots K , where R and c are unknotted components and c is a meridian to R . This link is such that, if the framing on R is zero and the framings on K and c are any integer numbers, an equivalent surgery diagram to L is a link $L' = B \cup R \cup G$ that satisfies Piccirillo's theorem, with K turning into B and c turning into G .

Any link can be described by a partitioned sequence of even numbers called its Dowker-Thistlethwaite code, or DT code. This is provided by the software SnapPy. In the case of a framed link L fitting the description above, we use the following lemma:

Lemma 3.1. *Up to isotopy, L can be uniquely described by its DT code.*

Proof. This lemma is a direct application of Doll and Hoste's Theorem 1.2 [CITE] to our link L . As defined, L will always be a nonsplit link whose orientation is irrelevant for the purposes of integral surgery. R and c are two unknots, so L is a prime link if and only if K is a prime knot. That is the single case we are interested in, since only prime knots are classified along with their invariants and diagrams. Thus, by Doll and Hoste, minimal projections of L are in one-to-one correspondence with valid DT codes. Finally, there is a single solution for the question of which of the components is K , R , and c . We use that c is strictly a meridian to R , so its only crossings are the two they share. Since the entries on the DT code come from crossings either within a single component or between two different ones, we notice that the code corresponding to the component c will have length 1 (since, by definition of a DT code, only even labeled crossings appear). R also does not have crossings with itself, but it has crossings with R and c , so its code has length greater than 1. Moreover, since K is not an unknot, $c(K) \geq 3$, so the code corresponding to K will always be longer than the code corresponding to R . Thus, it is clear that $l(K) > l(R) > l(c) = 1$, where $l(X)$ is the code length of the component X . \square

The script `integral_slopes.py`, given the DT code of a link L satisfying the conditions above, outputs a list of possible integral characteristic slopes for any knot K . It relies on writing a text file, running it on SnapPy, saving the output, and interpreting it repeatedly. There are certain system requirements explicit on the code. The script consists of the following steps:

- (1) From the DT code, the program writes the file `K [INFO].py`. This file asks for the volume of the knot K , for the volume of the manifold $Z = S_0^3(R)$ obtained by zero-surgery filling the 1-handle component, and for the length of the Seifert longitude in Z .
- (2) SnapPy opens, runs, and saves this file, producing a first output file named `[OUT] K [INFO].py`.
- (3) The program extracts this information and calculates the bound N using the theorems of hyperbolic Dehn surgery.
- (4) The program writes the file `K [TEST].py`. This file asks for the volume of each knot K'_n in the twist family of K , whose exterior is easily obtained by doing n -surgery on the first component of the manifold Z , for all n such that $|n| \leq N$.
- (5) SnapPy opens, runs, and saves this file, producing a second output file named `[OUT] K [TEST].py`.
- (6) The program verifies if this last file contains all $2N + 1$ cases. This step is included because the run time in SnapPy is sometimes unpredictable.
- (7) The program identifies the knots whose volume is less than or equal (up to approximations) to the volume of the knot K . Those cases are singled out in the output file and, if possible, identified by SnapPy.

All files corresponding to the knots listed in Theorem 1.2 can be found on [Kyle's website](#). These files prove the theorem.

Proof. We produced the link L aforementioned for all knots K mentioned in Theorem 1.2. All the DT codes are included in the file `DT_List.txt`, which is correctly formatted for the script `integral_slopes.py`. Running it outputs the `.py` files labeled `[OUT]`, and text that is reproduced in the file `output.txt`. From reading this file, we see that:

- (a) For all knots K of crossing number at most 10 with unknotting number $u(K) = 1$ and not twist knots, the program indicates that only $n = 2$ or $n = -2$ could be characterizing.
- (b) For K one of the knots $8_4, 8_6, 8_{10}$, or 8_{12} , the program indicates that all knots K'_n have different volume than K , so no integral slopes can be characterizing.
- (c) For the knot 8_1 , the program indicates that only $n = 0$ could be characterizing.

□

3.3. Possible extensions of Theorem 1.1. [Note that $u(K) = 1$ is not necessary for Piccirillo's construction. We only need to be able to produce

a link $R \cup B$ (with $B \simeq K$ and R an unknot) from handle slides on a Hopf link. Explain this procedure for knots with $u(K) > 1$.]

4. ADDITIONAL RESULTS

[Any additional findings, including HFK or Khovanov homology findings, or enhanced sliceness obstructions using Piccirillo’s technique, can go here.]

REFERENCES

- [BM18] Kenneth L. Baker and Kimihiko Motegi. Noncharacterizing slopes for hyperbolic knots. *Algebr. Geom. Topol.*, 18(3):1461–1480, 2018.
- [KMS91] Masaharu Kouno, Kimihiko Motegi, and Tetsuo Shibuya. Twisting and knot types. *Journal of the Mathematical Society of Japan*, 44(2):199–216, March 1991.
- [McC18] Duncan McCoy. On the characterising slopes of hyperbolic knots. Available as arXiv:1807.11099, August 2018.