
Cell Metabolism

Resource

A High-Density Human Mitochondrial Proximity Interaction Network

Graphical Abstract



Highlights

- We created a high-resolution human mitochondrial protein proximity map using BioID
- Bait-bait analysis shows that the map has sub-compartment resolution
- Prey correlation analysis identifies functional clusters and specific modules
- OMM baits show specific interactions reflecting contact sites and dual localization

Authors

Hana Antonicka, Zhen-Yuan Lin, Alexandre Janer, Mari J. Aaltonen, Woranontee Weraarpachai, Anne-Claude Gingras, Eric A. Shoubridge

Correspondence

gingras@lunenfeld.ca (A.-C.G.), eric.shoubridge@mcgill.ca (E.A.S.)

In Brief

Antonicka et al. used BioID, a proximity-dependent biotinylation assay, to create a high-density mitochondrial proximity interaction map. They were able to identify about half of the mitochondrial proteome, assign proteins to functional clusters and specific modules, show dual protein localization, and demonstrate specific interactions with other cellular organelles.

Antonicka et al., 2020, Cell Metabolism *32*, 479–497
September 1, 2020 © 2020 Elsevier Inc.
https://doi.org/10.1016/j.cmet.2020.07.017

CellPress

Cell Metabolism, 32 (2020) 479-506. doi:10.1016/j.cmet.2020.07.017

Cell Metabolism, 32 (2020) 479-506. doi:10.1016/j.cmet.2020.07.017

Cell Metabolism, 32 (2020) 479-506. doi:10.1016/j.cmet.2020.07.017

Cell Metabolism, 32 (2020) 479-506. doi:10.1016/j.cmet.2020.07.017

Cell Metabolism, 32 (2020) 479-506. doi:10.1016/j.cmet.2020.07.017

Cell Metabolism, 32 (2020) 479-506. doi:10.1016/j.cmet.2020.07.017

Cell Metabolism, 32 (2020) 479-506. doi:10.1016/j.cmet.2020.07.017

Cell Metabolism, 32 (2020) 479-506. doi:10.1016/j.cmet.2020.07.017

Cell Metabolism, 32 (2020) 479-506. doi:10.1016/j.cmet.2020.07.017

Cell Metabolism, 32 (2020) 479-506. doi:10.1016/j.cmet.2020.07.017

Cell Metabolism, 32 (2020) 479-506. doi:10.1016/j.cmet.2020.07.017

Cell Metabolism, 32 (2020) 479-506. doi:10.1016/j.cmet.2020.07.017

Cell Metabolism, 32 (2020) 479-506. doi:10.1016/j.cmet.2020.07.017

Cell Metabolism, 32 (2020) 479-506. doi:10.1016/j.cmet.2020.07.017

Cell Metabolism, 32 (2020) 479-506. doi:10.1016/j.cmet.2020.07.017

Cell Metabolism, 32 (2020) 479-506. doi:10.1016/j.cmet.2020.07.017

Cell Metabolism, 32 (2020) 479-506. doi:10.1016/j.cmet.2020.07.017

Cell Metabolism, 32 (2020) 479-506. doi:10.1016/j.cmet.2020.07.017

Cell Metabolism, 32 (2020) 479-506. doi:10.1016/j.cmet.2020.07.017

Cell Metabolism, 32 (2020) 479-506. doi:10.1016/j.cmet.2020.07.017

Cell Metabolism, 32 (2020) 479-506. doi:10.1016/j.cmet.2020.07.017

Cell Metabolism, 32 (2020) 479-506. doi:10.1016/j.cmet.2020.07.017

Cell Metabolism, 32 (2020) 479-506. doi:10.1016/j.cmet.2020.07.017

Cell Metabolism, 32 (2020) 479-506. doi:10.1016/j.cmet.2020.07.017

Cell Metabolism, 32 (2020) 479-506. doi:10.1016/j.cmet.2020.07.017

Cell Metabolism, 32 (2020) 479-506. doi:10.1016/j.cmet.2020.07.017

Cell Metabolism, 32 (2020) 479-506. doi:10.1016/j.cmet.2020.07.017

Cell Metabolism, 32 (2020) 479-506. doi:10.1016/j.cmet.2020.07.017