

$$1) f_0 = e^{-t^2/2} \quad f_1 = te^{-t^2/2} \quad f_2 = t^2 e^{-t^2/2} \quad \text{in } L_2; \quad \langle f, g \rangle = \int_{-\infty}^{\infty} f(t)g(t)dt$$

$$\int_{-\infty}^{\infty} e^{-t^2/2} dt = \sqrt{2\pi}$$

$$1. g_0 = f_0$$

$$\|g_0\|^2 = \langle f_0, f_0 \rangle = \int_{-\infty}^{\infty} e^{-t^2} dt = \sqrt{\pi}$$

$$2. g_1 = f_1 - \frac{\langle f_1, g_0 \rangle}{\langle g_0, g_0 \rangle} g_0$$

$$\langle f_1, g_0 \rangle = \int_{-\infty}^{\infty} te^{-t^2} dt = 0 \quad \text{te}^{-t^2} \text{ is an odd function, so}$$

$$g_1 = f_1$$

$$\|g_1\|^2 = \langle f_1, f_1 \rangle = \int_{-\infty}^{\infty} t^2 e^{-t^2} dt$$

$$\int_{-\infty}^{\infty} t^2 e^{-t^2} dt = \sqrt{\pi}/2$$

$$3. g_2 = f_2 - \frac{\langle f_2, g_0 \rangle}{\langle g_0, g_0 \rangle} g_0 - \frac{\langle f_2, g_1 \rangle}{\langle g_1, g_1 \rangle} g_1$$

$$\langle f_2, g_1 \rangle = \int_{-\infty}^{\infty} t^3 e^{-t^2} dt = 0, \text{ odd function}$$

$$\langle f_2, g_0 \rangle = \int_{-\infty}^{\infty} t^2 e^{-t^2} dt = \sqrt{\pi}/2$$

$$\langle g_0, g_0 \rangle = \sqrt{\pi}$$

$$g_2 = f_2 - \frac{\sqrt{\pi}/2}{\sqrt{\pi}} f_0 = t^2 e^{-t^2/2} - \frac{1}{2} e^{-t^2/2} = e^{-t^2/2} (t^2 - \frac{1}{2})$$

$$\|g_2\|^2 = \langle g_2, g_2 \rangle = \int_{-\infty}^{\infty} (t^2 - \frac{1}{2})^2 e^{-t^2} dt$$

$$\begin{aligned}
 \text{1) cont } \|g_2\|^2 &= \int t^4 e^{-t^2} dt - \int t^2 \frac{d^2}{dt^2} \int e^{-t^2} dt + \frac{1}{4} \int e^{-t^2} dt \\
 &= \frac{3\sqrt{\pi}}{4} - \frac{\sqrt{\pi}}{2} + \frac{\sqrt{\pi}}{4} = \sqrt{\pi}/2
 \end{aligned}$$

so an orthogonal set is

$$e^{-t^2/2}, t e^{-t^2/2}, (t^2 - \frac{1}{2}) e^{-t^2/2}$$

$$2) x(t) = \sin(4t) \quad L_2[0,1]$$

$$\langle f(t), g(t) \rangle = \int_0^1 f(t)g(t) dt \quad \text{in this space}$$

$$2.1) \text{ span } \{t, t^2\}$$

$$\begin{pmatrix} \langle t, t \rangle & \langle t, t^2 \rangle \\ \langle t^2, t \rangle & \langle t^2, t^2 \rangle \end{pmatrix} \begin{pmatrix} c_1 \\ c_2 \end{pmatrix} = \begin{pmatrix} \langle x, t \rangle \\ \langle x, t^2 \rangle \end{pmatrix}$$

$$= \underbrace{\begin{pmatrix} 1/3 & 1/4 \\ 1/4 & 1/5 \end{pmatrix}}_A \begin{pmatrix} c_1 \\ c_2 \end{pmatrix} = \begin{pmatrix} \int_0^1 t \sin(4t) dt \\ \int_0^1 t^2 \sin(4t) dt \end{pmatrix}$$

$$\int_0^1 t \sin(4t) dt = -\frac{1}{4} t \cos(4t) + \frac{1}{16} \sin(4t) \Big|_0^1$$

$$= -\frac{1}{4} \cos(4) + \frac{1}{16} \sin(4)$$

$$\int_0^1 t^2 \sin(4t) dt = \left(\frac{2}{4^3} - \frac{t^2}{4} \right) \cos(4t) + \frac{2t}{4^2} \sin(4t) \Big|_0^1$$

$$= \left(\frac{2}{64} - \frac{1}{4} \right) \cos(4) + \frac{2}{16} \sin(4) - \left(\frac{2}{4^3} \right)$$

$$= -7/32 \cos 4 + \frac{1}{8} \sin(4) - 1/32$$

$$A = \begin{pmatrix} 1/3 & 1/4 \\ 1/4 & 1/5 \end{pmatrix} \quad \det(A) = \frac{1}{15} - \frac{1}{16} = 1/240$$

$$A^{-1} = 240 \begin{pmatrix} 1/5 & -1/4 \\ -1/4 & 1/3 \end{pmatrix}$$

$$\begin{pmatrix} c_1 \\ c_2 \end{pmatrix} = 240 \begin{pmatrix} 1/5 & -1/4 \\ -1/4 & 1/3 \end{pmatrix} \begin{pmatrix} -\frac{1}{4} \cos 4 + \frac{1}{16} \sin 4 \\ -\frac{7}{32} \cos 4 + \frac{1}{8} \sin 4 - \frac{1}{32} \end{pmatrix}$$

$$C_1 = \frac{9}{8} \cos 4 - \frac{9}{2} \sin 4 + \frac{15}{8} \approx 4.5448$$

$$C_2 = -\frac{5}{2} \cos 4 + \frac{25}{4} \sin 4 - \frac{5}{2} \approx -5.5959$$

projection:

$$p(t) = 4.5448 t - 5.5959 t^2$$

2.2: $t, t^2, \tan t$

$$\langle t, t \rangle = 1/3 \quad \langle t, t^2 \rangle = 1/4 \quad \langle t^2, t^2 \rangle = 1/5$$

$$\langle t, \tan t \rangle = \int_0^1 t \tan(t) dt \quad \langle t^2, \tan(t) \rangle = \int_0^1 t^2 \tan(t) dt$$

$$\langle \tan(t), \tan(t) \rangle = \int_0^1 \tan^2(t) dt = \tan(1) - 1$$


Harder integrals that have to be solved with a Fourier series

↳ also, taking inverse of 3×3 will be worse, mentioning a move to solving using python

→ see appended code:

$$p(t) = C_1 t + C_2 t^2 + C_3 \tan(t)$$

$$p(t) = 2.489 t - 7.73 t^2 + 2.845 \tan(t)$$



3) $Ly = y'' = \lambda y \quad y(0) = y(1) = 0 \text{ on } [0,1]$

3 sign cases:

1. $\lambda = 0$

$$y'' = 0 \quad y' = A \quad y = Ax + B$$

$$0 = B$$

$$0 = A$$

> trivial solution
 $y = 0$, no eigenfunction

2. $\lambda > 0$

$$\lambda = a^2$$

$$y = Ae^{ax} + Be^{-ax}$$

$$0 = A + B \quad A = -B$$

$$0 = Ae^1 - Ae^{-1} \Rightarrow A = 0, B = 0$$

Again $y = 0$, no eigenfunction

3. $\lambda < 0$

$$\lambda = -k^2, \quad k > 0,$$

$$y = A \sin kx + B \cos kx$$

$$0 = B$$

$$0 = A \sin k \quad \text{either } A = 0 \text{ or } \sin k = 0$$

$$\sin k = 0$$

$$k = 0, \pi, 2\pi, \dots$$

$$k = n\pi \quad n = 1, 2, \dots$$

$$\lambda_n = -(n\pi)^2$$

λ_n eigenvalues

$$y_n(x) = \sin(n\pi x) \quad n = 1, 2, \dots$$

y_n eigenfunctions

$$4) \quad LT : -k \left(\frac{1}{r} \frac{\partial}{\partial r} \left(r \frac{\partial T}{\partial r} \right) + \frac{\partial^2 T}{\partial z^2} \right) = 0$$

$$\Rightarrow -k T_r(R, z) = \beta T(R, z)$$

Galerkin:

$$\hat{T}(r, z) = T_1(z) \phi_1(r) + T_2(z) \phi_2(r) \quad \phi_1 = 1$$

$$\langle u, v \rangle = \int_0^R r u v dr$$

$$\psi_n = \phi_m (m=1, 2)$$

$$\phi_2 = 2 \frac{r^2}{R^2} - 1$$

$$\langle \phi_m, L \hat{T} \rangle = 0 \quad m=1, 2 \quad (\text{enforce this condition}) \quad (*)$$

T_1, T_2 dependent on just z :

$$\hat{T}_r = T_2(z) \phi_2'(r) \quad \hat{T}_{zz} = T_1''(z) \phi_1(r) + T_2''(z) \phi_2(r)$$

$$\langle \phi_m, -k \frac{1}{r} \partial_r (r \hat{T}_r) \rangle = -k \int_0^R \phi_m \partial_r (r \hat{T}_r) dr$$

IBP:

$$= -k [\phi_m r \hat{T}_r]_0^R + k \int_0^R \phi_m' r \hat{T}_r dr$$

At $r=R$, we have $-k \hat{T}_r = \beta \hat{T}$ and $\phi_1(R) = \phi_2(R) \Rightarrow$
 $\hat{T}(R, z) = T_1 + T_2$ from $(*)$

$$\Rightarrow -k [\phi_m r \hat{T}_r]_0^R = \beta R \phi_m(R) (T_1 + T_2)$$

$$= \beta R (T_1 + T_2)$$

$$\hat{T}_r = T_2 \phi_2' \Rightarrow$$

$$k \int_0^R \phi_m' r \hat{T}_r dr = k T_2 \int_0^R r \phi_m' \phi_2' dr$$

From \hat{T}_{zz} , we have

$$\langle \phi_m, -k \hat{T}_{zz} \rangle = -k T_1'' \int_0^R r \phi_m \phi_1 dr - k T_2'' \int_0^R r \phi_m \phi_2 dr$$

$$4) \text{ cont. } M_{11} = \int_0^R r \phi_1 \phi_1 dr = R^2/2$$

$$M_{12} = M_{21} = \int_0^R r \phi_1 \phi_2 dr = 0 \quad (\text{orthogonal } \phi_1, \phi_2)$$

$$M_{22} = \int_0^R r \phi_2^2 dr = R^2/6$$

$$A_1 = \int_0^R r \phi_1' \phi_2' dr = 0$$

$$A_2 = \int_0^R r \phi_2' \phi_2' dr = \int_0^R r \left(\frac{4r}{R^2} \right)^2 dr = 4$$

Galerkin ODE's:

$$m=1$$

$$\beta R(T_1 + T_2) + kT_2 A_1 - kT_1'' M_{11} - kT_2'' M_{12}$$

$$\Rightarrow -\frac{kR^2}{2} T_1'' + \beta R(T_1 + T_2) = 0$$

$$m=2$$

$$\beta R(T_1 + T_2) + kT_2 A_2 - kT_1'' M_{21} - kT_2'' M_{22} = 0$$

$$-\frac{kR^2}{6} T_2'' + 4kT_2 + \beta R(T_1 + T_2) = 0$$