

Technological Civilization Collapse–Recovery Dynamics and Detectability

Celia Blanco^{1,2}

1. Centro de Astrobiología (CSIC-INTA),
Ctra de Torrejón a Ajalvir, km 4, 28850 Torrejón de Ardoz, Madrid, Spain

2. Blue Marble Space Institute of Science, Seattle, WA 98104 USA

April 10, 2026

1 Abstract

How long a technological civilization remains active, and what determines whether it collapses or persists, is a central question for both projecting humanity’s future and assessing the prevalence of detectable intelligence in the galaxy. We model collapse-recovery dynamics across ten plausible futures for Earth-originating civilization using a hybrid deterministic-stochastic simulation over a 1000-year window. Duty cycles range from ~ 0.38 to 1.00, with trajectory outcomes shaped by the interplay of governance structure, resource pressure, and hazard exposure. Several model parameters map onto actionable resilience levers, and modest improvements can qualitatively alter long-term trajectories. We discuss implications for Earth’s civilizational resilience and for the search for extraterrestrial technosignatures.

2 Introduction

One of the most puzzling questions in astrobiology is whether intelligent civilizations exist elsewhere in the galaxy, and if so, why we have not yet detected them. This apparent contradiction, often referred to as the Fermi Paradox or the Great Silence, raises the question of why, in a galaxy billions of years old, we observe no clear signs of advanced extraterrestrial life. Many solutions have been proposed to explain this Great Silence [Webb, 2010]. One class of explanations suggests that intelligent life is intrinsically rare, perhaps requiring an improbable convergence of planetary, chemical, or biological conditions [Wesson, 1990]. Another contends that while intelligence may emerge, technological civilizations are typically short-lived, succumbing to self-destruction through ecological collapse, conflict, or other systemic instabilities [Vinn, 2024].

The second class of explanation raises a question that is critical regardless of how common the emergence of intelligence may be: how long do technological civilizations last once they arise? Even if intelligent life is widespread, the number of civilizations detectable at any given time depends decisively on their longevity [Balbi and Ćirković, 2021]. This question is also central to projecting the long-term trajectory of our own civilization. Most treatments of longevity, however, assume that once a society becomes technologically active, it remains so continuously until it either collapses

or deliberately goes silent. But this assumption is difficult to justify. Historical studies of Earth’s civilizations reveal cycles of rise and fall with typical durations on the order of centuries [Shermer, 2002], and even within a single civilization’s lifespan, the temporal distribution of active periods may matter as much as their cumulative duration [Balbi, 2018]. Therefore, understanding civilizational longevity requires moving beyond static estimates and examining the temporal dynamics of technological activity itself.

There are good reasons to expect that civilizational collapse-recovery dynamics are complex. It has been argued that planetary civilizations face a fundamental bifurcation: either “asymptotic burnout” in which superlinear growth leads to singularity-driven collapse, or “homeostatic awakening”, in which a civilization deliberately curbs expansion to achieve long-term equilibrium [Wong and Bartlett, 2022]. Sustainability constraints make exponential expansion highly unlikely, so advanced societies may stabilize or collapse before spreading across the Galaxy [Haqq-Misra and Baum, 2009]. If collapse is common and recovery uncertain, then the technosphere (the totality of a civilization’s technological infrastructure and output) may best be understood not as a permanent state but as an intermittent phenomenon, alternating between periods of activity and dormancy in a pattern shaped by resource availability, governance structure, and exposure to existential risk.

Technological intermittency can be quantified through a civilization’s *duty cycle*, defined as the fraction of its total lifespan during which it is technologically active. The duty cycle captures a reality that static longevity estimates miss—that a civilization persisting for a thousand years may spend substantial portions of that time in dormancy, recovery, or regression. In the context of the search for extraterrestrial intelligence, the relevance of intermittent activity is already recognized. Interstellar signals may be intermittent due to power constraints, planetary rotation, or targeted sequential transmissions [Gray, 2020], and recent work applying Lindy’s law to technosignature longevities suggests that technologically active periods are more likely to be short-lived than long-lived [Balbi and Grimaldi, 2024]. But the implications of intermittency extend well beyond detectability. For our own civilization, understanding the dynamics of collapse and recovery, and how governance, resources, and resilience infrastructure shape the duty cycle, is directly relevant to long-term planning and existential risk mitigation.

In this paper, we develop a framework for modeling these dynamics. Building on ten plausible futures for Earth-originating civilization spanning 1000 years [Haqq-Misra et al., 2025a], we simulate collapse-recovery cycles across diverse sociotechnical trajectories and compute duty cycles for each scenario. We then derive an effective detectability duration, L_{eff} , and explore the implications for Earth’s own civilizational resilience, for the Drake Equation’s longevity term L , and for the broader search for technosignatures.

3 Model Description

A recently proposed framework outlines ten plausible futures for Earth-originating civilization 1000 years from now, structured around variations in governance type, resource regime, and systemic resilience [Haqq-Misra et al., 2025a]. These scenarios are grounded in Earth’s present-day sociotechnical conditions and extrapolated using techniques from futures studies, including PEST analysis (Political, Economic, Social, Technological), cross-consistency assessment, and structured scenario worldbuilding. Building on this qualitative scenario set, we implement a hybrid deterministic-stochastic simulation to represent the temporal dynamics of growth, collapse, and recovery.

3.1 Simulation framework

The simulation spans a fixed period of 1000 years, discretized into yearly time steps. At each step, a civilization is characterized by its technological capacity, $T(t)$, and its available resource stock, $R(t)$. Technological capacity increases linearly each year according to a fixed growth rate r , which we take directly from the values reported in Haqq-Misra et al., Table 9. Simultaneously, the available resource stock is depleted at a constant rate δ . If the resource level falls to zero (i.e., $R(t) < 0$), or if an existential risk event occurs (represented as a probabilistic hazard drawn at each time step) the civilization undergoes a collapse. Collapse represents a systemic breakdown in infrastructure, coordination, and technological continuity. It leads to a sharp reduction in technological capacity, determined by a collapse factor c_f , which defines the proportion of technology that survives the event. This loss can be interpreted as the result of disruptions to power grids, communication networks, knowledge retention, or the abandonment of complex systems.

Following collapse, the system enters a dormant recovery period of duration r_d . During this phase, the society is unable to grow or consume resources, reflecting a temporary halt in technological and economic activity. This delay allows time for institutional reorganization, population stabilization, or environmental recovery. Once the recovery period concludes, the resource stock is partially replenished to a fraction r_f of its original value R_0 . This recovery fraction represents retained infrastructure, stored reserves, or ecological regeneration. A full recovery ($r_f \approx 1$) implies robust contingency planning and strong institutional memory, while lower values indicate more severe losses and diminished rebuilding capacity. This cycle of growth, collapse, dormancy, and recovery can repeat multiple times during the simulation window, depending on the fragility of the system, the frequency of stochastic hazards, and the depletion rate of resources.

Formally, let the initial conditions be $T_0 = 1$, and $R_0 \in \mathbb{R}_+$, and define the indicator function as $\chi(t) \in \{0, 1\}$, with $\chi(t) = 1$ if the system is active (not collapsed or recovering) at time t , and $\chi(t) = 0$ otherwise. The update rules for each time step are given by

$$\begin{aligned} T_{t+1} &= \chi_t (T_t + r) + (1 - \chi_t) T_t, \\ R_{t+1} &= \chi_t (R_t - \delta) + (1 - \chi_t) R_t. \end{aligned} \tag{1}$$

We include both a deterministic and a stochastic trigger for collapse to reflect two distinct failure modes: internal depletion and external shocks. Collapse is deterministically triggered when the resource stock is fully depleted (i.e., $R(t) \leq 0$). It may also be triggered by a stochastic existential hazard, which we model as a Bernoulli process. At each time step t , a uniform deviate $U(t) \sim U(0, 1)$ is drawn. If $U(t) < h$, where $h \in [0, 1]$ is the hazard rate specific to the scenario, then a collapse is triggered independently of resource levels.

Upon collapse, the system state is reset according to

$$\begin{aligned} T_{t+1} &= c_f T_t, \\ R_{t+1} &= 0. \end{aligned} \tag{2}$$

The system then remains inactive for r_d time steps, after which resource availability is restored as

$$R_{t+r_d} = r_f R_0. \tag{3}$$

Together, Equations (1)–(3) define the full dynamical cycle of growth, collapse, and recovery in the model.

We define a civilization as being ON when it maintains a positive level of resources (i.e., $R(t) > 0$). This assumption reflects the idea that some minimal material throughput is necessary for societal function and detectability, including energy consumption, communication, and technosignature production. Conversely, when resources are fully depleted ($R(t) = 0$), the civilization is considered OFF, representing a dormant, collapsed, or otherwise undetectable state. This binary definition underlies the calculation of duty cycle and related metrics discussed in subsection 3.3.

3.2 Scenario typologies and parameter mapping

We apply our model to the ten future scenarios to simulate the long-term behavior of civilizations under diverse sociotechnical conditions. Each scenario describes a distinct vision for humanity’s trajectory, defined by combinations of resource abundance, governance structure, institutional resilience, and technological integration. Our goal is to translate these qualitative narratives into a coherent set of quantitative parameters that govern dynamics of growth, collapse, and recovery. Scenarios are characterized along three conceptual axes: resource regime (scarcity vs. non-scarcity), governance type (from autocratic to participatory), and structural design (hierarchical vs. distributed). Scenarios are also mapped onto six thematic clusters (Table 1).

We assign parameter values based on the sociotechnical features of each scenario. The growth rate r is taken directly from Haqq-Misra et al. (2025) and held fixed. We note that assuming technological capacity increases linearly each year via r is a simplification, especially given scenarios that imply discontinuities such as collapse points or radical transformation points; we retain it here for consistency with the source parameterization. The remaining parameters (initial resource stock R_0 , per-year depletion rate δ , collapse depth c_f , recovery delay r_d , recovery fraction r_f , and existential hazard rate h) are derived from the narrative structure of each scenario and converted into quantitative values through structured reasoning.

We assume that governance structure influences how technological civilizations respond to systemic stress and collapse. Highly centralized systems (rule by one) may retain moderate technological capacity through individual collapse events (higher c_f) due to concentrated infrastructure, and may recover relatively quickly (lower r_d) through top-down coordination. However, their rigidity and concentration of decision-making can make them more susceptible to repeated collapses when combined with resource scarcity or high external hazard exposure, consistent with the observation that increasing sociopolitical complexity yields diminishing returns, ultimately rendering centralized systems fragile [Tainter, 1988]. Oligarchic systems (rule by few) may experience deeper individual collapses (lower c_f) and slower recovery (higher r_d), reflecting fragmented coordination and weaker institutional memory. More distributed governance (rule by all) supports greater redundancy and flexibility, enabling a wider range of outcomes, from near-complete resilience in post-scarcity conditions to moderate collapse depth with relatively rapid recovery. This is consistent with the finding that polycentric governance systems mitigate the risk of institutional failure through overlapping jurisdictions and redundant decision-making structures [Ostrom, 2010]. To reflect these hypothesized patterns, we selected representative parameter value ranges for collapse depth c_f , recovery delay r_d , and recovery fraction r_f that align with the governance structure assumed in each scenario (Table 2). These values are not based on empirical observation, but represent internally consistent assumptions designed to explore contrasting trajectories in our scenario-based simulations.

One exception is the Restoration scenario (S7). While it is categorized as non-scarcity and governed by distributed institutions, it is conceptually framed as a rebuilding civilization emerging

Table 1: Scenario descriptions and sociotechnical tags

Scenario	Tags	Description
S1: Big Brother is Watching	Scarcity + Rule by One + Hierarchical Cluster 6 (Earth 2024)	Centralized authoritarian system under resource stress
S2: Wild West	Scarcity + Rule by Few + Hierarchical Cluster 6 (Earth 2024)	Deregulated, competitive, and fragmented society
S3: Golden Age	Non-scarcity + Rule by All + Distributed Cluster 6 (Earth 2024) [†]	Stable egalitarian society with abundant resources
S4: Living with the Land	Scarcity + Rule by Few + Hierarchical Cluster 5 (Simplicity)	Low-impact, cyclical, ecologically constrained society
S5: Transhumanism	Non-scarcity + Rule by All + Distributed Cluster 4 (Engineering)	Technologically integrated, post-biological civilization
S6: Sword of Damocles	Scarcity + Rule by Few + Hierarchical Cluster 4 (Engineering)	High-tech and high-risk system with cascading failure potential
S7: Restoration	Non-scarcity [‡] + Rule by All + Distributed Cluster 3 (Sustainability)	Recovered post-collapse world with resilience
S8: Ouroboros	Scarcity + Rule by Few + Hierarchical Cluster 3 (Sustainability)	Oscillatory civilization with cyclical collapse and regrowth
S9: Deus Ex Machina	Non-scarcity + Rule by All + Distributed Cluster 2 (Tech Escape)	Aggressive technological expansion with instability risk
S10: Out of Eden	Non-scarcity + Rule by All + Distributed Cluster 1 (Parallel Life)	Harmonized coexistence with nature and machines

[†]Cluster assignments follow [Haqq-Misra et al. \[2025a\]](#); S3 shares the Earth 2024 technology cluster with S1 and S2 despite differing in global factors.

[‡]S7 is tagged non-scarcity in the original framework but parameterized with scarcity-level resource pressure to model post-collapse recovery (see text).

from an earlier collapse. To reflect this, we initialize the simulation with a reduced effective resource stock and a moderate depletion rate (parameters that would otherwise suggest scarcity), thereby modeling a society recovering from prior systemic failure. This allows us to represent S7 as a post-collapse recovery regime without forcing a collapse to occur within the simulation window.

The Ouroboros scenario (S8) is treated differently. In the original framework, S8 is described as a civilization that repeatedly undergoes expansion, collapse, and regrowth, with the year 3000 situated partway through an ongoing recovery phase. Rather than imposing externally timed or periodic collapse events, we model this behavior as an emergent dynamical regime. Specifically, S8 is assigned moderate technological growth, partial technological survival during collapse, and a relatively high recovery fraction of planetary resources. This combination allows repeated collapse–recovery cycles to arise endogenously from the interaction between resource depletion and rebuilding, and permits exploration of whether oscillatory behavior remains stable or dampens over long timescales.

Similarly, Deus Ex Machina (S9) is tagged as non-scarcity in the original framework but is parameterized with moderate resource pressure ($R_0 = 900$, $\delta = 1.3$) to reflect the high extraction demands of its aggressive technological expansion program, which drives resource depletion despite the scenario’s nominally post-scarcity classification.

Table 2: Mapping sociotechnical structure to parameter logic

Governance Type	Collapse Depth c_f	Recovery Delay r_d	Recovery Fraction r_f
Rule by One	0.5–0.7	40–60	0.10–0.20
Rule by Few	0.2–0.6	50–100	0.10–0.35
Rule by All*	0.3–1.0	0–70	0.20–1.0

*Post-scarcity idealizations (S3, S10) use $c_f = 1.0$, $r_d = 0$, $r_f = 1.0$, reflecting no-collapse conditions. S7 and S9 use parameters outside the typical governance range as narrative exceptions (see text).

Scenarios also differ fundamentally in their assumptions about planetary resource conditions. Rather than define scarcity solely in terms of starting resources, we interpret it as a function of resource pressure (i.e., how quickly a civilization depletes its reserves). That is, systems with a large initial stock may still experience scarcity if it depletes resources at a high rate (e.g., S6). Conversely, a civilization with a modest stock may persist through repeated collapse–recovery cycles if post-collapse resource recovery partially replenishes its base (e.g., S8, S9), or may rebuild following an earlier systemic failure (e.g., S7).

Table 3: Interpreting resource pressure from R_0 and δ

Resource Pressure	R_0	δ	Scenarios
High	≤ 1000	1.0–2.5	S1, S2, S4, S7, S8, S9
Moderate	400–1600	0.5–1.0	S6
Low	≥ 1200	0.0–0.5	S3, S5, S10

We implement Monte Carlo simulations to explore how uncertainty in model parameters shapes the long-term behavior of each scenario. Each scenario’s baseline values (Table 4) are perturbed using probabilistic variation to capture plausible uncertainty (Table 5). Initial resource stock (R_0) and depletion rate (δ) are sampled from normal distributions with modest standard deviations to represent uncertainty in ecological baselines and consumption patterns. Collapse depth (c_f)

and recovery fraction (r_f) are sampled using triangular distributions centered on scenario-specific values, allowing outcomes to concentrate near expected levels while still capturing asymmetries in resilience and recovery capacity. Recovery delay (r_d) is sampled from a normal distribution with a standard deviation of ± 10 years, reflecting variability in rebuilding pace across different societal structures.

The existential hazard rate (h) is held fixed for each scenario because it encodes an assumed background level of external risk (such as AI misalignment, interstellar conflict, or catastrophic environmental feedbacks) rather than internal system dynamics. Unlike other parameters whose variability arises from structural or operational uncertainty, h reflects a narrative-level judgment about a civilization’s exposure to exogenous threats. Scenarios characterized by unstable geopolitical contexts, risky technological dependencies, or unresolved existential challenges (e.g., S1, S6, S7) are assigned higher hazard rates to reflect this vulnerability, while S5 carries a small residual risk due to high systemic complexity. In contrast, scenarios with societal stability, low-impact lifestyles, or resilient infrastructure (e.g., S3, S4, S8, S9, S10) are assigned zero or near-zero hazard rates. Keeping h fixed rather than sampling it probabilistically allows for clearer attribution of collapse outcomes to either endogenous fragility or exogenous shocks, rather than conflating the two.

Table 4: Final parameter values for each scenario

Scenario	r	R_0	δ	c_f	r_d	r_f	h
S1	0.0020	400	1.2	0.6	40	0.15	0.003
S2	0.0026	700	1.5	0.3	70	0.10	0.001
S3	0.0001	2400	0.0	1.0	0	1.0	0.0
S4	0.0060	600	2.5	0.2	100	0.10	0.0
S5	0.0003	2500	0.1	0.9	20	0.80	0.0005
S6	0.0018	1600	1.0	0.4	50	0.25	0.005
S7	0.0060	400	1.5	0.3	60	0.20	0.001
S8	0.0015	800	1.3	0.6	70	0.35	0.0
S9	0.0011	900	1.3	0.5	70	0.35	0.0
S10	0.0002	2700	0.0	1.0	0	1.0	0.0

Table 5: Parameter uncertainty assumptions for Monte Carlo sampling

Parameter	Distribution	Uncertainty
R_0	Normal	$\pm 5\%$ of mean
δ	Normal	$\pm 5\%$ of mean
c_f	Triangular	$\pm 5\%$ of mode
r_d	Normal	± 10 years
r_f	Triangular	$\pm 5\%$ of mode

3.3 Metrics

We define four metrics to reflect different aspects of technospheric persistence and collapse dynamics across Monte Carlo simulations. These metrics are designed to capture the onset, frequency, and continuity of civilization-level activity, with a focus on their implications for detectability.

The first metric, T_c , is the mean time to first collapse across replicates. It serves as a proxy for system fragility: scenarios with low values of T_c tend to break down early, while those with higher values demonstrate greater initial stability. When a simulation run does not experience collapse within the 1000-year window, the full timespan is used. The second metric, f_{nc} , denotes the fraction of runs that avoid collapse entirely. This value reflects the overall robustness of a scenario and helps differentiate consistently stable futures from those vulnerable to systemic failure. We also measure D_c , the average duty cycle, defined as the fraction of time a civilization remains active, that is, maintaining nonzero resource levels and thus potentially emitting technosignatures. This definition assumes that minimal resource availability is a prerequisite for detectable activity. Scenarios with high D_c maintain persistent activity, while those with low D_c may cycle through long dormant periods or experience early termination. Finally, N_c , the mean number of collapse-recovery cycles, captures whether a scenario tends to exhibit singular collapse events or undergo repeated disruptions over time. Together, these metrics provide a compact but informative summary of each scenario’s trajectory, enabling comparisons across sociotechnical regimes and offering insight into the reliability and intermittency of technosignature emission.

4 Results

4.1 Monte Carlo Outcomes and Temporal Profiles

We ran 200 independent simulations for each scenario, introducing probabilistic variation in model parameters as described in Section 3. Figure 1 shows the resulting trajectories of global technology $T(t)$ and available resources $R(t)$, averaged across runs. These profiles reveal the distinct collapse and recovery dynamics that characterize each scenario. As expected, Golden Age (S3) and Out of Eden (S10) show uninterrupted growth and resource stability throughout the 1000-year window, consistent with their post-scarcity narratives and reflected in perfect duty cycles ($\overline{D}_c = 1.0$) and no collapse events ($\overline{N}_c = 0$). In stark contrast, scenarios such as Big Brother (S1) and Sword of Damocles (S6) suffer early collapses ($\overline{T}_c < 210$) and high collapse frequency ($\overline{N}_c \approx 4.5$ for S6 and nearly 10 for S1), leading to sharp contractions in technological capacity and prolonged periods of inactivity. Deus Ex Machina (S9), although collapsing later ($\overline{T}_c \approx 689$), also experiences multiple interruptions and never recovers full continuity, with only modest duty cycle ($\overline{D}_c \approx 0.91$) and about 1.5 collapse events on average.

More complex dynamics emerge in intermediate cases. Transhumanism (S5) avoids collapse in roughly two-thirds of runs and achieves a near-continuous duty cycle ($\overline{D}_c \approx 0.99$), albeit with slower growth and occasional extinction. Living with the Land (S4), in contrast, undergoes frequent and deep collapses ($\overline{N}_c \approx 6.6$), producing the lowest sustained activity among all non-terminal scenarios ($\overline{D}_c \approx 0.38$) despite a steep initial growth phase. Restoration (S7), designed as a post-collapse recovery regime, displays consistent rebuilding behavior: early collapses ($\overline{T}_c \approx 223$), followed by approximately 7.5 recovery cycles and moderate continuity ($\overline{D}_c \approx 0.57$). Ouroboros (S8) exhibits a distinct pattern of low-frequency, self-stabilizing oscillations: it collapses only twice on average, yet maintains robust technological levels and a high duty cycle ($\overline{D}_c \approx 0.87$), reflecting a regime that cycles gracefully without exhausting resources or triggering runaway collapse.

Because the S8 narrative implies repeated collapse–recovery cycles, we ask whether those oscillations are always transient or can be sustained; we probe this with a sweep over r_f and δ (Figure 2). Panel A shows deterministic technology trajectories for varying r_f while holding all other S8 parameters at their baseline values. For low recovery fractions ($r_f \leq 0.15$), the system undergoes




Figure 1: Temporal dynamics of simulated civilizations across ten scenarios. (A) Mean trajectories of global technology $T(t)$ with ± 1 standard deviation shaded. (B) Mean resource levels $R(t)$ with ± 1 standard deviation shaded.

rapid, repeated collapse–recovery cycles (3 cycles within the 1000-year window), as the diminished resource base after each recovery is quickly exhausted. At intermediate values ($0.20 \leq r_f \leq 0.50$), the system sustains two collapse–recovery cycles with longer growth phases between collapses. At higher recovery fractions ($r_f = 0.60$), the first collapse is delayed significantly, and only a single cycle occurs within the simulation window. Panel B maps the number of collapse cycles across the (r_f, δ) parameter space, revealing a clear regime boundary: high depletion rates combined with low recovery fractions produce the most frequent oscillations (up to 8 cycles), while low depletion and high recovery fractions suppress collapse entirely. The baseline S8 parameters (marked with a star) sit near the boundary between 2- and 3-cycle regimes, suggesting that modest shifts in resource recovery capacity could qualitatively alter the civilization’s long-term trajectory. These results confirm that S8’s oscillatory behavior is not universally stable and it depends sensitively on the balance between resource depletion and post-collapse recovery, with some parameter combinations producing sustained cycling and others leading to dampened or terminal collapse.




Figure 2: Parameter sensitivity of S8 (Ouroboros) oscillatory dynamics. (A) Deterministic technology trajectories $T(t)$ for varying recovery fraction r_f , with all other parameters held at baseline values. The number of collapse–recovery cycles within the 1000-year window is indicated in the legend. (B) Number of collapse–recovery cycles as a function of recovery fraction r_f and depletion rate δ . The star marks the baseline S8 parameter combination.

Figure 3 summarizes the outcomes of these simulations using the average of the four metrics described above ($\bar{T}_c$, f_{nc} , $\bar{D}_c$, $\bar{N}_c$). While Golden Age (S3) and Out of Eden (S10) remain fully stable throughout the 1000-year window, with no collapses and continuous activity ($f_{nc} = 1.0$, $\bar{D}_c = 1.0$), other futures display more volatile dynamics. Big Brother (S1) and Sword of Damocles (S6) show early collapses ($\bar{T}_c \leq 220$) and high collapse frequency, while Deus Ex Machina (S9) collapses later ($\bar{T}_c \approx 689$) but still undergoes multiple recovery cycles, reflecting fragile systems with intermittent technosignature potential.

Among the ten scenarios, the mean duty cycle D_c spans a wide range. In the most collapse-prone scenario, S4, D_c was as low as 0.381, indicating these civilizations spend well over half of their time in a collapsed or dormant state. By contrast, scenarios S3 and S10 had $D_c = 1.00$,




Figure 3: Distributions of key collapse and recovery metrics across scenarios (200 runs each). (A) Mean time to first collapse $\bar{T}_c$ with standard deviation bars. (B) Fraction of simulations that never experienced collapse f_{nc} . (C) Mean duty cycle $\bar{D}_c$, defined as the fraction of time the technosphere remains active. (D) Mean number of collapse events $\bar{N}_c$.

never experiencing a collapse (continuous activity). Most scenarios fall somewhere in between. For example, S5 has a high $D_c \approx 0.99$, whereas S2 reaches only $D_c \approx 0.61$, reflecting prolonged instability.

5 Implications for Earth’s Future

A key insight from our modeling is that the parameters governing recovery (the recovery fraction r_f and the recovery delay r_d) are not fixed properties of a civilization but can be deliberately engineered. Investments in civilizational continuity infrastructure directly map onto these parameters: a higher r_f corresponds to greater preservation of knowledge, technology, and resources through collapse events, while a lower r_d corresponds to faster institutional and technological rebuilding. Together, improvements in these parameters increase the duty cycle D_c and thus, increase the fraction of time civilization remains functional and productive.

The concept of “recovery kits” (curated repositories of essential knowledge designed to accelerate post-collapse rebuilding) represents one practical approach to raising r_f . Dartnell [2014] has explored in detail the minimum knowledge base required to reconstruct industrial civilization from scratch, spanning agriculture, materials science, medicine, power generation, and communication. Such recovery kits, whether physical or digital, function as a form of civilizational insurance: they ensure that the knowledge accumulated over centuries of development is not lost when institutional continuity is broken. Ongoing efforts in biological resource preservation serve as real-world examples of resilience infrastructure in action. The Svalbard Global Seed Vault, which now holds over one million seed samples from nearly every country on Earth [Asdal and Guarino, 2020], preserves agricultural biodiversity against civilizational disruption, ensuring that post-collapse societies retain

access to the genetic resources needed to rebuild food systems. More recently, Colossal Biosciences has established a BioVault in Dubai dedicated to preserving broader biodiversity through cryopreservation of cellular material from threatened species [Colossal Biosciences]. In the language of our model, these vaults serve to increase r_f by preserving critical biological capital through collapse events.

Several other model inputs correspond to actionable levers for resilience policy. The depletion rate δ reflects the pace at which a civilization exhausts its resource base. Circular economy principles, renewable energy transitions, and efficiency improvements all serve to reduce δ , extending the time before resource-driven collapse. This connects directly to recent work reframing civilizational growth in terms of thermodynamic efficiency and exploration-exploitation trade-offs [Haqq-Misra et al., 2025b]. A civilization that pursues spatial expansion of its resource domain (exploration) effectively lowers its per-domain δ , while one that intensifies extraction within a fixed domain (exploitation) increases it. The existential hazard rate h can be reduced through investments in planetary defense (e.g., asteroid deflection programs such as NASA’s Double Asteroid Redirection Test, which successfully demonstrated kinetic impact as a viable deflection strategy [Daly et al., 2023]), pandemic preparedness, AI safety research, and nuclear risk reduction, the last of which has been shown capable of triggering global famine affecting over five billion people through climate disruption alone [Xia et al., 2022]. Ord [2020] provides a systematic assessment of existential risks and argues that humanity’s current exposure to catastrophic hazards is historically anomalous and reducible through deliberate policy. In our model, even small reductions in h can significantly decrease the frequency of stochastic collapse events, particularly for scenarios like S1 and S6 where exogenous hazards are a primary driver of instability. The collapse survival fraction c_f can be raised through distributed and redundant infrastructure. Decentralized energy grids, mesh communication networks, and dispersed governance structures all ensure that more technological capacity survives a systemic disruption. Finally, while the initial resource stock R_0 is treated as exogenous in our model, in practice it can be expanded through technologies that unlock previously inaccessible resources (e.g. asteroid mining, deep ocean extraction, nuclear fusion), effectively shifting a civilization into a lower resource-pressure regime.

Viewed through the duty-cycle framework, the question of civilizational resilience becomes quantifiable: what is the marginal improvement in D_c from a given resilience investment? As our parameter sensitivity analysis of S8 demonstrates (Section 4.1), modest shifts in r_f alone can qualitatively alter a civilization’s long-term trajectory. This suggests that targeted resilience investments could yield disproportionate returns in civilizational continuity, and that the duty-cycle framework provides a practical metric for evaluating such investments.

More broadly, our results help reconcile seemingly contradictory perspectives on civilizational longevity. One view posits that technological civilizations are fundamentally fragile and prone to self-destruction soon after emergence, driven by ecological, behavioral, or structural instabilities [Vinn, 2024]. Another suggests that once civilizations surpass key thresholds of resilience and coordination, they may persist indefinitely, steadily accumulating across cosmic time [Grinspoon, 2009]. Both patterns emerge naturally in our simulations. Scarcity-driven, hierarchically governed scenarios frequently enter collapse-recovery cycles, resulting in low duty cycles. In contrast, post-scarcity, distributed scenarios maintain stable resource use and continuous activity. These differences arise not from exogenous shocks, but from internal sociotechnical structure—suggesting that the long-term fate of a civilization is shaped by the interplay of its governance structure, resource management, and resilience infrastructure.

6 Implications for the Search for Technosignatures

6.1 Reframing the Drake Equation’s Longevity Term

The Drake Equation [Drake, 1965] estimates the number of detectable civilizations in our galaxy as

$$N = R_* f_p n_e f_l f_i f_c L, \quad (4)$$

where the final term L , the average duration that civilizations remain detectable, carries outsized importance and remains deeply uncertain. Our simulations suggest that the traditional interpretation of L as a continuous span of emission is unrealistic. To account for the intermittent nature of civilizational activity, we introduce an effective detectability duration,

$$L_{\text{eff}} = D_c T_{\text{span}}, \quad (5)$$

where D_c is the mean duty cycle and T_{span} is the observation window (1000 years in our simulations). If L is better understood as L_{eff} , then models that assume continuous detectability may significantly overestimate N . A galaxy populated by civilizations with low duty cycles would remain largely silent, even if intelligent life is common. We note that a related but distinct concept, a “reappearance factor” for planets where civilizations can arise multiple times in succession [Bhat-tacharya et al., 2011], which counts sequential independent civilizations over geological timescales, whereas our duty cycle refers to the active vs. inactive phases within a single civilization’s lifespan.

Moreover, technology can outlast its creators: the longevity of a technosignature may far exceed the lifespan of the civilization that produced it [Wright et al., 2022, Ćirković et al., 2019]. This further complicates the interpretation of L , which must account not only for active emission but also for the persistence of relic technosignatures, as we examine below.

These considerations help explain the Great Silence. Civilizations may exist in abundance, but if they are detectable only intermittently, the probability that their technosignatures overlap with our observational window is low [Balbi, 2018]. For two intermittent civilizations, the probability of mutual visibility is the product of their duty cycles, which can be very small. The simulation results raise a deeper question: should we expect civilizations to be uniform in character, or drawn from a diverse ensemble of trajectories? If most civilizations follow fragile, collapse-prone paths (e.g., S1, S2, S6), then long-duty-cycle civilizations would be statistical outliers, and the apparent absence of signals may reflect not the rarity of extraterrestrial technology but the dominance of intermittently silent technospheres. Our findings support this view: only S3 and S10 avoided collapse entirely, while most scenarios exhibited high collapse frequency and moderate or low duty cycles. These collapse-prone trajectories suggest that civilizations may not follow monotonic energy-growth trajectories, but instead oscillate or regress, challenging the static assumptions of the Kardashev scale [Kardashev, 1964].

Recent work by Balbi and Grimaldi [2024] provides further theoretical support for this view. Applying Lindy’s law to technosignature longevity, they show that for energy-intensive technoemissions the longevity distribution follows a power law $\rho_L(L) \propto L^{-\alpha}$ with $\alpha > 1$, strongly disfavoring long-lived technosignatures. Under realistic assumptions ($\alpha \geq 2$), the first detected technosignature is predicted to be relatively short-lived ($L < 10^2$ years at 95% confidence). If most detectable technosignatures are intrinsically short-lived, then the combination of Lindy-distributed longevity and low duty cycles (as observed in our collapse-prone scenarios) implies a doubly diminished probability of detection, offering a complementary explanation for the Great Silence.

6.2 Technosignature Persistence and Observation Probabilities

[PLACEHOLDER - Jacob’s per-technosignature detectability framework]

The L_{eff} formulation assumes that detectability ends when activity ceases. As noted above, this need not be the case [Balbi and Ćirković, 2021]. The persistence of different technosignature types varies enormously. Orbital debris and artificial satellites can last from decades to as long as 10,000 to 100,000 years. Satellites in low Earth orbit typically deorbit within decades due to atmospheric drag, while those in geosynchronous orbit, facing minimal resistance, may persist for much longer. Without maintenance, a dense band of satellites in the so-called Clarke belt would slowly degrade through collisions and orbital perturbations over tens of thousands of years, eventually becoming undetectable [Socas-Navarro, 2018]. Megastructures such as Dyson spheres, space mirrors, or stellar engines may persist longer still, though their survival depends on structural design and orbital stability. Rigid constructs without active station-keeping may collapse or drift within a few centuries to several thousand years [Wright, 2020], however, more modular configurations, like swarms of solar collectors, could remain in stable orbits for far longer timescales [Zac and Osmanov, 2022].

Atmospheric technosignatures provide the shortest windows of detectability. On Earth, CFC-11 and CFC-12 have atmospheric lifetimes of about 55 and 140 years, respectively, before being broken down by photolysis or other chemical processes [NOAA Global Monitoring Laboratory, 2025]. Other compounds are more resilient: tetrafluoromethane (CF_4) persists for $\sim 50,000$ years, hexafluoroethane (C_2F_6) for $\sim 10,000$ years, and octafluoropropane (C_3F_8) for $\sim 2,600$ years [Trudinger et al., 2016]. Longer-lived isotopic signatures are conceivable, but their remote detection remains technically challenging. Electromagnetic technosignatures such as radio or laser transmissions propagate indefinitely but are only practically detectable for finite periods. Focused, high-power signals might remain observable for a few hundred light-years over spans of centuries to millennia, depending on the strength of the signal and the sensitivity of receivers [Sheikh et al., 2025].

These persistence times imply that technosignature lifetimes can range from negligible (radio bursts) to dominant (orbital infrastructure). For short-lived civilizations with brief ON phases, persistent technosignatures could substantially extend the effective detectability window beyond what L_{eff} alone would suggest. In contrast, for continuously active civilizations, persistence contributes relatively little. If most detectable technosignatures are short-lived (e.g., radio leakage), then the duty cycle remains the key parameter, and our results suggest low duty cycles make detection unlikely. But if many technosignatures persist for millennia beyond collapse, then the silence we observe becomes harder to attribute solely to civilizational fragility; it would require that either technosignatures are intrinsically ephemeral, or civilizations rarely produce persistent artifacts.

In summary, our results suggest that if detectability is often sporadic, then long periods without contact may be the expected baseline. Intermittency may be typical, and recognizing that could refine our expectations and expand the scope of SETI. We also note that some persistent technologies, particularly autonomous systems such as Bracewell probes [Bracewell, 1960] or orbital maintenance infrastructure, may not decay passively but instead cycle through their own active and dormant states, creating layered duty-cycle structures that merit further investigation.

7 Limitations

Several limitations warrant discussion. Our modeling framework is grounded in Earth-originating scenario typologies, drawing from narrative futures methodologies informed by contemporary geopo-

litical, ecological, and technological trends. These projections embed assumptions about governance structures, resource constraints, and recovery dynamics that may not extend to non-terrestrial civilizations or to those that do not follow Earth-like developmental pathways. While this grounding enables empirical plausibility, it limits extrapolation beyond our biosphere’s trajectory [Rubin, 2001, Denning, 2011]. Our simulations assume bounded planetary systems and do not account for galactic colonization, interstellar migration, or modes of expansion that could alter the spatial and temporal visibility of civilizations [Walters et al., 1980, Papagiannis, 1983]. We also exclude the effects of cultural convergence, information saturation, or technological transcendence (e.g., the emergence of post-biological intelligence), all of which could suppress or qualitatively change technosignature output. Our operational definition of detectability (based on internal sociotechnical continuity) does not differentiate among specific technosignature classes (e.g., radio emissions, waste heat, megastructures), nor does it address detection thresholds or instrumental sensitivity.

Additionally, our assumption that technological capacity increases linearly each year according to a fixed growth rate r is a significant simplification. The many discontinuities implied by the scenarios (not only collapse points but also radical transformation points such as the emergence of artificial general intelligence or breakthroughs in energy harvesting) suggest that a linear growth model may not adequately capture the dynamics of technological change across all regimes. Moreover, the available resource stock R_0 is treated as an exogenous parameter, but in practice it is contingent on the technological capacity to identify and utilize resources. For example, a civilization that develops asteroid mining or nuclear fusion effectively increases its accessible resource base, fundamentally altering the growth-depletion dynamics. Recent work by Haqq-Misra et al. [2025b], which shares the same ten-scenario framework employed here, reframes the Kardashev scale as a luminosity limit constrained by thermodynamic efficiency (exergy) and explores exploration-exploitation trade-offs for long-lived technospheres. Their framework shows that growth trajectories depend on whether a civilization pursues spatial expansion (exploration) or intensive resource use within a fixed domain (exploitation), a distinction that our linear growth model does not capture. Future work could incorporate exergy constraints and non-linear growth functions informed by their luminosity-limit model, as well as feedback between technological capacity and accessible resource stocks, to produce more physically grounded projections of civilizational trajectories.

8 Conclusions

We have developed a hybrid deterministic-stochastic model to simulate collapse-recovery dynamics across ten plausible futures for Earth-originating civilization, and used these simulations to quantify the duty cycles of technological civilizations. Our principal findings are as follows.

First, collapse-recovery dynamics produce a wide range of duty cycles across plausible civilizational futures, with D_c spanning from approximately 0.38 to 1.00 across the ten scenarios. Our simulations naturally produce both fragile, collapse-prone trajectories [Vinn, 2024] and indefinitely persistent ones [Grinspoon, 2009], with outcomes shaped by the interplay of governance structure, resource pressure, and hazard exposure.

Second, the duty-cycle framework reveals that several model parameters correspond to actionable levers for civilizational resilience. The recovery fraction r_f and recovery delay r_d can be deliberately raised through knowledge preservation initiatives, genetic resource vaults, and distributed infrastructure. The depletion rate δ can be reduced through circular economy principles and efficiency improvements. The existential hazard rate h can be lowered through investments in

planetary defense, pandemic preparedness, and nuclear risk reduction. Our parameter sensitivity analysis demonstrates that modest shifts in these parameters (particularly r_f) can qualitatively alter a civilization’s long-term trajectory, suggesting that targeted resilience investments could yield disproportionate returns in civilizational continuity.

Third, the effective detectability duration $L_{\text{eff}} = D_c T_{\text{span}}$ provides a more realistic replacement for the Drake Equation’s static longevity term L . Combined with power-law longevity distributions that disfavor long-lived technosignatures [Balbi and Grimaldi, 2024], low duty cycles imply a doubly diminished probability of detection, helping explain the observed absence of extraterrestrial signals.

Several avenues for future work emerge from this study. Incorporating exergy constraints and non-linear growth functions informed by thermodynamic limits to growth [Haqq-Misra et al., 2025b] could yield more physically grounded simulations. Introducing feedback between technological capacity and accessible resource stocks would address a key simplification of the current model. Finally, extending the framework to account for per-technosignature observation probabilities and for the possibility that autonomous remnant technologies may cycle through their own active and dormant states would add further realism to the detectability analysis.

References

- Åsild Asdal and Luigi Guarino. The Svalbard Global Seed Vault: 10 years—1 million samples. *Biopreserv. Biobank.*, 18(5):391–392, 2020. doi:[10.1089/bio.2018.0025](https://doi.org/10.1089/bio.2018.0025).
- Amedeo Balbi. The impact of the temporal distribution of communicating civilizations on their detectability. *Astrobiology*, 18(1):54–58, January 2018. doi:[10.1089/ast.2017.1652](https://doi.org/10.1089/ast.2017.1652).
- Amedeo Balbi and Milan M Ćirković. Longevity is the key factor in the search for technosignatures. *Astron. J.*, 161(5):222, May 2021. doi:[10.3847/1538-3881/abec48](https://doi.org/10.3847/1538-3881/abec48).
- Amedeo Balbi and Claudio Grimaldi. Technosignature longevity and lindy’s law. *Astron. J.*, 167(3):119, March 2024. doi:[10.3847/1538-3881/ad217d](https://doi.org/10.3847/1538-3881/ad217d).
- A B Bhattacharya, A Sarkar, and J Pandit. Some computations on the drake equation to encapsulate the probable number of broadcasting civilizations. *International Journal of Applied Engineering Research*, 2(2):590–594, 2011.
- Ronald N Bracewell. Communications from superior galactic communities. *Nature*, 186(4726):670–671, May 1960. doi:[10.1038/186670a0](https://doi.org/10.1038/186670a0).
- Milan M Ćirković, Branislav Vukotić, and Milan Stojanović. Persistence of technosignatures: A comment on lingam and loeb. May 2019. doi:[10.48550/arXiv.1905.03146](https://doi.org/10.48550/arXiv.1905.03146).
- Colossal Biosciences. Colossal biovault. <https://colossalfoundation.org/project/species-biobank/>. Accessed: 2026-4-7.
- R Terik Daly, Carolyn M Ernst, Olivier S Barnouin, Nancy L Chabot, et al. Successful kinetic impact into an asteroid for planetary defence. *Nature*, 616(7957):443–447, April 2023. doi:[10.1038/s41586-023-05810-5](https://doi.org/10.1038/s41586-023-05810-5).
- Lewis Dartnell. *The Knowledge: How to Rebuild Our World from Scratch*. Penguin Press, 2014.

- Kathryn Denning. Being technological. *Acta Astronaut.*, 68(3-4):372–380, February 2011. doi:[10.1016/j.actaastro.2010.02.027](https://doi.org/10.1016/j.actaastro.2010.02.027).
- F D Drake. The radio search for intelligent extraterrestrial life. In *Current Aspects of Exobiology*, pages 323–345. Elsevier, 1965. doi:[10.1016/b978-1-4832-0047-7.50015-0](https://doi.org/10.1016/b978-1-4832-0047-7.50015-0).
- Robert H Gray. Intermittent signals and planetary days in SETI. *Int. J. Astrobiology*, 19(4): 299–307, August 2020. doi:[10.1017/s1473550420000038](https://doi.org/10.1017/s1473550420000038).
- Robert H Gray. Intermittent signals and planetary days in SETI. September 2021. doi:[10.48550/arXiv.2109.06175](https://doi.org/10.48550/arXiv.2109.06175).
- David Grinspoon. *Lonely planets*. HarperCollins eBooks, March 2009.
- Jacob Haqq-Misra, George Profitiliotis, and Ravi Kopparapu. Projections of earth’s technosphere: Scenario modeling, worldbuilding, and overview of remotely detectable technosignatures. *Technol. Forecast. Soc. Change*, 218(124194):124194, September 2025a. doi:[10.1016/j.techfore.2025.124194](https://doi.org/10.1016/j.techfore.2025.124194).
- Jacob Haqq-Misra, Clément Vidal, and George Profitiliotis. Projections of earth’s technosphere: Luminosity and mass as limits to growth. *Acta Astronaut.*, 229:831–838, 2025b. doi:[10.1016/j.actaastro.2025.01.048](https://doi.org/10.1016/j.actaastro.2025.01.048).
- Jacob D Haqq-Misra and Seth D Baum. The sustainability solution to the fermi paradox. *J. Br. Interplanet. Soc.*, 62(2):47–51, 2009.
- N Kardashev. Transmission of information by extraterrestrial civilizations. *SvA*, 8:217, October 1964.
- NOAA Global Monitoring Laboratory. Chlorofluorocarbons (CFCs). <https://gml.noaa.gov/hats/publicctn/elkins/cfcs.html>, 2025. Accessed: 2025-6-16.
- Toby Ord. *The Precipice: Existential Risk and the Future of Humanity*. Hachette Books, 2020.
- Elinor Ostrom. Polycentric systems for coping with collective action and global environmental change. *Glob. Environ. Change*, 20(4):550–557, October 2010. doi:[10.1016/j.gloenvcha.2010.07.004](https://doi.org/10.1016/j.gloenvcha.2010.07.004).
- M D Papagiannis. The importance of exploring the asteroid belt. *Acta Astronaut.*, 10(10):709–712, October 1983. doi:[10.1016/0094-5765\(83\)90071-1](https://doi.org/10.1016/0094-5765(83)90071-1).
- Charles T Rubin. L factor: hope and fear in the search for extraterrestrial intelligence. In Stuart A Kingsley and Ragbir Bhathal, editors, *The Search for Extraterrestrial Intelligence (SETI) in the Optical Spectrum III*, volume 4273, pages 230–239. SPIE, August 2001. doi:[10.1117/12.435379](https://doi.org/10.1117/12.435379).
- Sofia Z Sheikh, Macy J Huston, Pinchen Fan, Jason T Wright, Thomas Beatty, Connor Martini, Ravi Kopparapu, and Adam Frank. Earth detecting earth: At what distance could earth’s constellation of technosignatures be detected with present-day technology? *Astron. J.*, 169(2): 118, February 2025. doi:[10.3847/1538-3881/ada3c7](https://doi.org/10.3847/1538-3881/ada3c7).
- Michael Shermer. Why ET hasn’t called. *Sci. Am.*, 287(2):33–33, August 2002. doi:[10.1038/scientificamerican0802-33](https://doi.org/10.1038/scientificamerican0802-33).

- Hector Socas-Navarro. Possible photometric signatures of moderately advanced civilizations: The clarke exobelt. *Astrophys. J.*, 855(2):110, March 2018. doi:[10.3847/1538-4357/aaae66](https://doi.org/10.3847/1538-4357/aaae66).
- Joseph A Tainter. *The Collapse of Complex Societies*. Cambridge University Press, 1988. doi:[10.1017/CBO9780511564505](https://doi.org/10.1017/CBO9780511564505).
- Cathy M Trudinger, Paul J Fraser, David M Etheridge, William T Sturges, Martin K Vollmer, Matt Rigby, Patricia Martinerie, Jens Mühle, David R Worton, Paul B Krummel, L Paul Steele, Benjamin R Miller, Johannes Laube, Francis S Mani, Peter J Rayner, Christina M Harth, Emmanuel Witrant, Thomas Blunier, Jakob Schwander, Simon O’Doherty, and Mark Battle. Atmospheric abundance and global emissions of perfluorocarbons CF_4 , C_2F_6 and C_3F_8 since 1800 inferred from ice core, firn, air archive and in situ measurements. *Atmos. Chem. Phys.*, 16(18):11733–11754, September 2016. doi:[10.5194/acp-16-11733-2016](https://doi.org/10.5194/acp-16-11733-2016).
- Olev Vinn. Potential incompatibility of inherited behavior patterns with civilization: Implications for fermi paradox. *Sci. Prog.*, 107(3):368504241272491, July 2024. doi:[10.1177/00368504241272491](https://doi.org/10.1177/00368504241272491).
- Clifford Walters, Raymond A Hoover, and R K Kotra. Interstellar colonization: A new parameter for the drake equation? *Icarus*, 41(2):193–197, February 1980. doi:[10.1016/0019-1035\(80\)90003-2](https://doi.org/10.1016/0019-1035(80)90003-2).
- Stephen Webb. *If the universe is teeming with aliens ... WHERE IS EVERYBODY?* Springer, New York, NY, December 2010. doi:[10.1007/b97464](https://doi.org/10.1007/b97464).
- P Wesson. Cosmology, extraterrestrial intelligence, and a resolution of the fermi-hart paradox. *Quarterly journal of the royal astronomical society*, 31:161–170, June 1990.
- Michael L Wong and Stuart Bartlett. Asymptotic burnout and homeostatic awakening: a possible solution to the fermi paradox? *J. R. Soc. Interface*, 19(190):20220029, May 2022. doi:[10.1098/rsif.2022.0029](https://doi.org/10.1098/rsif.2022.0029).
- Jason Wright, T. Dyson spheres. *Serbian Astron. J.*, (200):1–18, 2020. doi:[10.2298/saj2000001w](https://doi.org/10.2298/saj2000001w).
- Jason T Wright, Jacob Haqq-Misra, Adam Frank, Ravi Kopparapu, Manasvi Lingam, and Sofia Z Sheikh. The case for technosignatures: Why they may be abundant, long-lived, highly detectable, and unambiguous. *Astrophys. J. Lett.*, 927(2):L30, March 2022. doi:[10.3847/2041-8213/ac5824](https://doi.org/10.3847/2041-8213/ac5824).
- Lili Xia, Alan Robock, Kim Scherrer, Cheryl S Harrison, Benjamin Leon Bodirsky, Isabelle Weindl, Jonas Jägermeyr, Charles G Bardeen, Owen B Toon, and Ryan Heneghan. Global food insecurity and famine from reduced crop, marine fishery and livestock production due to climate disruption from nuclear war soot injection. *Nat. Food*, 3(8):586–596, August 2022. doi:[10.1038/s43016-022-00573-0](https://doi.org/10.1038/s43016-022-00573-0).
- Seth Zac and Zaza Osmanov. Review and viability of a Dyson swarm as a form of Dyson sphere. *Phys. Scr.*, 97(12):122001, 2022. doi:[10.1088/1402-4896/ac9e78](https://doi.org/10.1088/1402-4896/ac9e78).