



Your new experience awaits. Try the new design now and help us make it even better

Switch to the ...

5,8K

Total views

1,4K

Downloads

24

Citations



Download article ▾

[View article impact >](#)

Supplemental data



BRIEF RESEARCH REPORT article

Front. Vet. Sci., 10 July 2020

Sec. Veterinary Infectious Diseases

Volume 7 - 2020 |

<https://doi.org/10.3389/fvets.2020.00346>

This article is part of the Research Topic

Antibiotic Resistance and its Continuity in the Environmental Niche

[View all 19 articles >](#)

Widespread Environmental Presence of Multidrug-Resistant *Salmonella* in an Equine Veterinary Hospital That Received Local and International Horses



Paula Soza-Ossandón¹



Dácil Rivera^{1,2}



Rodolfo Tardone¹

We use cookies

Our website uses cookies that are essential for its operation and additional cookies to track performance, or to improve and personalize our services. To manage your cookie preferences, please click [Cookie Settings](#). For more information on how we use cookies, please see our [Cookie Policy](#)

[Cookies Settings](#)

Reject non-essential cookies

Accept cookies

morbidity and mortality rates, nosocomial transmission to other patients, zoonotic transmission to hospital personnel, and even closure of facilities. In this study, 545 samples (environmental and hospitalized patients) were collected monthly during a 1-year period from human and animal contact surfaces in an equine hospital that received local and international horses. A total of 22 *Salmonella* isolates were obtained from human contact surfaces (e.g., offices and pharmacy) and animal contact surfaces (e.g., stalls, surgery room, and waterers), and one isolate from a horse. Molecular serotyping revealed 18 isolates as *Salmonella* Typhimurium and three as *Salmonella* Infantis. Nineteen isolates were resistant to at least one antimicrobial class, and only two isolates were susceptible to all antimicrobials tested. In addition, we identified nine multidrug-resistant (MDR) isolates in *S. Typhimurium*, which displayed resistance to up to eight antimicrobials (i.e., amoxicillin/clavulanate, ampicillin, ciprofloxacin, chloramphenicol, streptomycin, gentamicin, trimethoprim/sulfamethoxazole, and tetracycline). Pulsed-field gel electrophoresis (PFGE) revealed the presence of three PFGE patterns permanently present in the environment of the hospital during our study. The persistent environmental presence of MDR *Salmonella* isolates, along with the fact that local and international horses are attended in this hospital, highlights the importance of improving biosecurity programs to prevent disease in horses and the hospital personnel and also for the global dissemination and acquisition of MDR *Salmonella*.

Introduction

Salmonella enterica, a Gram-negative bacteria of the family Enterobacteriaceae, is an important zoonotic pathogen that causes an estimated of 93.8 human cases and 150,000 deaths every year worldwide (1). *Salmonella* is usually transmitted to humans as foodborne and through contact with infected animals (2). This pathogen is a microorganism responsible for gastrointestinal disease affecting equines (among other animals) of all ages (3). Clinical symptoms include diarrhea, fever, and dehydration, with severity ranging from a subclinical colonization to a severe systemic illness (4). As a highly contagious disease, it can be reported as

We use cookies

Our website uses cookies that are essential for its operation and additional cookies to track performance, or to improve and personalize our services. To manage your cookie preferences, please click [Cookie Settings](#). For more information on how we use cookies, please see our [Cookie Policy](#)

Salmonellosis outbreaks in animal health facilities are full of challenges beside the sole medical treatment and control the outbreak *per se*; they also involve communication with owners and referring veterinarians of infected horses (10). On the other hand, the consequences are serious including hospital-acquired infections of patients and hospital personnel, the establishment of expensive infection control programs, and decrease in clients' trust and hospitals' revenues and may even lead to litigation procedures (11, 12). Infection control programs should be an integral part of every animal health facility (16, 17). Several studies have reported outbreak control measurements (7, 12, 18) and assessment of protocols of contamination, which have been adopted by many facilities (16, 19). To date, there are no reports of salmonellosis in veterinary hospitals in Chile, and therefore, scarce biosecurity protocols have been established. Hence, this study was performed to determine the presence, antimicrobial resistance, and subtypes of *Salmonella* in the environment and patients from an EVH without reported history of outbreaks or hospital-acquired infections.

Materials and Methods

Description of the Setting and Location

The EVH is located at a thoroughbred horse racetrack at the center of the city of Santiago (Chile). It has an average flow of 100 incoming patients daily, providing equine health services to Thoroughbred, Arabian, Chilean rodeo, and Warmblood horses. This veterinary hospital has no records of outbreak or hospital-acquired infections due to *Salmonella* spp., and this information is remarkable in view of the lack of biosecurity measures or infection control programs (e.g., isolation of infected patients and protocols for cleaning and sanitation).

Sampling Procedure

A total of 545 samples were obtained in a longitudinal study conducted from July 2015 to June 2016. With the corresponding consent from the Chief Director, we collected both environmental ($n = 61$, for details see Table 1) samples and patient fecal samples, from one to nine, depending on hospitalized horses at a given time

We use cookies

Our website uses cookies that are essential for its operation and additional cookies to track performance, or to improve and personalize our services. To manage your cookie preferences, please click [Cookie Settings](#). For more information on how we use cookies, please see our [Cookie Policy](#)

Table 1

Sample origin	No. of samples	No. positive samples	% positive samples
Animal feces	53	1	1.88
Environmental/surgery (SA)^a			
Stalls (1–4)	48	1	2.08
Surgery room floor	12	2	16.67
Bed	12	0	0
Pharmacy	12	1	8.33
Washing room	12	1	8.33
Dressing room	12	0	0
Personal entrance	12	0	0
Office	12	0	0
Induction/recovery room	12	0	0
Area Floor	12	0	0
Environmental/hospitalization (HA)^a			
Stalls (5–10)	72	3	4.17
Floor	12	1	8.33
Environmental/proceeding (PA)^a			
Pharmacy	12	1	8.33
Floor	12	1	8.33
Main office	12	2 ^b	16.67
Environmental/equipment (EQ)^a			
Twitches (3×)	36	1	2.78
Endoscope	12	1	8.33
Gastroscope	12	1	8.33
Pitchforks (2×)	24	2	8.33
Waterers (1×)	120	1	0.83
Environmental/exterior (EA)^a			
Manure collection site	12	1	8.33
Total	545	21	3.85

^aEnvironmental samples were classified according to how the hospital was divided into four main areas, plus equipment (see **Figure 1** and Materials and Methods).

^bTwo different isolates were obtained from one sample taken on September 2015.

Table 1. Results of *Salmonella* spp. on samples collected in the equine veterinary hospital during the study.

Figure 1

A EA

Figure 1. Schematic diagram of the

We use cookies

Our website uses cookies that are essential for its operation and additional cookies to track performance, or to improve and personalize our services. To manage your cookie preferences, please click [Cookie Settings](#). For more information on how we use cookies, please see our [Cookie Policy](#).

entrance; Bth, bathroom; Dr, dressing room; MOF, main office; SYr, surgery yard; HYr, hospitalization yard; mcs, manure collection site.

Figure 2

Figure 2. Dendrogram representation of *Salmonella enterica* isolates clustered using unweighted pair group method with arithmetic mean (UPGMA) method. Five pulse types of *Salmonella* were identified at the right; colors have been assigned for each pulse type (A–D), matching **Figure 1B**.

Bacterial Culture and Molecular Identification

Salmonella isolation was conducted as previously described (22). In brief, all samples were cultured in peptone water at 37°C overnight, and 100 µl and 1 ml were transferred into Rappaport–Vassiliadis (RV) (BD, Franklin Lakes, NJ) supplemented with novobiocin (20 mg/ml) and 100 µl of Tetrathionate (TT) (BD,

We use cookies

Our website uses cookies that are essential for its operation and additional cookies to track performance, or to improve and personalize our services. To manage your cookie preferences, please click [Cookie Settings](#). For more information on how we use cookies, please see our [Cookie Policy](#)

disk dispenser was used, along with the antimicrobial disks, detailed as follows: amikacin (AMK; 30 µg), amoxicillin/clavulanate (AMC; 30 µg), ampicillin (AMP; 10 µg), ceftiofur (FOX; 30 µg), ceftriaxone (CTR; 30 µg), ciprofloxacin (CIP; 5 µg), chloramphenicol (CHL; 30 µg), streptomycin (STR; 300 µg), gentamicin (GEN; 10 µg), kanamycin (KAN; 30 µg), trimethoprim/sulfamethoxazole (SXT; 23.75 µg), and tetracycline (TET; 30 µg). The agar plates were incubated at 37°C overnight.

Escherichia coli American Type Culture Collection (ATCC) 25922 was used as control. Interpretations were made based on the guidelines of Clinical Laboratory Standard Institute (25). The samples were classified according to Magiorakos's criteria as MDR when resistant to at least one agent in three or more antimicrobial classes (26).

Molecular Characterization of *Salmonella* Serotype

A previously described molecular method for serotype prediction was used (27, 28). Briefly, DNA extraction of the isolates was conducted using the DNeasy Blood and Tissue kit (QIAGEN, Hilden, Germany). The molecular scheme included an initial multiplex PCR, conducted to identify the serogroup of each isolate, followed by PCR-sequencing approaches to determine H1 and H2 antigens (27, 28). PCR products were sent to MACROGEN™ (Korea) for Sanger sequencing. Consensus sequences were obtained using CAP3 Sequence Assembly Program (<http://douna.prabi.fr/software/cap3>); the complementary reverse was obtained by using Bioinformatics.org. The results were analyzed using basic local alignment tool (BLAST) on the National Center for Biotechnology Information (NCBI).

Molecular Typing

Molecular typing of the isolates was conducted by pulsed-field gel electrophoresis (PFGE), using the CDC PulseNet standard protocol (29). For this, overnight cultures in brain heart infusion broth (BHI, BD, Germany) were embedded in 1% of SeaKem® Gold Agarose (Lonza, Rockland, ME, USA). Upon lysis and washing, the plugs were digested with *Xba*I (Thermo Fisher Scientific Inc., Waltham, MA). The CHEF-DR® III System (Bio-Rad Laboratories, Hercules, CA) was used for the electrophoresis for 20 h. A standard, *Salmonella* Braenderup

We use cookies

Our website uses cookies that are essential for its operation and additional cookies to track performance, or to improve and personalize our services. To manage your cookie preferences, please click [Cookie Settings](#). For more information on how we use cookies, please see our [Cookie Policy](#)

10/21 (47.6%) samples, *Salmonella* isolates were obtained from both enrichments conducted. On positive samples, one isolate was selected, except for one sample, in which two different colonies were obtained; therefore, a total of 22 *Salmonella* colonies were further characterized. From the 22 isolates, 1/22 (4.5%) was obtained from a sick Chilean rodeo patient, which died of peritonitis after colic surgery (no positive foreign patients were found), and the other 21 (21/23; 95.4%) were obtained from 20 environmental samples (i.e., stalls, surgery room floor, surgical pharmacy, washing room, hospitalization area floor, main office, pitchforks, endoscope, gastroscope, twitches, waterers, manure collection site, proceeding area floor, and pharmacy) ([Table 1](#)). Regarding the type of contact surface, 13/396 (3.28%) isolates were obtained from animal contact surfaces and 8/96 (9.38%) from human contact surfaces ([Table 2](#)). About the dates of isolation, two peaks were seen during the months of September 2015 and May 2016, where 9/22 and 8/22 isolates of *Salmonella* spp. were obtained, respectively ([Supplementary Figure 1](#)). A few isolates were also obtained during October 2015 ($n = 1$), December 2015 ($n = 1$), April 2016 ($n = 1$), and May 2016 ($n = 2$) ([Table 2](#), [Supplementary Figure 1](#)).

Table 2

Table 2. Characteristics, serotypes, PFGE patterns, and antimicrobial resistance of *Salmonella* isolates.

Presence of Multidrug-Resistant *Salmonella* Isolates

We use cookies

Our website uses cookies that are essential for its operation and additional cookies to track performance, or to improve and personalize our services. To manage your cookie preferences, please click [Cookie Settings](#). For more information on how we use cookies, please see our [Cookie Policy](#)

isolates belonging to O:7 (C1) serogroup (2/22) were predicted as *Infantis* serotype (Table 2).

Five Different Pulsed-Field Gel Electrophoresis Types of *Salmonella* Were Identified

According to the PFGE, four PFGE patterns were identified in 19 *Salmonella typhimurium* isolates, and one PFGE type was found in three *S. Infantis* isolates. Among *S. Typhimurium*, seven isolates (1, 2, 3, 4, 7, 10, and 13) were indistinguishable from each other and classified as PFGE pattern A. In three isolates (5, 6, and 8), PFGE patterns were also indistinguishable from each other and related to PFGE pattern A, which was therefore classified as A1. All PFGE patterns A and A1 were detected only in the sampling of September 2015. Eight isolates (9, 12, 14, 15, 17, 18, 19, and 20) were indistinguishable from each other and different from all others, classified as PFGE pattern B; these isolates were obtained in samplings of September 2015 and in April and May 2016. One additional PFGE pattern D of isolate 16 was found in *S. Typhimurium*. Isolates 11, 21, and 22 were indistinguishable from each other and different from all others, classified as PFGE pattern C. Importantly, these isolates were classified as *S. Infantis* (Table 2).

Discussion

This study examined the environmental presence of *Salmonella* in an equine hospital with no history of outbreak or hospital-acquired infections. Here, we identified two serotypes that were widely distributed. The major findings of this study are the following: (i) wide spatial distribution of *Salmonella* in the hospital, mainly in spring and autumn; (ii) MDR *Salmonella* Typhimurium accounted for most of the isolates; and (iii) multiple *Salmonella* PFGE patterns present in human contact surfaces highlight the need of developing biosecurity standard protocols.

Wide Spatial Distribution of *Salmonella* in the Hospital, Mainly in Spring and Autumn

In this study, we found a considerable presence of *Salmonella* in the EVH environment, compared with the equine's samples. The prevalence of *Salmonella* in equine subclinical shedders (1–2%) tends to increase under stress conditions

We use cookies

Our website uses cookies that are essential for its operation and additional cookies to track performance, or to improve and personalize our services. To manage your cookie preferences, please click [Cookie Settings](#). For more information on how we use cookies, please see our [Cookie Policy](#)

but further investigation is needed, which is beyond the scope of this study. Importantly, these isolates represented a closely related PFGE pattern. Nevertheless, neither official information nor patient history could be collected to explain the second peak, in June 2016. Although it is uncertain about the origin of these isolates, shedding patients present during non-sampling periods could be a common source of dissemination (5). Other possible sources of contaminations, such as other animals (rodents), feed, or even environmental persistent strains (34), are also plausible and have to be considered.

Multidrug-Resistant *Salmonella* Typhimurium Accounted for Most Isolates

Reported outbreaks of *Salmonella* in EVHs have involved serotypes such as Typhimurium, Newport, Agona, Anatum (12, 35), Infantis (36), Heidelberg (37), and Oranienburg (38). Here, we found that 87% of the isolates were represented by *S. Typhimurium*. This serotype has been commonly isolated from horses, causing severe clinical signs, along with high morbidity and mortality rates (7, 8, 33). In Chile, only one outbreak of *S. Typhimurium* has been reported, which affected weanling foals with a morbidity rate of 87% and mortality rate of 13% (39). Regarding *Salmonella* Infantis, which is less commonly reported compared with *S. Typhimurium*, only three isolates were found. Nonetheless, there is a report of a serious outbreak in 1996, which caused important economic losses and even the closure of the facilities (36).

Antimicrobial resistance profiles, which include resistance to AMP (10 isolates), as the most common profile, followed by the profile AMC–AMP–CRO–CHL–STR–TE (six isolates), include antimicrobials in which resistance has already been described in other salmonellosis outbreaks (38), not only in equine hospitals but also in small animal shelters (13). Notably, we found that almost half of the isolates ($n = 10$) displayed an MDR phenotype, showing resistance to three or more antimicrobial classes (26), which is a major concern for the public health, the personnel at the hospital, and the treatment of hospitalized horses.

Multiple *Salmonella* Pulsed-Field Gel Electrophoresis Patterns Present in Human Contact Surfaces Highlight the Need of Developing Biosecurity Standards

We use cookies

Our website uses cookies that are essential for its operation and additional cookies to track performance, or to improve and personalize our services. To manage your cookie preferences, please click [Cookie Settings](#). For more information on how we use cookies, please see our [Cookie Policy](#)

implementation of biosecurity protocols is quite expensive, it is much less than controlling an outbreak itself, especially considering the fact that the EVH located at a thoroughbred racetrack, harbors nearly 1,500 horses together with hospital personnel (17).

Conclusions

This study has revealed the importance of implementing mitigation strategies and biosecurity protocols to control MDR *Salmonella* to ensure the safety of patients and hospital personnel. Also, this could set an example for other veterinary facilities to establish or recheck their functioning biosecurity protocols, especially in developing countries.

Data Availability Statement

The datasets generated for this study are available on request to the corresponding author.

Ethics Statement

The animal study was reviewed and approved by The University Andres Bello Bioethics Committee, Santiago, Chile. Written informed consent for participation was not obtained from the owners because fresh fecal samples were obtained from the floor.

Author Contributions

PS-O designed the study, conducted the experiments, and wrote the manuscript. AM-S wrote the manuscript, analyzed data, and designed the study. DR and RT conducted the experiments. RR-N critically reviewed the manuscript. AA, GG-R, and CH-W analyzed the data. PG conducted the experiments. All authors contributed to the article and approved the submitted version.

Funding

We thank the following funding sources: ANID Millennium Science

We use cookies

Our website uses cookies that are essential for its operation and additional cookies to track performance, or to improve and personalize our services. To manage your cookie preferences, please click [Cookie Settings](#). For more information on how we use cookies, please see our [Cookie Policy](#)

References

1. Majowicz SE, Musto J, Scallan E, Angulo FJ, Kirk M, O'Brien SJ, et al. The global burden of nontyphoidal *salmonella* gastroenteritis. *Clin Infect Dis*. (2010) 50:882–9. doi: 10.1086/650733

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

2. Hoelzer K, Switt AIM, Wiedmann M. Animal contact as a source of human non-typhoidal salmonellosis. *Vet Res*. (2011) 42:1–27. doi: 10.1186/1297-9716-42-34

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

3. FDA - USA Food and Drug Administration. *Get the Facts About Salmonella! Anim Heal Lit*. (2017). Available online at: <https://www.fda.gov/animal-veterinary/animal-health-literacy/get-facts-about-salmonella> (accessed February 15, 2020).

[Google Scholar](#)

4. Timoney J. *Salmonella* in domestic animals. 2nd edition. In: Paul A, Barrow U, editors. *Methner*. Oxfordshire: CABI. (2013). 305–17 p.

[PubMed Abstract](#) | [Google Scholar](#)

5. Ernst NS, Hernandez JA, MacKay RJ, Brown MP, Gaskin JM, Nguyen AD, et al. Risk factors associated with fecal *salmonella* shedding among hospitalized horses with signs of gastrointestinal tract disease. *J Am Vet Med Assoc*. (2004) 225:275–81. doi: 10.2460/javma.2004.225.275

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

We use cookies

Our website uses cookies that are essential for its operation and additional cookies to track performance, or to improve and personalize our services. To manage your cookie preferences, please click [Cookie Settings](#). For more information on how we use cookies, please see our [Cookie Policy](#)

multidrug-resistant *salmonella* typhimurium in a population of hospitalized horses. *Vet Microbiol.* (2005) 107:233–40. doi: 10.1016/j.vetmic.2005.01.019

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

9. Dallap Schaer BL, Aceto H, Rankin SC. Outbreak of salmonellosis caused by *Salmonella enterica* serovar newport MDR-AmpC in a large animal veterinary teaching hospital. *J Vet Intern Med.* (2010) 24:1138–46. doi: 10.1111/j.1939-1676.2010.0546.x

[CrossRef Full Text](#) | [Google Scholar](#)

10. Dargatz D A, Traub-Dargatz JL. Multidrug-resistant *Salmonella* and nosocomial infections, *Vet Clin North Am Equine Prac.* (2004) 20:587–600. doi: 10.1016/j.cveq.2004.07.008

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

11. Johnson J A. Nosocomial infections. *Vet Clin North Am Small Anim Pract.* (2002) 32:1101–26. doi: 10.1016/S0195-5616(02)00038-4

[CrossRef Full Text](#) | [Google Scholar](#)

12. McMillan NS. *The Management of a Multidrug-Resistant Salmonella Agona Outbreak at Large Animal Teaching Hospital.* (2004) Available online at: <http://scholar.lib.vt.edu/theses/available/etd-05272004-144444/unrestricted/Chapter2NSM.pdf> (accessed May 11, 2015).

[Google Scholar](#)

13. Wright JG, Tengelsen LA, Smith KE, Bender JB, Frank RK, Grendon JH, et al. Multidrug-resistant *salmonella* typhimurium in four animal facilities. *Emerg Infect Dis.* (2005) 11:1235–41. doi: 10.3201/eid1108.050111

We use cookies

Our website uses cookies that are essential for its operation and additional cookies to track performance, or to improve and personalize our services. To manage your cookie preferences, please click [Cookie Settings](#). For more information on how we use cookies, please see our [Cookie Policy](#)

16. Benedict KM, Morley PS, Van Metre DC. Characteristics of biosecurity and infection control programs at veterinary teaching hospitals. *J Am Vet Med Assoc.* (2008) 233:767–73. doi: 10.2460/javma.233.5.767

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

17. Morley PS, Anderson MEC, Burgess BA, Aceto H, Bender JB, Clark C, et al. Report of the third havemeyer workshop on infection control in equine populations. *Equine Vet J.* (2013) 45:131–6. doi: 10.1111/evj.12000

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

18. Steneroden KK, Van Metre DC, Jackson C, Morley PS. Detection and control of a nosocomial outbreak caused by *salmonella* newport at a large animal hospital. *J Vet Intern Med.* (2010) 24:606–16. doi: 10.1111/j.1939-1676.2010.0484.x

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

19. Alinovi CA, Ward MP, Couëtil LL, Wu CC. Detection of *salmonella* organisms and assessment of a protocol for removal of contamination in horse stalls at a veterinary teaching hospital. *J Am Vet Med Assoc.* (2003) 223:1640–4. doi: 10.2460/javma.2003.223.1640

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

20. Hoet AE, Johnson A, Nava-Hoet RC, Bateman S, Hillier A, Dyce J, et al. Environmental methicillin-resistant *staphylococcus aureus* in a veterinary teaching hospital during a nonoutbreak period. *Vector Borne Zoonotic Dis.* (2011) 11:609–615. doi: 10.1089/vbz.2010.0181

[CrossRef Full Text](#) | [Google Scholar](#)

We use cookies

Our website uses cookies that are essential for its operation and additional cookies to track performance, or to improve and personalize our services. To manage your cookie preferences, please click [Cookie Settings](#). For more information on how we use cookies, please see our [Cookie Policy](#)

Escherichia Coli O157:H7, *Salmonella*, *Staphylococcus aureus*, *listeria monocytogenes*, and *vibrio parahaemolyticus*. *J Food Prot.* (2007) 70:1656–62. doi: 10.4315/0362-028X-70.7.1656

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

24. Hudzicki J (University of KMC. *Kirby Bauer Disk Diffusion Susceptibility Test Protocol*. (2016) Available online at: <https://www.asmscience.org/content/education/protocol/protocol.3189> (accessed January 1, 2016).

[PubMed Abstract](#) | [Google Scholar](#)

25. Clinical and Laboratory Standards Institute (CLSI). *Performance Standards for Antimicrobial Susceptibility Testing*. M100, 27th edition. Wayne, PA: Clinical and Laboratory Standards Institute. (2017).

[Google Scholar](#)

26. Magiorakos A, Srinivasan A, Carey RB, Carmeli Y, Falagas ME, Giske CG, et al. Multidrug-resistant, extensively drug-resistant and pandrug-resistant bacteria: an international expert proposal for interim standard definitions for acquired resistance. *Clin Microbiol Infect.* (2011) 18:268–81. doi: 10.1111/j.1469-0691.2011.03570.x

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

27. Herrera-Leon S, Ramiro R, Arroyo M, Diaz R, Usera MA, Echeita MA. Blind comparison of traditional serotyping with three multiplex PCRs for the identification of *salmonella* serotypes. *Res Microbiol.* (2007) 158:122–7. doi: 10.1016/j.resmic.2006.09.009

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

We use cookies

Our website uses cookies that are essential for its operation and additional cookies to track performance, or to improve and personalize our services. To manage your cookie preferences, please click [Cookie Settings](#). For more information on how we use cookies, please see our [Cookie Policy](#)

[Google Scholar](#)

31. Tenover FC, Arbeit RD, Goering R V, Mickelsen PA, Murray BE, Persing DH, et al. Interpreting chromosomal DNA restriction patterns produced by pulsed-field gel electrophoresis: criteria for bacterial strain typing. *J Clin Microbiol.* (1995) 33:2233–9. doi: 10.1128/JCM.33.9.2233-2239.1995

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

32. Albers AL, Mollenkopf DF, Mathys DA, Daniels JB, Wittum TE. *Salmonella Enterica Prevalence in the Ohio State University Veterinary Medical Center Environment.* (2017) Available online at: <https://kb.osu.edu/handle/1811/80678> (accessed February 03, 2018).

[Google Scholar](#)

33. Jones SL. Inflammatory Diseases of the Gastrointestinal Tract Causing Diarrhea, In Reed S, editors. *Equine Internal Medicine*, (St. Louis: Saunders), p. 884–888.

[Google Scholar](#)

34. Wales A, Davies RH. Environmental Aspects of *Salmonella*, In Barrows PA, Methner U, editors. *Salmonella in Domestic Animals*. Oxfordshire: CABI. p. 399–425. doi: 10.1079/9781845939021.0399

[CrossRef Full Text](#) | [Google Scholar](#)

35. Hartmann FA, West SE. Antimicrobial susceptibility profiles of multidrug-resistant *salmonella* anatum isolated from horses. *J Vet Diagn Invest.* (1995) 7:159–61. doi: 10.1177/104063879500700128

We use cookies

Our website uses cookies that are essential for its operation and additional cookies to track performance, or to improve and personalize our services. To manage your cookie preferences, please click [Cookie Settings](#). For more information on how we use cookies, please see our [Cookie Policy](#)

veterinary medical teaching hospital with evidence of nosocomial and on-farm transmission. *Vector Borne Zoonotic Dis.* (2014) 14:496–502. doi: 10.1089/vbz.2013.1467

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

39. Soza-Ossandón P, Toledo V, Moreno-Switt A. Brote de *Salmonella* spp. en potrillos al pie, en un criadero de caballos fina sangre de carrera (fsc) de la región metropolitana en Chile. in *XXIII Congreso Latino Americano de Microbiología* (Rosario, Argentina), 1160. Available online at: <http://www.alam-cam2016.aam.org.ar/descarga/LibroCAMALAM2016.pdf> (accessed February 03, 2018).

[Google Scholar](#)

40. Ekiri AB, Morton AJ, Long MT, MacKay RJ, Hernandez JA. Review of the epidemiology and infection control aspects of nosocomial *Salmonella* infections in hospitalised horses. *Equine Vet Educ.* (2010) 22:631–41. doi: 10.1111/j.2042-3292.2010.00144.x

[CrossRef Full Text](#) | [Google Scholar](#)

41. Duce G, Fabry J, Nicolle L. *Prevention of Hospital Acquired Infection: A Practical Guide*. World Heal Organ. (2002). p. 1–72. Available at: <http://www.who.int/csr/resources/publications/whocdscsreph200212.pdf> (accessed May 11, 2015).

[Google Scholar](#)

Keywords: *Salmonella enterica*, multidrug-resistant, equine hospital, hospital-acquired infections, biosecurity

We use cookies

Our website uses cookies that are essential for its operation and additional cookies to track performance, or to improve and personalize our services. To manage your cookie preferences, please click [Cookie Settings](#). For more information on how we use cookies, please see our [Cookie Policy](#)

article distributed under the terms of the [Creative Commons Attribution License \(CC BY\)](#). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.

***Correspondence:** Andrea I. Moreno-Switt, andrea.moreno@unab.cl; andrea.moreno@uc.cl

Disclaimer: All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article or claim that may be made by its manufacturer is not guaranteed or endorsed by the publisher.

Guidelines



Explore



Outreach



Connect



Follow us

We use cookies

Our website uses cookies that are essential for its operation and additional cookies to track performance, or to improve and personalize our services. To manage your cookie preferences, please click [Cookie Settings](#). For more information on how we use cookies, please see our [Cookie Policy](#)

We use cookies

Our website uses cookies that are essential for its operation and additional cookies to track performance, or to improve and personalize our services. To manage your cookie preferences, please click [Cookie Settings](#). For more information on how we use cookies, please see our [Cookie Policy](#)