



Bioactive immunostimulants as health-promoting feed additives in aquaculture: A review

Seerengaraj Vijayaraj^{a,b}, Yun-Zhang Sun^{a,c,**}, Antonio Zuurro^d, Hamed Ghafarifarsani^e, Hien Van Doan^{f,g,*}, Seyed Hossein Hoseinifar^h

^a Xiamen Key Laboratory for Feed Quality Testing and Safety Evaluation, Fisheries College, Jimei University, Xiamen, China

^b Department of Environmental Studies, School of Energy Environment and Natural Resources, Madurai Kamaraj University, Madurai, India

^c The Key Laboratory of Healthy Mariculture for the East China Sea, Ministry of Agriculture, Fisheries College, Jimei University, Xiamen, China

^d Department of Chemical Engineering, Materials and Environment, Sapienza University, Via Eudossiana 18, 00184, Rome, Italy

^e Department of Fisheries, Faculty of Natural Resources, Urmia University, Urmia, Iran

^f Department of Animal and Aquatic Sciences, Faculty of Agriculture, Chiang Mai University, Chiang Mai, Thailand

^g Science and Technology Research Institute, Chiang Mai University, Suthep, Muang, Chiang Mai, Thailand

^h Department of Fisheries, Faculty of Fisheries and Environmental Sciences, Gorgan University of Agricultural Sciences and Natural Resources, Gorgan, Iran

ARTICLE INFO

Keywords:

Bioactive immunostimulants
Immune system
Disease resistance
Feed additives
Aquaculture

ABSTRACT

Bioactive immunostimulants could be derived from different sources like plants, animals, microbes, algae, yeast, etc. Bioactive immunostimulants are the most significant role to enhance aquatic production, as well as the cost of this method, which is effective, non-toxic, and environment-friendly. These immunostimulants are supportive to increase the immune system, growth, antioxidant, anti-inflammatory, and disease resistance of aquatic animals' health and also improve aquatic animal feed. Diseases are mainly targeted to the immune system of aquatic organisms in such a way that different processes of bioactive immunostimulants progress are considered imperative techniques for the development of aquaculture production. Communicable infections are the main problem for aquaculture, while the mortality and morbidity connected with some outbreaks significantly limit the productivity of some sectors. Aquaculture is considered the mainly developing food production sector globally. Protein insists is an important issue in human nutrition. Aquaculture has been an exercise for thousands of years, and it has now surpassed capture fisheries as the most vital source of seafood in the world. Limited study reports are available to focal point on bioactive immunostimulants in aquaculture applications. This review report provides information on the nutritional administration of bioactive immunostimulants, their types, functions, and beneficial impacts on aquatic animals' health as well as for the feed quality development in the aquaculture industry. The scope of this review combined to afford various kinds of natural derived bioactive molecules utilization and their beneficial effects in aquaculture applications.

1. Introduction

Bioactive immunostimulants have been utilized to improve aquatic animal health, a broad spectrum of bioactive active molecules are used to enhance aquatic animals' health in many ways such as antimicrobial, antioxidant, immunostimulant, growth-promoting, and anti-inflammatory properties as well as bioactive molecules applied in aquatic feed is the novel technique, this bioactive immunostimulant approach reliable and reproducing methods to enhance the feed quality [1–3]. In addition, dietary administration of various bioactive

immunostimulants namely FOS, butyric acid, propionic acid, polyphenol, chitosan, soybean isoflavones, lentinan, lipoteichoic acid, lactoferrin, and fucoidan have several positive effects such as immune system enhancement, growth, survival and disease confrontation against harmful microbes in aquatic organisms, natural bioactive immunostimulant dietary additive technique is the new way to enhance aquatic productions [4–8]. Microbial disease (bacteria, fungi, and viruses) outbreaks are considered the most imperative problem of aquaculture production, which can eventually lead to devastating financial losses [11]. The modes of action of immunostimulants are boosting the

* Corresponding author. Department of Animal and Aquatic Sciences, Faculty of Agriculture, Chiang Mai University, Chiang Mai, Thailand.

** Corresponding author. Xiamen Key Laboratory for Feed Quality Testing and Safety Evaluation, Fisheries College, Jimei University, Xiamen, China.

E-mail addresses: jmusunyunzhang@163.com (Y.-Z. Sun), hien.d@cmu.ac.th (H. Van Doan).

<https://doi.org/10.1016/j.fsi.2022.09.011>

Received 6 June 2022; Received in revised form 3 September 2022; Accepted 5 September 2022

Available online 10 September 2022

1050-4648/© 2022 Elsevier Ltd. All rights reserved.

immune system of aquatic animals and also improving the immunity level against harmful microbes, bioactive immunostimulants are already been used to control pathogens in shrimp and fish including bacteria (*Aeromonas hydrophila*, *Aeromonas salmonicida*, *Vibrio anguillarum*, *Vibrio vulnificus*, *Vibrio salmonicida*, *Streptococcus* sp., *Yersinia ruckeri*) and viruses, which cause diseases such as *hematopoietic necrosis*, *yellow head virus*, *viral hemorrhagic septicemia* and the parasite *Ichthyophthirius multifiliis*, these kinds of immunostimulants are supportive to boost survival and reduced mortality rate in aquatic animals [9,10]. Aquaculture is a vital and rapidly developing sector as it performs a very important responsibility to attain widespread protein foodstuff demands. The responsibility of aquaculture to expand the socio-economic implication of some places is mainly obvious since it's not only able to provide essential nutrients but also a variety of employment opportunities are created [12]. This review provides information about dietary administration, functions, and doses of different bioactive immunostimulants to increase healthy aquatic production and also enhance feed quality levels in the aquaculture industry.

2. The roles of bioactive immunostimulants in aquaculture

2.1. Bioactive immunostimulants improve growth and digestive

Bioactive immunostimulants are present in plants, animals, and microbes or can be synthetically produced; the natural derived bioactive immunostimulants have various biological actions such as antioxidant, antitumor, cytotoxic, an inducer of apoptosis and cell cycle arrest, antimicrobial, larvicidal, chemotactic, antifouling, antimelanogenic, and anti-inflammatory activities [13]. Bioactive immunostimulants can be classified into two types such as macromolecules (carbohydrates, proteins, nucleic acids, lipids, and polysaccharides) and micro molecules (natural products, primary and secondary metabolites), both the biomolecules play a major role in aquaculture. Bioactive immunostimulants are complex organic molecules, which are present in all living organisms, while such bioactive immunostimulant substances maintain the metabolic process of all living organisms [15]. Moreover, bioactive immunostimulants are shaped in the body and maintain the structure and growth, while they could be characterized as necessary for body functions [16]. The bioactive immunostimulants contain numerous positive responses such as the immune system strengthening, antioxidant activity, mediation of hormones, and butyric acid production in the colon as well as combination and/or dilution of structure in the gut, the small intestine is not only the fatal position for nutrient digestion and absorption but also strongly connected with a different region of intestinal antigens and bacteria to control gut and whole-body health [14]. Similarly, Ashraf et al. (2020) defined bioactive immunostimulant substances as the material that can cause stimulating effects on living tissue [17].

2.2. Bioactive immunostimulants modulate gut microbiota

Dietary immunostimulants feed additives were described as of quite chemical characteristic nature, containing nutritional and non-nutritional possessions as well as bioactive immunostimulants, which may connect with direct or indirect physiological responses of fish species [18]. Moreover, dietary feed additives of carvacrol, and/or thymol koi carp (*Cyprinus carpio*) [20] from (*Allium sativum*) garlic rainbow trout (*Oncorhynchus mykiss*) [21] have revealed noteworthy possessions on gut health, digestive enzyme activity, intestinal immunity, and extra supportive photogenic targets for the progress of aquatic feed, the dietary immunostimulant additive stimulates and modifies the intestinal microbiota conferring health benefits of the host, but notably down-regulated the relative manner of tumor necrosis factor α (TNF- α) ($P = 0.0002$) and transforming enlargement factor β (TGF- β) and remarkably decreased malonaldehyde (MDA) levels in fish species. Heterogeneous groups of functional feed additives, such as essential oils

(derived from aromatic plants) are widely used to enhance animal nutrition since these bioactive immunostimulants are applied in livestock nutrition as well as in animal feed. Essential oils are an important bioactive immunostimulant group; as such molecules are potential chemotherapeutic alternatives without side effects like immunostimulant, antimicrobial, anti-stress, antioxidant, and growth-stimulating effects while there is no ecological or harmful side effects linkage. Additionally, essential oils may exhibit a positive impact on the gut health of livestock as well as fish species [22]. Additionally, the colonization of gut microbiota has an important function in the regulation of non-specific and specific pathways of the immune system [24]. The intestinal microbiota of fish has an extremely imperative responsibility due to gastrointestinal growth, higher digestive enzyme action, mucosal tolerance maintenance, immunoregulation, and infection treatment enhancement [25–28]. Similarly, the gut microbial composition changes in the host's microbiota through the utilization of bioactive immunostimulants from various sources [23]. Different types of dietary functional feed additives such as probiotics, prebiotics, phytogenics, and immunostimulants are used to enhance healthy aquatic animals' production and also enhance the feed quality in aquaculture, the dietary immunogen additive increased growth performance, feed utilization, protein efficiency, and also decrease feed conversion ratio consequently, dietary immunogen additive increased higher body protein content [19].

2.3. Bioactive immunostimulants improve immunity

The nutritional additives of β -glucan, chitosan, or raffinose have been used to improve the non-specific immune system of koi (*Cyprinus rubrofasciatus*) by amplifying various resistant criteria such as phagocytic activity, respiratory burst activity, lysozyme action, WBC, and SOD action, finally, the bioactive immunostimulants are supportive to get better the immune system of aquatic animals [31]. MyD88 signaling pathway is directly triggered by MAMPs of intestinal microbiota strains. Different kinds of substances (lysozyme, protease, complements, lectins, and immunoglobulins) are significant for skin mucosal immunity and also for disease resistance in fish [29]. For example, the MyD88 signaling pathway plays a major part in the innate immune system control, while on other hand some microbial components controlled the adaptive immune system, increased digestive enzyme, protein content, and reduced lipid content level, there are no differences in immune gene expression those all the parameters confirmed by the control groups (IgM and IgT) of rainbow trout (*Oncorhynchus mykiss*) head kidney [30]. Fishes treated with bioactive immunostimulants have generally confirmed by phagocytic cell performance increase [32], as according to literature the phagocytic cell action can be better by bioactive immunostimulants due to the activation of a natural immune response, for example, the inflammatory reaction before adaptive response such as antibody production, excessive doses level of some immunostimulants provoke immunosuppression in fish. The side effects of immunostimulants have not been well-studied. The study concluded that immunostimulants are a vital role in to control of fish infections and the optimum dietary immunostimulants are helpful to improve the growth of farming fish culture [33]. Mendoza Rodriguez et al. (2017) examined that dietary additives of organic acids (polyhydroxy butyrate (PHB 2%), potassium diformate (KDF 0.6%)), other biomolecules (carrageenan (0.5%) and alginic acid (1%)) enhanced the immunological system of red drum (*S. ocellatus*), as well as feed efficiency and weight gain, was notably decreased in fish feed compared to the basal diet and together bioactive immunostimulants [34]. Consequently, the bioactive immunostimulant supplements have been shown to noticeably resilient structure stimulation of fish [35]. In aquaculture, bioactive immunostimulants and their derivatives enhanced non-specific immune responses against various diseases, while these molecules have exhibited effective immunostimulant properties in fish and shrimps [10].

2.4. Bioactive compounds as immunostimulants in aquaculture

2.4.1. Plants derived bioactive immunostimulants

Plant-based bioactive immunostimulants as supplements could show the potential impact in aquatic organisms like enhanced gut health, immune system, antimicrobial action and act as appetite stimulators, the herbal bioactive immunostimulants only give positive effects; they cannot be suggested owing to their remaining and other side effects in aquatic animals [36]. Similarly, the active dietary supplementation of herb and spices were capable to provoke and release intestinal enzymes, mucosal serum immunities, and antioxidant response that will stimulate the appetite; thus, rising food consumption and subsequently improving the efficiency in Nile tilapia (*Oreochromis niloticus*), this dietary additive has no significant changes in feed conversion ratio and also dietary inclusion was not affected to fish health [37]. In addition, various plants derived bioactive elements such as flavonoids, alkaloids, phenolics, terpenoids, tannins, steroids, glycosides, saponins, and essential oils have been described to show positive effects namely immunostimulation, growth promotion, anti-pathogenic, appetite stimulation, and anti-stress in fish species, this study pointed out that dietary additives of herbal bioactive immunostimulant as helpful to enhanced physiological and hematological responses in fishes [36]. The plant-based bioactive immunostimulants were commonly tonics for the release of digestive enzymes and have a direct effect on intestinal microflora, while according to literature these compounds can increase the digestibility, feed conversion, and protein synthesis of the rainbow trout (*Oncorhynchus mykiss*) [38]. Different types of plant bioactive immunostimulants have been added to feed supplements, demonstrating significant impacts in farmed animals, while such dietary additives are safe for animals, consumers, and the environment [39]. Abha and Abhisweta (2020) reported that the animal feed industry utilized plant bioactive immunostimulants to develop animal nutrition and health [40]. Additionally, Baleta et al. (2022) suggested that dietary additives of 1% ethanolic katuk extract bioactive immunostimulant enhanced growth, appetite, and improved food utilization in Orange-spotted grouper (*Epinephelus coioides*) [41].

2.5. Polyphenols

The polyphenols group is one of the major classifications of bioactive immunostimulants produced by plants to defend against different pathogenic bacteria, fungi, and viruses and also defend response to various types of abiotic stress [42]. Therefore, polyphenols are suggested as alternative options for the (partial) replacement of chemicals and antibiotics in aquaculture development [43,44]. In addition, polyphenol dietary additive (tannins from 2.5 to 20.0 g/kg) was used for immunological parameters enhancement in rohu (*Labeo rohita*) fingerlings, while these supplements have shown numerous positive effects such as resistance against oxidative stress improvement, reproductive performance, growth performance and infection resistance in different fish species such as European sea bass (*Dicentrarchus labrax*), black carp (*Mylopharyngodon piceus*), Wuchang bream (*Megalobrama amblycephala*), common carp (*Cyprinus carpio*), convict cichlid (*Amatitlania nigrofasciata*) and beluga sturgeon (*Huso huso*) [10,45–51]. Dietary polyphenol supplement (0.2%) was extracted from chestnut and olive mill wastewater, this polyphenol additive improved the growth performance, immunological parameters, and antioxidant status, and also feed conversion ratio of fish-fed polyphenols notably diminished and differentiated from the control groups in common carp (*Cyprinus carpio*) fingerlings [52]. Dietary *Clarius lazera* phenol additives have been used in various concentrations (35, 75, 150, and 300 mg/L) and enhanced hematological parameters and also elevated LDH, glucose, and cortisol, and significantly reduced serum cholesterol concentration in catfish (*Siluriformes*). The dietary polyphenol was supportive to improve the fish's immune system [53]. In addition, dietary tannins polyphenol additive (0.1%) was derived from chestnut wood and were in various analyzes such as growth parameters, growth gene expression, enzymatic

activity, and hematological and non-specific immune parameters in beluga under experimental conditions. Finally, the optimum level of 0.1% dietary polyphenol was supportive to improve fish health [51].

2.6. Flavonoids

Flavonoids are the most important ingredients of plants [54]. Dietary *Allium mongolicum* Regel flavonoids (AMRF) supplement (40 mg/kg) enhanced the immune system, antioxidant status, regulation of gene activation, and infection resistance of *Cephalopholis argus* (Peacock hind) against *Aeromonas hydrophila* as well as considerably decreased in serum cortisol, liver MDA content, serum ALT and AST activity [55]. Dietary supplementation of *R. verniciflua* extract flavonoids enhanced feed quality level and fish production, exhibiting also a significant impact on antiviral effects in fish viral diseases like infectious hematopoietic necrosis virus (IHNV) and viral hemorrhagic septicemia virus (VHSV) [56].

2.7. Alkaloids

Alkaloids are found in plant tissues as a type of water-soluble salts of organic acids, esters, or sugars rather than as free bases [57]. Moreover, alkaloids contain different types of protein bands, which have a significant role in immune system enhancement and infection resistance against various pathogens in fish [58]. Subsequently, dietary additives of *Gelsemium elegans* alkaloids (40 mg/kg) increased intestinal villus length, villus number, muscle growth, antioxidant activity, intestinal immune activity, cytokine-related genes expression, and also to improved the disease resistance of Wuchang bream (*Megalobrama amblycephala*), while firmicutes abundance was reduced at phylum level after the alkaloid feeding [59]. Zhou et al. (2016) reported that dietary additives (25 g/kg) of the fibrous root of *Rhizoma Coptidis* containing alkaloids could enhance the immune system and disease resistance in common carp (*Cyprinus carpio*) [60].

2.8. Fructooligosaccharides

FOS is derived from blue agave plants, vegetables, fruits, cereals, and grains [61], composed of D-fructans, in which fructose is attached by b-(2–1) glycosidic connections and associated with an end glucose unit. Additionally, FOS may be linked via tolls like receptors (TLR2) pointing to macrophages [62] and stimulating the expression of antimicrobial peptides (Leap), which have a significant role in natural immune protection and disease resistance in fish (*Megalobrama amblycephala*) [63]. Likewise, dietary supplementation of FOS (1%, 2%, and 3%) increased lysozyme action, salinity stress resistance, leukocyte production, and elevated digestive enzyme function in Caspian roach fry (*Rutilus rutilus*) [64]. Similarly, our research group described that diet containing (5 g/kg) FOS improved intestinal enzyme activity, and lysozyme action, and also decreased triglyceride (TG) levels compared to the control diet in Japanese flounder (*Paralichthys olivaceus*) [59,65]. Dietary supplementation of FOS (0.3 and 0.6%) has shown a beneficial effect on the innate immune response of Nile tilapia (*Oreochromis niloticus*), confirmed by the major progress of serum lysozyme action and IgM level and also both plasma and phenoloxidase performance were noticeably ($P < 0.05$) affected only by dietary FOS levels with the maximum standards experiential in fish fed 0.6 and 0.3% FOS, respectively [66].

2.9. Galactooligosaccharides

Galactooligosaccharides (GOS) incorporate 2–20 different prebiotic substances of galactose and glucose and are generally formed by the enzymatic hydrolysis of lactose [67]. GOS has a small point of polymerization, which improves its metabolism rate by probiotics [68]. The use of GOS (1% and 2%) dietary supplement enhanced humoral and mucosal immune response, survival rate, and increased salinity resistance in Caspian roach (*Rutilus rutilus*) fry as well as there is no negative

impacts in the experimental condition [69]. Nutritional management of GOS as an immunostimulant in rockfish (*Sebastes schlegelii*) presented that feeding rockfish with (1%) of GOS for 8 weeks, notably improved skin mucus as well as humoral immune criteria, and also this bioactive immunostimulant not affected to the physiological and hematological responses in fish species [70]. Yukgehnash et al. (2020) reported that the dietary supplement of GOS (10 g/kg⁻¹) acted as a significantly important prebiotic for the Atlantic salmon (*Salmo salar*) and also this bioactive immunostimulant affected protein concentration in the wet body and protein retention were reduced by 6 and 9%, respectively, relative to the basal fish [71], Caspian roach (*Rutilus rutilus*) [72], rainbow trout (*Oncorhynchus mykiss*) [73], common carp (*Cyprinus carpio*) [25,74], Caspian white fish (*Rutilus frisii kutum*) [75], narrow-clawed crayfish (*Astacuseptodactylus leptodactylus*) [76], and rockfish (*Sebastes schlegelii*) [70].

2.10. Inulin

Inulin exists in several fruits, vegetables, and cereals, like leeks, onions, chicory, bananas, wheat, garlic, and artichokes. It is reported that dietary inulin supplement (5 g/kg) could improve hematocrit and NBT activity levels and lysozyme activity as well as there is no negative impact on physiological and hematological response in Nile tilapia (*Oreochromis niloticus*) [77]. Gupta et al. (2020) stated that dietary inulin additives can control the hematological and/or serum biochemical parameters (erythrocyte counts, mean corpuscular volume (MCV), hematocrit, mean corpuscular hemoglobin (MCH), hemoglobin, mean corpuscular hemoglobin concentration (MCHC), leukocyte counts, serum glucose or serum total protein levels, differential leukocyte counts and serum cholesterol); thus this indicator has a prominent role in fish health [78]. Furthermore, Mo et al. (2015) presented that dietary inulin additive (2%) improved growth action and the natural immune system of grass carp (*Ctenopharyngodon idella*) [79]. Another study about inulin supplement (15–20 g/kg) in Barramundi (*L. calcarifer*) juveniles; reported significant effects in histology, biochemical parameters, and immunohistology [80].

2.11. Salidroside

Salidroside is a glucoside of tyrosol, which is present in the plant species of *Rhodiola rosea*. It has been considered, along with rosin, as one of the most significant factors responsible for the known antidepressant and anxiolytic actions of this plant [81]. Zhang et al. (2021) reported that salidroside was found not only in the plant *Rhodiola rosea* but also in a few Chinese herbs like *Angelica root* [82], *Ganoderma lucidum* [83], *Prunella vulgaris*, etc., as well as this bioactive immunostimulant enhanced the immune system and antiviral effects challenging against coxsackievirus B3 and dengue virus [84]. Xie et al. (2020) pointed out that salidroside (0.1% and 1.0%) components were highly present in *Prunella vulgaris* extracts, and these factors enhanced non-specific immune activity, respiratory burst activity, phagocytic activity reduced mortality rate, and treat Olive flounder (*Paralichthys olivaceus*) challenge with *Uronema marinum*. Finally [4], Yang et al. (2020) mentioned that dietary additive of salidroside improved the immune system in crucian carp (*Carassius carassius*) and enhanced the defense action against harmful microbes, main at the concentration of 100 mg/kg for four weeks. The protective effect of salidroside on crucian carp (*Carassius auratus*) might be considered as an alternative to antibiotics for controlling fish infections in aquaculture as well as the quantity of *A. hydrophila* in the kidney and spleen was considerably decreased in salidroside additive diet groups ($P < 0.05$).

2.12. Geniposide

Geniposide is a bioactive iridoid glycoside; it is present in different types of medicinal herbs, namely *Gardenia jasminoides* (fruits).

Geniposide showed a variety of pharmacological responses both *in vitro* and *in vivo* applications, such as antidiabetic, anti-inflammatory, neuroprotective, hepatoprotective, analgesic, antidepressant-like, cardioprotective, antioxidant, immune-regulatory, antithrombotic, and antitumor activity. Geniposide is derived from *Gardenia jasminoides* Ellis, a Chinese herbal medicine [85]. Nevertheless, few studies demonstrate the efficacy of geniposide as an immunostimulant in fish species, dietary administration of geniposide 400–800 mg/kg increased total collagen and alkaline-insoluble collagen content in grass carp (*Ctenopharyngodon idella*) muscle than control and also fish fed diet containing geniposide notably reduced muscle crude lipid content compared to control groups [86]. Geniposide was shown as an efficient diet supplement to control the apoptosis-related factor, reduce WSSV (white spot syndrome virus) simulation and increase the survival rate of WSSV-challenged red swamp crayfish (*Procambarus clarkii*) [87]. Similar beneficial effects of lysozyme reaction were also observed in crucian carp (*Carassius carassius*) [88], and Nile tilapia (*Oreochromis niloticus*) [66] feeding geniposide-containing food. According to the literature, the use of 100 mg geniposide/kg was proved positive effects on immune reaction improvement and infection resistance in crucian carp (*Carassius carassius*), one of the most (freshwater) cultivated fish species in China. Furthermore, this dietary supplement was tested for innate immune response improvement, upregulated resistant connected gene exhibition, and infection resistance against *Aeromonas hydrophila* in crucian carp (*Carassius carassius*) [89].

2.13. Lectins

Lectins are carbohydrate-binding proteins, present in most plants and some animals [90]. Lectins are considered as a carbohydrate detection domain, mainly binding sugars while there are different classifications of animal lectins namely C-type, I-type, F-type, L-type, P-type, M-type, R-type, F-box lectins, chitinase-like lectins, galectins, intelectins, calnexin, and ficolins. Lectins play a major role in intracellular and extracellular, extracellular lectins are vital for cell signaling and pathogen recognition while intracellular lectins play a major part in the convey of proteins movement throughout the cells and in protein sorting [91]. C-type lectins, which are responsible for stimulating the immune system, are particularly linked with PAMPs on the surfaces of several harmful microbes, providing them the ability to detect a broad diversity of pathogens [92]. Several classifications of lectins have been recognized in rainbow trout serum through calcium ion chains to *A. salmonicida* lipopolysaccharide [93]. On the other hand, opsonizing lectins, which are present in fish serum, may be beneficial for fish health, as are considered for the immune mechanism and prevention of aquatic animal diseases. The components play an imperative function in protecting the host by identifying pathogens and then increasing their elimination and appearance in phagocytic cells of the immune system. The mammalian mannose-associated lectin can opsonize in different types of organisms such as yeasts, bacteria, and surrounded viruses, like HIV or influenza [94]. Similarly, the potential function of lectins as non-recognized substances in all living organisms' immunity [95]. Likewise, the crucial role of lectins in molecular biology and immunological analysis in animals [96], while the importance of lectins in the innate immune reaction of mammals is well known [97]. In addition, Liang et al. (2020) pointed out that lectins from immune cells of marine invertebrates were effective in improving disease resistance against *V. vulnificus* and *V. pelagicus* in aquatic animals and also reduced mortality rate in aquatic animals [98].

2.14. Carotenoids

Carotenoids are natural pigments, with a significant role in all living organisms like plants, algae, fungi, bacteria, and animals. Animals cannot synthesize carotenoids, so they obtain them from their diet [99]. Shrimp-extracted carotenoid supplements of 100 and 200 mg/kg could

lead to immune system enhancement and infection resistance of Carp against *A. hydrophila* [100]. The addition of carotenoids in feed exhibited a beneficial role in both fish and fishery products as pathogenic organisms' inhibition was achieved. Carotenoids' dietary additives range (50 and 100 mg/kg) varies from species to species [101]. Many researchers exhibited that a dietary additive of 100 mg carotenoids/kg can boost the immune system, survival, and growth as well as avoid harmful effects of lipid peroxidation in different species such as rainbow trout (*Oncorhynchus mykiss*), giant tiger prawn (*Penaeus monodon*) and European bullhead (*Cottus gobio*) [102–104]. Natural immunostimulants, namely dietary supplements of carotenoids boosted the immune system, disease resistance, and antioxidant function of aquatic animals and also downregulated in superoxide dismutase but high catalase activity compared to the control one [105]. The function of plants based bioactive compounds as immunostimulants in aquatic animals has been summarised in Table 1.

2.15. Animal originated bioactive immunostimulants

2.15.1. Chitosan

Chitosan is a type of alkaline polysaccharide existing in the shell of aquatic organisms namely crabs, shrimps, and shellfish. Chitosan is a conventional non-toxic, safe, bio-compatible, and bio-degradable polymer substance. The main effect of chitosan molecule progression is to significantly increase the growth of aquatic species [106], reduce the growth of harmful microbes [107], purify the aquaculture water [108], and improve the resistance of aquatic organisms to infections [106]. Chitosan can chelate metal ions via the role of amino acid groups and hydroxyl substances and might decrease the heavy metal ions in water; thus chitosan is considered a major defender factor for the aquatic species [109]. In addition, immunostimulants are responsible for the improvement of inflammatory cell function namely polymorphonuclear

leucocytes, cytokine, and macrophages [110]. In aquaculture, dietary chitosan improved the protective effect against bacterial infections and enhanced the immune response in hybrid striped bass (*Morone saxatilis*) [111], while recently, chitosan was presented as an effective immunostimulant in different fish species [112]. Dietary supplement of chitosan enhanced non-specific immune response under experimental conditions of healthy control, and additionally, chitosan was tested against the immune response modulation in rohu (*Labeo rohita*) [113]. Diets with chitosan for rainbow trout (*Oncorhynchus mykiss*) [114], olive flounder (*Paralichthys olivaceus*) [115], koi (*Cyprinus rubrofuscus*) [116], and kelp grouper (*Epinephelus bruneus*) [107] demonstrated that such a bioactive immunostimulant could improve growth, natural immunity, infection, stress resistance, enhance immunological criterion and increase water quality. The mechanism of chitosan action as an immunostimulant in fish has not been fully established yet. Similar studies showed that chitosan additives might increase the innate immune system response of various fish species [117]. Oushani et al. (2020) reported the action of chitosan as an immunostimulant in rainbow trout (*Oncorhynchus mykiss*), as they observed that a chitosan (2.5 g/kg) diet was favorable on immunological criteria and increased the survival rate of rainbow trout (*O. mykiss*) against environmental stress [114]. Wu et al. (2020) reported that the immunostimulatory feature of chitosan for olive flounder (*Paralichthys olivaceus*) activates the innate immune system of fish species, thus this action exhibits a defense role in the fish body through amino moieties [115].

2.16. Lactoferrin

Lactoferrin is a globular glycoprotein with a molecular mass of about 80 kDa that is widely represented in different secretory fluids namely milk, saliva, tears, and nasal secretions. Lactoferrin is one of the significant bioactive mechanisms of the immune system in the body; it has

Table 1

Plants originated bioactive compounds as immunostimulants in aquaculture.

Origin	Bioactive immunostimulants	Aquatic organisms	Immune response	References
Plant	Fructooligosaccharides	Caspian roach	↑lysozyme function, improvement of IgM level, No significant different ACP and increases PO	[63,65]
		Common carp		
		Nile tilapia		
		Japanese sea bass	↑growth, digestion and immune activities	[190]
		Tambaqui	Promote growth and improved healthiness	[191]
	Galactooligosaccharides	Juvenile grouper <i>Epinephelus coioides</i>	Improved immune system and intestinal morphology	[192]
		Caspian roach	↑ Humoral and mucosal immune reaction	[69,71,72, 75]
		Rainbow trout		
		Narrow-clawed Crayfish		
		Common Carp	Controlled innate immune response in skin mucosa	[193]
	Inulin	Pacific white shrimp	Modulate gut microbiota composition, reduce stress and enhanced immune system	[194]
		<i>L. calcarifer</i>	↑ hematocrit, NBT and lysozyme activity, biochemical parameters and immuno-haematology	[63,79]
		Rainbow trout		
	Salidroside	common carp	↑growth performance, innate immune response and antioxidant activity	[195]
		Nile tilapia	Improved fish health under salinity stress condition	[196]
	Geniposide	Crucian carp	↑ immune related gene and non specific immune system	[4,105]
		Paralichthysolivaceus		
	Polyphenols	Crucian carp	↑ innate immune response, upregulated immune gene expression, RPA	[65,188]
		Nile tilapia		
		Crayfish	Anti inflammatory, antioxidant and antiviral activity	[197]
	Flavonoids	Common carp fingerlings	↑haematological parameters(erythrocytes and haemoglobin), anti oxidant status	[51]
		<i>Amatitlania nigrofasciata</i>	Improved humoral and mucosal response and antioxidant activity	[198]
	Alkaloids	<i>Lates Calcarifer</i>	Improved growth, mucosal immune response and liver oxidative status	[199]
		<i>C.argus</i>	↑immune system, antioxidant activity, regulation of immune gene expression	[7,54]
	Alkaloids	<i>Trachinotusovatus</i>		
		<i>Ctenopharyngodon idella</i>	↑Antioxidant, anti inflammatory and immune response	[200]
		<i>Megalobramaamblycephala</i>	↑antioxidant activity, Intestinal immune activity, RPA	[58,59]
		common carp		
		Pacific white shrimp	Improved growth response, survival rate, immune response and antibacterial activity	[201]

↑- Enhance or Increase or Improve.

antimicrobial activity (bactericide, fungicide) and is part of the innate defense, mainly against mucose. Lactoferrin connects with DNA and RNA, polysaccharides, and heparin, and performs the biological response in complexes with these ligands [118]. The dietary additive of 100 mg/kg Lf enhanced the immune response, respiratory burst, leukocytes peroxidase content, and phagocytic index in gilthead sea bream (*Sparus aurata*) [119]. [120] Morshedi et al. (2020) observed that the dietary supplement (1200 mg/kg) of Lf could enhance low salinity stress, tolerance of air exposure, natural immune response, and mucus secretion of juvenile orange-spotted grouper (*Epinephelus coioides*), as well as dietary LF, did not affect the growth rate of this species under the condition applied in this study. Similarly [35], Luna-Castro et al. (2021) represented that the nutritional additive of Lf at a level of 100 mg/kg increased the innate immunity and infection resistance of sea bream (*Pagrus major*) and Asian catfish (*Siluriformes*), as well as this study, described that support the possible use of Lf as an immunostimulant for farmed fish species. Several Lactoferrin (Lf) products have been classified and evaluated, but bovine lactoferrin (BLf) has received attention for its potential for animal protection. Owing to its immunoregulatory function, BLf may stimulate antiviral, antifungal, and antibacterial functions and subsequently may activate efficient defense against different microbial infections in some fish species according to the dose and administration methods [121]. Additionally, dietary Lf supplementation of 100 mg/kg for 1 week the immune system improvement of diverse fish species such as goldfish (*Carassius auratus*), rainbow trout (*Oncorhynchus mykiss*), Atlantic salmon (*Salmo salar*), red sea bream (*Pagrus major*) and Asian catfish (*Siluriformes*) [122]. The function of animals-base bioactive compounds as immunostimulants in aquatic animals has been summarised in Table 2.

2.17. Bacterium originated bioactive immunostimulants

2.17.1. Peptidoglycan

Peptidoglycan or murein is a polymer consisting of sugars and amino acids that forms a mesh-like peptidoglycan layer outside the plasma membrane of most bacteria, forming the cell wall. The sugar component consists of alternating residues of β -(1, 4) linked N-acetyl glucosamine (NAG) and N-acetylmuramic acid (NAM), and attached to the NAM is a peptide chain of three to five amino acids. Peptidoglycan performs the structural function, strengthens the bacterial cell wall, and prevents the

osmotic pressure of the cytoplasm. Peptidoglycan was also involved in the binary division during bacterial cell reproduction [123]. Oral dietary supplements of peptidoglycan (0.2 mg/kg) extracted from *Bifidobacterium thermophilum* could act against vibriosis and white spot syndrome in kuruma shrimps (*Marsupenaeus japonicas*) as well as reduce mortality and increase the survival rate in shrimp [124]. Likewise, dietary additives of cellular components like cell wall (CW), peptidoglycan (PG), and lipoteichoic acid (LTA) enhanced physiological and immunological response, while notably decreased feed conversion ratio (FCR) was noticed in all the treatments compared with the control group in orange-spotted grouper (*Epinephelus coioides*) [125]. [126] Kondo et al. (2021) demonstrated that dietary additives of A3a-PG (8 g A3a-PG/kg) led to the immune system enhancement, disease resistance, and growth performance of Japanese flounder (*Paralichthys olivaceus*) as well as the highest dose explored in the previous report, (16 g A3a-PG/kg), also failed to show any significant influences, although no obvious pathology was observed. Another case of dietary additive of peptidoglycan (0.18 g/kg for 5 days/week) exhibited an increase in the immune system and infection resistance of black tiger shrimp (*Penaeus monodon Fabricius*) and there is no significant changes to the growth [127]. The immunostimulatory effect of microbial peptidoglycan is acknowledged from the laboratory verification process, which can stimulate innate immune responses, phagocytic index, and survival rate and infection resistance against aquatic pathogens in aquatic animals [128].

2.18. Lipopolysaccharides

Lipopolysaccharides (LPS) are huge molecules incorporating a lipid and a polysaccharide controlled of O-antigen, outer core, and inner core connected with a covalent bond, while they are attending in the outer layer membrane of the Gram-negative bacteria. LPS are the components of the gram-negative bacteria cell wall and also turn on B cell proliferation. LPS injected into red sea bream (*Pagrus major*) exhibited improved macrophage phagocytic action [129]. The LPS are very effective even in extremely low doses; while they can be derived from bacteria by chemical extraction methods and present better immunostimulating properties during their use in fish species [130]. Ali et al. (2021) suggested that LPS (1250 μ g) played a significant role in rainbow trout (*Oncorhynchus mykiss*) and carp fish (*Cyprinus carpio*) species, as they are good immunomodulators and beneficial response in the disease resistance of fish species, the lipopolysaccharides improved phagocytic activity, total serum Ig and get better mucus lysozyme compared to untreated fishes mortality level is increased [32,131]. Abd El-Kader et al. (2021) recommended that dietary supplement of LPS (50 and 100 μ g of LPS/fish) was injected intraperitoneally and successfully activated innate cellular and secondary immune response as well O-specific linkage for activating the adaptive immune system and disease resistance against *A. hydrophila* [132]. In contrast, oral consumption did not alter any of the aforementioned criteria. Similarly [133], Holen et al. (2021) reported that dietary additive of lipopolysaccharides was derived from *Salmonella Typhimurium*. The lipopolysaccharide component enhanced better immunostimulant, phagocytic activity, and disease resistance against *Escherichia coli* and *A. salmonicida* disease in trout (*Oncorhynchus mykiss*). In addition [134], Mohammadi et al. (2020) reported that a nutritional LPS additive of 0.1 mL was derived from *Aeromonas hydrophila* enhanced disease resistance against *A. hydrophila* in (*Oreochromis niloticus*) Nile tilapia as well as for disease prevention during unfavorable environmental conditions.

2.19. Lipoteichoic acid

Dietary supplementation of lipoteichoic acid (LTA) was derived from probiotics and is used to regulate immune gene expression as well as in ligand binding connections. Since LTA can stimulate the immune response in fish, it might be favorable for the control of disease in Gram-positive bacteria such as *Streptococcus iniae* and *Streptococcus agalactiae*

Table 2
Animal originated bioactive compounds as immunostimulants in aquaculture.

Origin	Bioactive immunostimulants	Aquatic organisms	Immune response	References
Animal	Chitosan	Rainbow trout	\uparrow haematological parameters, RPA	[105, 112–114]
		Olive flounder		
		Koi kelp grouper		
	Lactoferrin	Asian catfish Atlantic salmon	\uparrow non specific immune response, respiratory burst, leukocytes peroxidase content and phagocytic index, RPA	[34,117, 118]
	Carotenoids	gilthead sea bream Carp rainbow trout giant tiger prawn European bullhead	\uparrow antioxidant activity, RPA. immune system response, Disease resistance	[98, 100–102]

\uparrow - Enhance or Increase or Improve.

[135]. Similarly [136], Yang et al. (2019) observed that dietary additives of both peptidoglycan and LTA extracted from *B. pumilus* SE5 can stimulate TLRs/MyD88 signal pathway and the condition of antibacterial agents and thus form the gut microbiota in orange-spotted grouper (*E. coioides*). Additionally, dietary additive of LTA extracted from probiotic *B. pumilus* SE5 was utilized to uncontrolled three antibacterial peptides genes (epinecidin-1, hepcidin-1, and β -defensin) and immune gene expression (TLR2, NOD2, IL-8, IgM) in the head kidney of Grouper (*Epinephelus coioides*) [137]. Earlier studies have revealed that LTA is the main cell wall element of Gram-positive bacteria, can associate with the TLR-2 receptor, and triggers a cascade of responses. *L. Plantarum* LTA extensively improved the feature of TLR2 and *V. anguillarum* flagellin particularly improved TLR5. It is differentiated that LTA, flagellin, and LPS bacterial components are mainly stimulated by TLR2, TLR5, and TLR9 receptor pathways [138,139]. [140] Gao et al. (2016) suggested that dietary LTA enhanced the natural immune system of silver pomfret (*Pampus argenteus*). Probiotic dietary additives of LTA can regulate the activation of several inflammatory factors and provoke apoptosis to a lower level, as might also be determined by the physiological characterization of the bacterium. Dietary supplements of LTA an imperative function in the host cell TLR and also enhanced immune response, probiotic colonization, and decreased pathogenic load while sustaining a strong and active strength among the host and its gut bacteria.

2.20. Levan

Levan is a naturally occurring fructan, the homopolymer of fructose. The molecular weight and degree of branching of levan differ depending on the production organism. Moreover, the molecular weights of bacterial levan are much higher than the plant levan due to the multiple branching [141]. Levan is the component of natural polymeric chain linkages, which are mainly present in the microbial extracellular products of *Bacillus subtilis*, *Bacillus polymyxa*, *Zymomonas mobilis*, *Acetobacter xylinum*, and a few plant species [78]. Similarly, dietary additives (25 g/kg) of microbial levan exhibited beneficiary effects in aquatic animals like improved non-specific immune response, disease resistance, growth, survival, and enhanced tolerance to thermal stress, and also levan additive diets notably reduced the count of total viable aerobic bacteria and *Vibrio* spp. in the gut region of orange-spotted grouper (*Epinephelus coioides* H.) [142]. Moreover, the dietary additive of microbial levan (1.25%) enhanced the immune response and thermal stress tolerance effect of rohu (*L. rohita*) juveniles, this optimum dietary additive can confer protection against thermal stress tolerance in fish species [143].

2.21. Fungal originated bioactive immunostimulants

Higher fungi are considered important sources of natural products and have also exhibited various biological properties like anti-microbial, anti-oxidant, immunostimulation, hypoglycaemic activities, and anti-cancer [144]. Numerous higher fungi (*Pleurotus ostreatus*, *Lentinula edodes*, *Grifola frondosa*, *Coriolus Versicolor*, *Ganoderma lucidum*, and *Schizophyllum commune*) have bioactive polysaccharides in their fruiting bodies; mycelium and broth culture [145]. Consequently, polysaccharides can significantly improve the growth and natural immune response of fish and shrimp species [146–148].

2.22. Lentinan

Lentinan is a biological active transformer, which is shown to be effective in protecting the host from oxidative damage without obvious adverse activity and also diet additives of lentinan were used to modify the gastrointestinal microbiota, which would be useful to sustain the epithelial barrier role and enhance intestinal immunity. Lentinan is a type of β -glucans with safe and pharmacological properties as well as immunomodulatory, anti-oxidant, anti-inflammatory, anti-microbial,

and metabolic regulatory functions [147,149]. The nutritional additive of lentinan (100 μ g/ml) has notably increased the intestinal anti-oxidant, and inflammatory capacity by enhancing the performance of SOD, CAT, and GSH-Px, and suppressing lipid peroxidation and also lentinan may be effective for the treatment of gut inflammation including IBD in juvenile taimen (*Hucho taimen*) [148,149]. [150]. Galindo-Villegas et al. (2016) reported that lentinan can induce anti-microbial activity and stimulate the immune system through *in vitro* methods. Moreover, dietary additives of lentinan (100–1000 μ g/ml) increased the immune system and disease resistance in Common carp (*Cyprinus carpio*) by *in vitro* methods [151].

2.23. Mannan oligosaccharides

Mannan oligosaccharides (MOS) are extracted from the cell wall of yeasts and recently researchers indicated the beneficial properties of MOS in fish and shrimp species [152]. The possible effect of MOS (0.6%) in aquaculture content is determined by the mannose receptor (endocytic receptor), while these receptors communicate with macrophages and endothelial cells to be identified together with individual glycoproteins and microbial glycan ligands [153]. MOS (0.6%) presented a related consequence on the function of lysozyme in Nile tilapia (*Oreochromis niloticus*), Japanese flounder (*Paralichthys olivaceus*), Rainbow trout (*Oncorhynchus mykiss*), and Channel catfish (*Ictalurus punctatus*), while the dietary additives of MOS were not affected in fish growth and immune system [154]. MOS as feed supplements (6 g/kg) in aquaculture of fish and shellfish may demonstrate to be supportive to the host by increasing innate immunity, although more studies described that it is needed to infer the appropriate level of MOS addition to diverse fish species and also MOS dietary supplement was significantly decreased lipogenic enzymes response in liver and resulted in a lower lipid vacuolization and more regular-shaped morphology of hepatocytes approximately the sinusoidal spaces, which could be associated to better energy and good feed utilization because of the improvement of feed utilization and the decrease of feed intake found in the current experiment [155]. The dietary additive of MOS (0.4%) notably activated hemolytic and phagocytic reactions in rainbow trout (*Oncorhynchus mykiss* Walbaum) [156], while an amount of 2 g MOS/kg for 60 days was used to evaluate certain immune components and immuno-associated enzymes in red sea bream (*Pagrus major*) as well as the specific immunostimulatory activity of MOS in the gastrointestinal tract of fish and also the performance of lysozyme, bactericidal and peroxidase in serum and mucus of fish fed MOS diets improved over the control [157]. Similarly, the dietary feed supplement of MOS (0.6%–1%) was experienced to increase blood protein levels in rohu (*Labeo rohita*) and Nile tilapia (*Oreochromis niloticus*), and also dietary additive of MOS at a 0.6% and 1% dietary level stimulates growth and survival in rohu (*Labeo rohita*) and Nile tilapia (*Oreochromis niloticus*). In difference, elevated addition levels of immunostimulants led to an immunosuppressive effect in rohu (*Labeo rohita*) and Nile tilapia (*Oreochromis niloticus*) [158].

2.24. β -Glucans

Glucans are primarily present in the cell walls of bacteria, fungi, algae, lichens, and yeasts they obtain a helical or spiral backbone suitable to specific intra-factor hydrogen bonding. These β -glucans (10 mg/kg) have been commonly expressed to control the immune function as well as blood neutrophils, oxidative radical production, and superoxide anion production in stimulated macrophages of *Litopenaeus vannamei* juveniles [159]. The main component of β glucans is high-quality immunostimulating factors. The abundance and nature of side chains strongly affect the capability of the glucans to interfere with binding to external receptors on the particular cells influencing the reaction of the glucans as an immunostimulant [160]. These changes may indicate the way to a mortality rate reduction of fish species against various pathogens. Innovative β -glucan receptors namely Dectin-1 and Toll-like

receptors on leukocyte surfaces of vertebrates have provided various insights into the functions of immune responses induced by β -glucans [161]. Oral dietary administration of yeast-derived beta-glucans was showed that numerous positive affects such as enhance phagocytosis, lysozyme levels, respiratory burst activity, and resistance against bacterial pathogens in several cultured fish species. The function of glucans have been extensively studied in aquatic animals and the findings indicate that β -glucans improved growth and significantly reduced mortality rate in certain species of aquatic animals [162,163]. β -glucans have been mentioned for their valuable role in several fish species such as red sea bream (*Pagrus major*) [164], Atlantic Salmon (*Salmo salar* L.) is utilized to improve growth, survival, immune system, disease resistance against aquatic pathogens and reduced mortality rate in aquatic animals [165].

The function of microbial-based bioactive compounds as immunostimulants in aquatic animals has been summarised in Table 3.

2.25. Algal bioactive immunostimulants

2.25.1. Alginate

Alginate was derived from brown seaweed, and it is a polysaccharide component controlled by β -1, 4-D-mannuronic acid (M), and C5-epimer α -L-glucuronic acid (G). A dietary additive of sodium alginate was used to increase the non-specific immune system and infection resistance in Tilapia (*Oreochromis niloticus*) [125,166]. In addition, a nutritional additive of sodium alginate (2 g/kg) enhanced the survival of grouper *E. cooides* by increasing its phagocytic activity, respiratory bursts, lysozyme activity, alternative complement activity, and SOD activity, disease resistance against *Streptococcus* sp. and iridovirus infection in grouper [167]. It has been reported that the nutritional additive of sodium alginate (2 g/kg) could improve phagocytic activity, respiratory bursts, lysozyme activity, alternative complement activity, SOD activity, and disease resistance against *V. alginolyticus* but decreased GPX activity in Pacific white shrimp (*L. vannamei*) [168].

2.26. Alginic acid

Alginic acid (or algin) is a naturally occurring, edible polysaccharide, present in brown algae. Such compound is hydrophilic and forms a viscous gum when hydrated. Accompanied by different metals such as calcium and sodium, forms salts, which are recognized as alginates. Its color ranges from white to yellowish-brown. Concerning its beneficial

results through consumption; previous studies demonstrated that alginic acid could be used to enhance the nonspecific immune response in fish species [125]. Dietary supplements of alginic acid (1 g/kg) improved immune response, growth performance, and infection resistance against WSSV disease in *P. monodon*, suitable dietary level of alginic acid is protecting immune system response against WSSV infection in juvenile *P. monodon* [169]. Similarly, ergosan is an algal derivative with alginic acid (2 g/kg), which could be used to improve innate immune response and infection resistance in different types of fish species [170]. Additionally, ergosan containing (0.5% and 1%) alginic acid derived from *Laminaria digitata*, could enhance the immune response, phagocytic activity, respiratory activity, and expression of interleukins and less response in a humoral immune parameter in rainbow trout (*Oncorhynchus mykiss*) [171]. The dietary additive of ergosan (5 g/kg) enhanced growth performance, protease, esterase activities, alkaline phosphatase, lysozyme actions, and disease resistance against *Yersinia ruckeri* in rainbow trout (*Oncorhynchus mykiss*) (110 g) [172].

2.27. Fucoidan

Fucoidan is a complex long-chain sulfated polysaccharide, present in different species of brown algae. Commercially available fucoidan is usually derived from the different seaweed species namely *Fucus vesiculosus*, *Cladosiphon okamuranus*, *Laminaria japonica*, and *Undaria pinnatifida*. Fucoidan (3% and 6%) promotes several biological effects like immunostimulation, disease resistance improvement, growth, and survival, whereas serum Albumin/Globulin (A/G) ratio and blood glucose level exhibited decreasing trend in diverse fishes and shrimps [173]. Saeed et al. (2021) reported that dietary fucoidan (200 mg/kg of body weight, group 12–15 g) boosted phagocytic activity and disease resistance of shrimps as well as overdoses of several immunostimulants to induce immunosuppression in fish [174]. The side effects of immunostimulants have not been well-studied. Dietary supplements of fucoidan (1 g/kg) increased the immune system (circulating hemocytes, PO activity, RBs, phagocytic activity) and disease resistance against *V. alginolyticus* in shrimps. The study reports concluded that dietary fucoidan's most important role is to enhance the immune system of shrimp species [175]. Finally, oral administration of fucoidan (15 g/kg) derived from *Sargassum polycystum* could reduce the infection of *White Spot Syndrome Virus* (WSSV) and reduce mortality rate, increase survival level in black tiger shrimp *P. monodon* [176].

Table 3
Microbial bioactive compounds as immunostimulants in aquaculture.

Origin	Bioactive immunostimulants	Aquatic organisms	Immune response	References
Microbes (Bacteria and fungi)	Peptidoglycan	kuruma shrimps	↑immune system response, RPA, antibacterial and antiviral effects	[122,124,125]
		Japanese flounder		
		<i>Penaeus monodon</i>		
	Lipopolysaccharides	sea cucumber	Potential immunomodulatory factor	[202]
			Regulation of innate immune response	
		spotted sea bass		[203]
	Exopolysaccharides	Red sea bream	↑phagocytic activity and disease resistance against antibacterial activity	[127,131]
		trout		
		<i>Channa argus</i>	Induced antioxidant and immune response	[204]
	Lipoteichoic Acid	Zebrafish	↑immune system response, antibacterial effects, higher expression of TLR1 and TLR 2	[189]
	Levan	Pacific white shrimps	Good immune modulator and preventive factor against vibriosis	[205]
		Zebrafish	Modification changes of gut microbiota but not affected SCFAs	[206]
		Silver pomfret	↑TLR, RPA	[137]
	Lentinan	Labeo rohita	↑Lys, TP, A: G, RBA, RPA	[140]
		common carp	Potent nutraceutical and immuno additive agents	[207]
		<i>Cyprinus carpio</i>	Growth promoter and antibacterial agents	[208]
		juvenile taimen	↑TGF- β , TNF- α , IL-1 β , IL-6 and IL-8, ↑SOD, CAT, and GSH-Px, RPA	[148,149]
		Senegalese sole	Modulate immune system and regulate gut microbiota	[209]

↑- Enhance or Increase or Improve.

2.28. Laminarin

Laminarin is the most significant component derived from plants and seaweeds; Laminarin has a lot of bio-functional performances such as antitumor, anti-apoptotic, anti-inflammatory, anticoagulant, and antioxidant activity. The biological activities of laminarin can be improved after appropriate chemical modifications, sulphation, and novel processing techniques. Studies on feeding laminarin-rich extracts to animals point out its suitability as an efficient component for food applications [177,178]. Laminarin is generally low molecular weight (e.g., 5 kDa) and their inter-chain linkages with glucans residues are of great importance [179]. [180] Cui et al. (2021) demonstrated that laminarin injection can change the expression of Arginine kinase (AK), and antioxidant activity, and reduced the activity of *Staphylococcus aureus*, *Listeria monocytogenes*, *Escherichia coli*, and *Salmonella typhimurium*. Similarly [181], Shanmugavel and Krishnamoorthy (2021) reported that laminarin may act as an immunostimulant and the energy source of crustaceans [179]. Meng et al. (2020) demonstrated that dietary laminarin of 1 g/kg enhanced growth activity, feed efficiency, and the nonspecific immune reaction of *E. coioides* and also The levels of Creatinine (CREA) and UREA, as well as the action of alkaline phosphatase (ALP), were lower than of the control. There was no major difference in the levels of alanine aminotransferase (ALT) and, aspartate aminotransferase (AST) between control groups and treated groups. This study recommended that laminarin regulates the immune response and induces growth of the fish.

2.29. Ulvans

Ulvans are derived from the cell wall of green seaweeds; they contain sulfated polysaccharides, *Ulva* was elevated in dietary fiber which stimulates gastrointestinal health and is linked to a reduction in the incidence of chronic diseases as well as biological activities of dietary ulvans contains several positive responses such as immunomodulating, antiviral, antioxidant, antihyperlipidemic and anticancer activity in human and animals (including fish) [181,183]. In literature, it was reported for the first time that dietary supplementation (1 g/kg) of ulvans component could enhance immune system response and disease resistance against bacterial and parasitic infection and there was no significant difference in growth performance among fish for any of the diets tested of Nile tilapia (*Oreochromis niloticus*) [184]. [185] Ponce et al. (2020) suggested that *U. ohnoi* containing ulvans element was nontoxic and potential application as a vaccine adjuvant and/or nutraceuticals in aquaculture. Ulvans (0.5 mg/fish) was used to enhance immune gene

expression and regulate immune pathways in flatfish (*Senegalese sole*) juveniles. Dietary supplements of ulvans as sulfated polysaccharide elements contain numerous health-beneficial effects such as anticoagulant, antiviral, antioxidative, anticancer, and immunomodulating activities. In addition, dietary additives of ulvans help improve aquatic animal health and there is no negative impact on aquatic animals [186]. Dietary supplementation range of ulvans (0.05%) improved immunological parameters (respiratory burst, total hemocyte count, and phenoloxidase activities) of shrimps and also survival and challenging tests in giant tiger prawn (*P. monodon*) and Pacific white shrimp (*L. vannamei*). The dietary administration of ulvan was an effective immunostimulant for *L. vannamei* and *P. monodon* [187]. The function of other original bioactive compounds as immunostimulants in aquatic animals has been summarised in Table 4.

3. Conclusion

This review is based on the last few years' literature data analysis and provides an enlightening direction for bioactive immunostimulant improvement in the aquaculture industry. As a comprehensive study was needed to improve bioactive immunostimulants, this review concluded that bioactive immunostimulants sources, types, functions, and administration methods are beneficial for the improvement of healthy aquatic production and also for the increase of feed quality levels. Due to this review report, information was presented about bioactive immunostimulants, which are beneficial with a decreased negative impact on aquatic animal health. Thus, such a report mainly focused on positive responses analysis. The utilization of plant, animal, bacterial, fungal, and algal-originated bioactive immunostimulants is strongly recommended for improved aquatic animal health and aquatic environment enhancement. Biomolecules' functional feed additives improve healthy aquatic productions and enhanced aquatic feed quality levels and are also less toxic, cost-effective, and environmentally friendly. The next steps of our research group will focus on the extraction, characterization, and identification of biomolecules from different natural sources, biomolecules interaction analysis from the body, identification of novel bioactive immunostimulants, able to improve functional nutritional supplements in the modern aquaculture industry. This study information is enlightening to the future research direction of novel bioactive immunostimulants in the aquaculture industry.

Declaration of competing interest

Authors have no conflict of interest to declare for the publication of

Table 4
Various bioactive compounds as immunostimulants in aquaculture.

Origin	Bioactive immunostimulants	Aquatic organisms	Immune response	References
Algae	Alginate	<i>Penaeus monodon</i> Rainbow trout	↑ immune system, growth and antiviral activity	[169,172]
Yeast	β-Glucans	<i>Litopenaeus vannamei</i> Red sea bream Atlantic Salmon	↑phagocytic activity, RPA, resistance against bacterial pathogens	[164,165]
	Mannanoligosaccharides	Rainbow trout Nile tilapia	↑TLC ↑ blood protein, RPA	[154,156]
Seaweed	Alginate	Tilapia Grouper <i>E. Coioides</i>	PA, PoA, RBA	[166,167]
	Fucoidan	<i>P. monodon</i>	↑ immune system, RPA	[176]
	Laminarin	<i>E. coioides</i>	↑ non specific immunity, RPA	[179]
	Ulvans	<i>P. monodon</i> <i>L. vannamei</i> Nile tilapia	THC, RBA, PoA ↑immune response, RPA	[184,187]

↑- Enhance or Increase or Improve.

Abbreviations of immunological parameters: ACP- Acid phosphatase; RPA-reduce pathogenic activity; IgM-ImmunoglobulinM; TLC-Total leukocyte count, THC-Total haemocyte count; SOD-Superoxide dismutase; GSH-Px, Glutathione-peroxidase; CAT-Catalase; TAC - Total antioxidant capacity; Lys- Lysozyme activity; PA-Phagocytic activity; PI- Phagocytic activity PoA-phenoloxidase activity; TP-Total protein; A:G, Albumin/Globulin; RBA- Respiratory burst activity; SOD- Super oxide dismutase; RBC- Red blood cell; WBC - White blood cell; IL-1a; Interleukin 1a; TNF-a, Tumor Necrosis Factor a; TAS-Total antioxidant status; ACP- the alternative complement pathway; PO activities-total peroxidase enzyme concentration; ACPA-the alternative complement pathway activity.

the present work.

Data availability

Data will be made available on request.

Acknowledgments

We wish to acknowledge the national natural science foundation of China (Grant no. 32072990), Xiamen Marine and Fisheries Development Fund (Grant No. 19CZP018HJ04), Industry-University Cooperation Project of Fujian Province (Grant No. 2018N5011) for supporting this research work. This research work was partially supported by Chiang Mai University, Chiang Mai, Thailand.

References

- [1] C.A. Wang, J. Li, L. Wang, Z. Zhao, L. Luo, X. Du, Q. Xu, Effects of dietary phosphorus on growth, body composition and immunity of young taimen *Hucho taimen* (Pallas, 1773), *Aquacult. Res.* 48 (6) (2017) 3066–3079, <https://doi.org/10.1111/are.13138>.
- [2] C. Ragunath, V. Ramasubramanian, Dietary effect of padina boergeseni on growth, immune response, and disease resistance against *Pseudomonas aeruginosa* in *Cirrhinus mrigala*, *Appl. Biochem. Biotechnol.* 194 (5) (2022) 1881–1897, <https://link.springer.com/article/10.1007/s12010-021-03770-y>.
- [3] J.P. Firmino, J. Galindo-Villegas, F.E. Reyes-López, E. Gisbert, Phytochemical bioactive compounds shape fish mucosal immunity, *Front. Immunol.* 12 (2021), 695973, <https://doi.org/10.3389/fimmu.2021.695973>.
- [4] K. Yang, X. Qi, M. He, K. Song, F. Luo, X. Qu, F. Ling, Dietary supplementation of salidroside increases immune response and disease resistance of crucian carp (*Carassius auratus*) against *Aeromonas hydrophila*, *Fish Shellfish Immunol.* 106 (2020) 1–7, <https://doi.org/10.1016/j.fsi.2020.07.054>.
- [5] T. Magrone, S. Fontana, F. Laforgia, T. Dragone, E. Jirillo, L. Passantino, Administration of a polyphenol-enriched feed to farmed sea bass (*Dicentrarchus labrax* L.) modulates intestinal and spleen immune responses, *Oxid. Med. Cell. Longev.* (2016), <https://doi.org/10.1155/2016/2827567>, 2016.
- [6] T. Magrone, M.A. Russo, E. Jirillo, Dietary approaches to attain fish health with special reference to their immune system, *Curr. Pharmaceut. Des.* 24 (41) (2018) 4921–4931, <https://doi.org/10.2174/1381612825666190104121544>.
- [7] C. Zhou, H. Lin, X. Ge, J. Niu, J. Wang, Y. Wang, X. Tan, The effects of dietary soybean isoflavones on growth, innate immune responses, hepatic antioxidant abilities and disease resistance of juvenile golden pompano *Trachinotus ovatus*, *Fish Shellfish Immunol.* 43 (1) (2015) 158–166, <https://doi.org/10.1016/j.fsi.2014.12.014>.
- [8] G. Ren, K. Li, Y. Hu, M. Yu, J. Qu, X. Xu, Optimization of selenizing conditions for Seleno-Lentinan and its characteristics, *Int. J. Biol. Macromol.* 81 (2015) 249–258, <https://doi.org/10.1016/j.ijbiomac.2015.08.012>.
- [9] N. Sadat Hoseini Madani, T.J. Adorian, H. Ghafari Farsani, S.H. Hoseinifar, The effects of dietary probiotic Bacilli (*Bacillus subtilis* and *Bacillus licheniformis*) on growth performance, feed efficiency, body composition and immune parameters of whiteleg shrimp (*Litopenaeus vannamei*) postlarvae, *Aquacult. Res.* 49 (5) (2018) 1926–1933, <https://doi.org/10.1111/are.13648>.
- [10] S.H. Hoseinifar, Y.Z. Sun, Z. Zhou, H. Van Doan, S.J. Davies, R. Harikrishnan, Boosting immune function and disease bio-control through environment-friendly and sustainable approaches in finfish aquaculture: herbal therapy scenarios, *Rev. Fish. Sci. Aquacul.* 28 (3) (2020) 303–321, <https://doi.org/10.1080/23308249.2020.1731420>.
- [11] O. Alfred, A. Shaahu, D.A. Orban, M. Egwenomhe, An overview on understanding the basic concept of fish diseases in aquaculture, *IRE J.* 4 (6) (2020), <http://www.irejournals.com/formatedpaper/1702541.pdf>.
- [12] J.A. Gephart, C.D. Golden, F. Asche, B. Belton, C. Brugere, H.E. Froehlich, E. H. Allison, Scenarios for global aquaculture and its role in human nutrition, *Rev. Fish. Sci. Aquacul.* 29 (1) (2020) 122–138, <https://doi.org/10.1080/23308249.2020.1782342>.
- [13] R.N. Roy, Bioactive natural derivatives of phthalate ester, *Crit. Rev. Biotechnol.* 40 (7) (2020) 913–929, <https://doi.org/10.1080/07388551.2020.1789838>.
- [14] G. Wu, Nutrition and metabolism: foundations for animal growth, development, reproduction, and health, in: *Recent Advances in Animal Nutrition and Metabolism*, Springer, Cham, 2022, pp. 1–24, https://doi.org/10.1007/978-3-030-85686-1_1.
- [15] V. Šimat, N. Elabed, P. Kulawik, Z. Ceylan, E. Jamroz, H. Yazgan, F. Özogul, Recent advances in marine-based nutraceuticals and their health benefits, *Mar. Drugs* 18 (12) (2020) 627, <https://doi.org/10.3390/md18120627>.
- [16] S.A. Ashraf, M. Adnan, M. Patel, A.J. Siddiqui, M. Sachidanandan, M. Snoussi, S. Hadi, Fish-based bioactives as potent nutraceuticals: exploring the therapeutic perspective of sustainable food from the sea, *Mar. Drugs* 18 (5) (2020) 265, <https://doi.org/10.3390/md18050265>.
- [17] P. Encarnação, Functional feed additives in aquaculture feeds, in: *Aquafeed Formulation*, Academic Press, 2016, pp. 217–237, <https://doi.org/10.1016/B978-0-12-800873-7.00005-1>.
- [18] M. Khodadadi, N. Abbasi, T.J. Adorian, H.G. Farsani, A. Hedayat, S.M. Hoseini, Growth performance, survival, body composition, hematological parameters, intestinal histomorphology, and digestive enzymes' activity in juvenile rainbow trout (*Oncorhynchus mykiss*) fed dietary Immunogen, *J. Appl. Aquacult.* 30 (2) (2018) 174–186, <https://doi.org/10.1080/10454438.2017.1420515>.
- [19] R. Zhang, X.W. Wang, L.L. Liu, Y.C. Cao, H. Zhu, Dietary oregano essential oil improved the immune response, activity of digestive enzymes, and intestinal microbiota of the koi carp, *Cyprinus carpio*, *Aquaculture* 518 (2020), 734781, <https://doi.org/10.1016/j.aquaculture.2019.734781>.
- [20] M.E. Büyükdöveci, J.L. Balcázar, I. Demirkale, S. Dikel, Effects of garlic-supplemented diet on growth performance and intestinal microbiota of rainbow trout (*Oncorhynchus mykiss*), *Aquaculture* 486 (2018) 170–174, <https://doi.org/10.1016/j.aquaculture.2017.12.022>.
- [21] F.J. Sutili, D.M. Gatlin III, B.M. Heinzmann, B. Baldissarotto, Plant essential oils as fish diet additives: benefits on fish health and stability in feed, *Rev. Aquacult.* 10 (3) (2018) 716–726, <https://doi.org/10.1111/raq.12197>.
- [22] E. Ringo, Z. Zhou, J.G. Vecino, S. Wadsworth, J. Romero, Å. Krogdahl, D. L. Merrifield, Effect of dietary components on the gut microbiota of aquatic animals. A never-ending story? *Aquacult. Nutr.* 22 (2) (2016) 219–282, <https://doi.org/10.1111/anu.12346>.
- [23] X. Wu, T. Teame, Q. Hao, Q. Ding, H. Liu, C. Ran, Z. Zhang, Use of a paraprobiotic and postbiotic feed supplement (HWF™) improves the growth performance, composition and function of gut microbiota in hybrid sturgeon (*Acipenser baeri* x *Acipenser schrenckii*), *Fish Shellfish Immunol.* 104 (2020) 36–45, <https://doi.org/10.1016/j.fsi.2020.05.054>.
- [24] S.H. Hoseinifar, A. Ahmadi, M. Raeisi, S.M. Hoseini, M. Khalili, N. Behnampour, Comparative study on immunomodulatory and growth enhancing effects of three prebiotics (galactooligosaccharide, fructooligosaccharide and inulin) in common carp (*Cyprinus carpio*), *Aquacult. Res.* 48 (7) (2017) 3298–3307, <https://doi.org/10.1111/are.13156>.
- [25] S.H. Hoseinifar, M. Dadar, E. Ringo, Modulation of nutrient digestibility and digestive enzyme activities in aquatic animals: the functional feed additives scenario, *Aquacult. Res.* 48 (8) (2017) 3987–4000, <https://doi.org/10.1111/are.13368>.
- [26] S.H. Hoseinifar, A. Mirvaghefi, M.A. Amoozegar, D.L. Merrifield, E. Ringo, In vitro selection of a synbiotic and in vivo evaluation on intestinal microbiota, performance and physiological response of rainbow trout (*Oncorhynchus mykiss*) fingerlings, *Aquacult. Nutr.* 23 (1) (2017) 111–118, <https://doi.org/10.1111/anu.12373>.
- [27] S.H. Hoseinifar, R. Safari, M. Dadar, Dietary sodium propionate affects mucosal immune parameters, growth and appetite related genes expression: insights from zebrafish model, *Gen. Comp. Endocrinol.* 243 (2017) 78–83, <https://doi.org/10.1016/j.ygcen.2016.11.008>.
- [28] A. Rodríguez, J. Velázquez, L. González, T. Rodríguez-Ramos, B. Dixon, F. H. Miyares, Y. Carpio, PACAP modulates the transcription of TLR-1/TLR-5/MyD88 pathway genes and boosts antimicrobial defenses in *Clarias gariepinus*, *Fish Shellfish Immunol.* 115 (2021) 150–159, <https://doi.org/10.1016/j.fsi.2021.06.009>.
- [29] S.H. Hoseinifar, G. Rashidian, H. Ghafari Farsani, M.A. Jahazi, M. Soltani, H. V. Doan, M. Paolucci, Effects of apple (*Malus pomila*) pomace-derived pectin on the innate immune responses, expressions of key immune-related genes, growth performance, and digestive enzyme activity of rainbow trout (*Oncorhynchus mykiss*), *Animals* 11 (7) (2021) 2117, <https://www.mdpi.com/2076-2615/11/7/2117>.
- [30] U.S. Sagaram, M.S. Gaikwad, R. Nandru, S. Dasgupta, Microalgae as feed ingredients: recent developments on their role in immunomodulation and gut microbiota of aquaculture species, *FEMS (Fed. Eur. Microbiol. Soc.) Microbiol. Lett.* 368 (11) (2021), <https://doi.org/10.1093/femsle/fnab071>.
- [31] M.F.Z. Ali, S. Nakahara, Y. Otsu, A. Ido, C. Miura, T. Miura, Effects of functional polysaccharide from silkworm as an immunostimulant on transcriptional profiling and disease resistance in fish, *J. Insects as Food and Feed* (2021) 1–14, <https://doi.org/10.3920/JIFF2021.0108>.
- [32] V.K. Gurjar, D. Pal, Natural compounds extracted from medicinal plants and their immunomodulatory activities, in: *Bioactive Natural Products for Pharmaceutical Applications*, Springer, Cham, 2021, pp. 197–261, https://doi.org/10.1007/978-3-030-54027-2_6.
- [33] M.G. Mendoza Rodríguez, C. Pohlenz, D.M. Gatlin III, Supplementation of organic acids and algae extracts in the diet of red drum *Sciaenops ocellatus*: immunological impacts, *Aquacult. Res.* 48 (4) (2017) 1778–1786, <https://doi.org/10.1111/are.13015>.
- [34] S. Luna-Castro, I. Ceballos-Olvera, F. Benavides-González, Z. Blanco-Martínez, G. Sánchez-Martínez, M.D.L.L. Vázquez-Sauceda, M. de la Garza, Bovine lactoferrin in fish culture: current research and future directions, *Aquacult. Res.* 53 (3) (2022) 735–745, <https://doi.org/10.1111/are.15621>.
- [35] H. Minářová, L. Bláhová, J. Kalina, I. Papežíková, J. Mareš, R. Kopp, M. Palíková, Plant-based and immunostimulant-enhanced diets modulate oxidative stress, immune and haematological indices in rainbow trout (*Oncorhynchus mykiss*), *Acta Vet.* 90 (2) (2021) 233–253, <https://actavet.vfu.cz/90/2/233/>.
- [36] C. Lumsangkul, W. Tapingkae, K. Sringarm, S. Jaturasitha, C. Le Xuan, S. Wannavijit, H. Van Doan, Effect of dietary sugarcane bagasse supplementation on growth performance, immune response, and immune and antioxidant-related gene expressions of Nile tilapia (*Oreochromis niloticus*) cultured under biofloc system, *Animals* 11 (7) (2021) 2035, <https://doi.org/10.3390/ani11072035>.
- [37] P. Yarahmadi, H.G. Farsani, A. Khazaei, M. Khodadadi, G. Rashidian, M. A. Jalali, Protective effects of the prebiotic on the immunological indicators of rainbow trout (*Oncorhynchus mykiss*) infected with *Aeromonas hydrophila*, *Fish*

- Shellfish Immunol. 54 (2016) 589–597, <https://doi.org/10.1016/j.fsi.2016.05.010>.
- [38] R.B. Dixit, U.S. Sagaram, C. Gocher, G.R. Krishna Kumar, S. Dasgupta, Biomolecular characterisation of marine microalga in comparison to fishmeal and soymeal as an alternative feed ingredient, *Phytochem. Anal.* 33 (3) (2022) 365–372, <https://doi.org/10.1002/pca.3094>.
- [39] M. Abha, S. Abhisweta, Effect of curcumin, ascorbic acid against chlorpyrifos toxicity on histotexture and bimolecular content of fresh water catfish, *Heteropneustes fossilis*, *J. Exp. Zool. India* 23 (2) (2020) 1043–1049, <https://www.cabdirect.org/cabdirect/abstract/20203390895>.
- [40] F. Baleta, P. Candelaria, D.E. Baldo, L.J. Baleta, L. Plantado, M. Navarro, Dietary incorporation of Sweet Potato Ipomoea batatas shoots improved growth performance and haematological profile of Tilapia *Oreochromis niloticus* in Hapa Nets, *Aquat. Res.* 5 (1) (2022) 1–10, <https://doi.org/10.3153/AR22001>.
- [41] M.T. El-Saadony, N.M. Zaberemawi, N.M. Zaberemawi, M.A. Burollus, M.E. Shafi, M. Alagawany, M.E. Abd El-Hack, Nutritional aspects and health benefits of bioactive plant compounds against infectious diseases: a review, *Food Rev. Int.* 1–23 (2021), <https://doi.org/10.1080/87559129.2021.1944183>.
- [42] M. Paolucci, M.G. Volpe, S. Costantini, Polyphenol enriched diets for aquaculture applications, *J. Int. Soc. Antioxid. Nutr. Health* 3 (4) (2016).
- [43] M.A. Dawood, S. Koshio, M.A. Esteban, Beneficial roles of feed additives as immunostimulants in aquaculture: a review, *Rev. Aquacult.* 10 (4) (2018) 950–974, <https://doi.org/10.1111/raq.12209>.
- [44] M. Arieli, D. Fiocco, S. Fontana, M.P. Arena, M.A. Frassanito, A. Gallone, Administration of a polyphenol-enriched feed to farmed sea bass (*Dicentrarchus labrax* L.): kidney melanomacrophages response, *Fish Shellfish Immunol.* 68 (2017) 404–410, <https://doi.org/10.1016/j.fsi.2017.07.043>.
- [45] M. Long, W. Lin, J. Hou, H. Guo, L. Li, D. Li, F. Yang, Dietary supplementation with selenium yeast and tea polyphenols improve growth performance and nitrite tolerance of Wuchang bream (*Megalobrama amblycephala*), *Fish Shellfish Immunol.* 68 (2017) 74–83, <https://doi.org/10.1016/j.fsi.2017.07.017>.
- [46] M.G. Volpe, S. Costantini, E. Coccia, L. Parrillo, M. Paolucci, Evaluation of metabolic changes induced by polyphenols in the crayfish *Astacus leptodactylus* by metabolomics using Fourier transformed infrared spectroscopy, *J. Biosci.* 43 (4) (2018) 585–596, <https://doi.org/10.1007/s12038-018-9774-1>.
- [47] H. Van Doan, S.H. Hoseinifar, K. Sringarm, S. Jaturasitha, B. Yuangsoi, M. A. Dawood, C. Faggio, Effects of Assam tea extract on growth, skin mucus, serum immunity and disease resistance of Nile tilapia (*Oreochromis niloticus*) against *Streptococcus agalactiae*, *Fish Shellfish Immunol.* 93 (2019) 428–435, <https://doi.org/10.1016/j.fsi.2019.07.077>.
- [48] T. Magrone, A. Spagnoletta, M. Magrone, M.A. Russo, A. Corriero, E. Jirillo, L. Passantino, Effects of polyphenol administration to European farmed sea bass (*Dicentrarchus labrax* L.): special focus on hepatopancreas morphology, *Endocr. Metab. Immune Disord. - Drug Targets* 19 (4) (2019) 526–533, <https://doi.org/10.2174/187153031866618100911214>.
- [49] L. Zhong, Y. Hu, Y. Hu, J. Li, Y. Tian, J. Chen, T. Xiao, Effects of dietary tea polyphenols on growth, immunity and lipid metabolism of juvenile black carp *Mylopharyngodon piceus*, *Aquacult. Res.* 51 (2) (2020) 569–576, <https://doi.org/10.1111/are.14403>.
- [50] R. Safari, S.H. Hoseinifar, M.R. Iranpour, M. Mazandarani, H. Sanchouli, M. Paolucci, Effects of dietary polyphenols on mucosal and humoral immune responses, antioxidant defense and growth gene expression in beluga sturgeon (*Huso huso*), *Aquaculture* 528 (2020), 735494, <https://doi.org/10.1016/j.aquaculture.2020.735494>.
- [51] M.A. Jahazi, S.H. Hoseinifar, V. Jafari, A. Hajimoradloo, H. Van Doan, M. Paolucci, Dietary supplementation of polyphenols positively affects the innate immune response, oxidative status, and growth performance of common carp, *Cyprinus carpio* L., *Aquaculture* 517 (2020), 734709, <https://doi.org/10.1016/j.aquaculture.2019.734709>.
- [52] S.M. Hussain, H. Gohar, M. Asrar, M.M. Shahzad, A. Rasul, M. Hussain, A. Sharif, Effects of Polyphenols Supplemented Canola Meal Based Diet on Proximate Composition, Minerals Absorption and Hematology of *Cyprinus carpio* Fingerlings, 2021. https://researcherslinks.com/uploads/articles/1620236631/PJZ_MH20190214170258-R3_Hussain%20et%20al.pdf.
- [53] U. Sarker, S. Oba, Polyphenol and flavonoid profiles and radical scavenging activity in leafy vegetable *Amaranthus gangeticus*, *BMC Plant Biol.* 20 (1) (2020) 1–12, <https://doi.org/10.1186/s12870-020-02700-0>.
- [54] M. Li, X. Zhu, J. Tian, M. Liu, G. Wang, Dietary flavonoids from Allium mongolicum Regel promotes growth, improves immune, antioxidant status, immune-related signaling molecules and disease resistance in juvenile northern snakehead fish (*Channa argus*), *Aquaculture* 501 (2019) 473–481, <https://doi.org/10.1016/j.aquaculture.2018.12.011>.
- [55] Z.G. Sun, Z.N. Li, X.W. Miao, S. Li, H.L. Zhu, Recent advances in natural products with antiviral activities, *Mini Rev. Med. Chem.* 21 (14) (2021) 1888–1908, <https://doi.org/10.2174/138955752166621030410824>.
- [56] C.G. Awuchi, The biochemistry, toxicology, and uses of the pharmacologically active phytochemicals: alkaloids, terpenes, polyphenols, and glycosides, *J. Food Pharm. Sci.* (2019) 2. -2, [file:///C:/Users/Hii/AppData/Local/Temp/666-Article%20Text-2613-4-10-20210723.pdf](https://doi.org/10.1016/j.fsi.2019.07.077).
- [57] S. Andayani, M. Fadjar, A.P. Rahayu, M.F. Rahman, E. Kurniati, Potency of jellyfish alkaloid (*Aeginura* sp) on kidney histopathology and relative percent survival (RPS) of tiger grouper (*Epinephelus fuscoguttatus*) against *Vibrio harveyi*, No. 1, in: IOP Conference Series: Earth and Environmental Science, vol. 584, 2020, October, 12065. IOP Publishing, <https://iopscience.iop.org/article/10.1088/1755-1315/584/1/012065/meta>.
- [58] Q. Ye, Y. Feng, Z. Wang, A. Zhou, S. Xie, L. Fan, J. Zou, Effects of dietary Gelsemium elegans alkaloids on intestinal morphology, antioxidant status, immune responses and microbiota of *Megalobrama amblycephala*, *Fish Shellfish Immunol.* 94 (2019) 464–478, <https://doi.org/10.1016/j.fsi.2019.09.048>.
- [59] X. Zhou, Y. Peng, L. Li, K. He, T. Huang, S. Mou, X. Li, Effects of dietary supplementations with the fibrous root of *Rhizoma Coptidis* and its main alkaloids on non-specific immunity and disease resistance of common carp, *Vet. Immunol. Immunopathol.* 173 (2016) 34–38, <https://doi.org/10.1016/j.vetimm.2016.03.014>.
- [60] M. Kherade, S. Solanke, M. Tawar, S. Wankhede, Fructooligosaccharides: a comprehensive review, *J. Ayu. Herb. Med* 7 (2021) 193–200, <https://doi.org/10.31254/jahm.2021.7305>.
- [61] A. Nawaz, S. Irshad, S.H. Hoseinifar, H. Xiong, The functionality of prebiotics as immunostimulant: evidences from trials on terrestrial and aquatic animals, *Fish Shellfish Immunol.* 76 (2018) 272–278, <https://doi.org/10.1016/j.fsi.2018.03.004>.
- [62] D.B. Bebo, J. Zhu, S.M. Sun, C.Y. Song, Effects of Antibiotics on the Growth, Antioxidant Capacity, Activity of Digestive Enzymes and Serum Biochemical Indices, Liver and Intestine Morphology of *Megalobrama amblycephala* Fry, 2020. <https://www.fisheriesjournal.com/archives/2020/vol8issue3/PartG/8-3-39-522.pdf>.
- [63] H. Ghafarifarani, S.H. Hoseinifar, M. Aftabgard, H. Van Doan, The improving role of savory (*Satureja hortensis*) essential oil for Caspian roach (*Rutilus rutilus*) fry: growth, haematological, immunological, and antioxidant parameters and resistance to salinity stress, *Aquaculture* 548 (2022), 737653, <https://doi.org/10.1016/j.aquaculture.2021.737653>.
- [64] M.T. Hasan, W. Je Jang, J.M. Lee, B.J. Lee, S.W. Hur, I.S. Kong, Effects of immunostimulants, prebiotics, probiotics, synbiotics, and potentially immunoreactive feed additives on olive flounder (*Paralichthys olivaceus*): a review, *Rev. Fish. Sci. Aquacul.* 27 (4) (2019) 417–437, <https://doi.org/10.1080/23308249.2019.1622510>.
- [65] L. Poolsawat, X. Li, H. Yang, P. Yang, M.A. Kabir Chowdhury, A. Yusuf, X. Leng, The potentials of fructooligosaccharide on growth, feed utilization, immune and antioxidant parameters, microbial community and disease resistance of tilapia (*Oreochromis niloticus* × *O. aureus*), *Aquacult. Res.* 51 (11) (2020) 4430–4442, <https://doi.org/10.1111/are.14786>.
- [66] A. Raza, S. Iqbal, Z. Ahmad, M.A. Rehman, M. Waseem, M. Usman, Conversion of milk lactose to galacto-oligosaccharides by enzymes to produce prebiotic enriched cheese, *Future Foods* 4 (2021), 100097, <https://doi.org/10.1016/j.fufo.2021.100097>.
- [67] A. Rashidinejad, A. Bahrami, A. Rehman, A. Rezaei, A. Babazadeh, H. Singh, S. M. Jafari, Co-encapsulation of probiotics with prebiotics and their application in functional/synbiotic dairy products, *Crit. Rev. Food Sci. Nutr.* 62 (9) (2022) 2470–2494, <https://www.tandfonline.com/doi/abs/10.1080/10408398.2020.1854169>.
- [68] S.H. Hoseinifar, F. Zoheiri, M. Dadar, R. Rufchaei, E. Ringø, Dietary galactooligosaccharide elicits positive effects on non-specific immune parameters and growth performance in Caspian white fish (*Rutilus frisii kutum*) fry, *Fish Shellfish Immunol.* 56 (2016) 467–472, <https://doi.org/10.1016/j.fsi.2016.08.001>.
- [69] M.T. Hasan, W.J. Jang, H. Kim, B.J. Lee, K.W. Kim, S.W. Hur, I.S. Kong, Synergistic effects of dietary *Bacillus* sp. SJ-10 plus β-galactooligosaccharides as a synbiotic on growth performance, innate immunity and streptococcosis resistance in olive flounder (*Paralichthys olivaceus*), *Fish Shellfish Immunol.* 82 (2018) 544–553, <https://doi.org/10.1016/j.aquaculture.2017.09.020>.
- [70] K. Yukgehnaish, P. Kumar, P. Sivachandran, K. Marimuthu, A. Arshad, B.A. Paray, J. Arockiaraj, Gut microbiota metagenomics in aquaculture: factors influencing gut microbiome and its physiological role in fish, *Rev. Aquacult.* 12 (3) (2020) 1903–1927, <https://doi.org/10.1111/raq.12416>.
- [71] R. Tarkhani, A. Imani, S.H. Hoseinifar, O. Ashayerizadeh, K.S. Moghanlou, R. Manaffar, M. Reverter, Comparative study of host-associated and commercial probiotic effects on serum and mucosal immune parameters, intestinal microbiota, digestive enzymes activity and growth performance of roach (*Rutilus rutilus*) fingerlings, *Fish Shellfish Immunol.* 98 (2020) 661–669, <https://doi.org/10.1016/j.fsi.2019.10.063>.
- [72] S.H. Hoseinifar, A. Mirvaghefi, M.A. Amoozegar, M. Sharifian, M.A. Esteban, Modulation of innate immune response, mucosal parameters and disease resistance in rainbow trout (*Oncorhynchus mykiss*) upon synbiotic feeding, *Fish Shellfish Immunol.* 45 (1) (2015) 27–32, <https://doi.org/10.1016/j.fsi.2015.03.029>.
- [73] M. Modanloo, S. Soltanian, M. Akhlaghi, S.H. Hoseinifar, The effects of single or combined administration of galactooligosaccharide and *Pediococcus acidilactici* on cutaneous mucus immune parameters, humoral immune responses and immune related genes expression in common carp (*Cyprinus carpio*) fingerlings, *Fish Shellfish Immunol.* 70 (2017) 391–397, <https://doi.org/10.1016/j.fsi.2017.09.032>.
- [74] S.H. Hoseinifar, F. Zoheiri, C.M. Caipang, Dietary sodium propionate improved performance, mucosal and humoral immune responses in Caspian white fish (*Rutilus frisii kutum*) fry, *Fish Shellfish Immunol.* 55 (2016) 523–528, <https://doi.org/10.1016/j.fsi.2016.06.027>.
- [75] O. Safari, M. Paolucci, Modulation of growth performance, immunity, and disease resistance in narrow-clawed crayfish, *Astacus leptodactylus* (*leptodactylus* (Eschscholtz, 1823)) upon synbiotic feeding, *Aquaculture* 479 (2017) 333–341, <https://doi.org/10.1016/j.aquaculture.2017.05.049>.
- [76] H. Ghafarifarani, G. Rashidian, T. Bagheri, S.H. Hoseinifar, H. Van Doan, Study on growth enhancement and the protective effects of dietary prebiotic inulin on

- immunity responses of rainbow trout fry infected with, *Ann. Anim. Sci.* 21 (2) (2021) 543–559, <https://doi.org/10.2478/aoas-2020-0074>.
- [77] S.K. Gupta, B. Sarkar, M. Priyam, N. Kumar, S. Naskar, M.J. Foysal, T.R. Sharma, Inflammatory and stress biomarker response of *Aeromonas hydrophila* infected rohu, *Labeo rohita* fingerlings to dietary microbial levan, *Aquaculture* 521 (2020), 735020, <https://doi.org/10.1016/j.aquaculture.2020.735020>.
- [78] W.Y. Mo, Z. Cheng, W.M. Choi, C.H. Lun, Y.B. Man, J.T. Wong, M.H. Wong, Use of food waste as fish feeds: effects of prebiotic fibers (inulin and mannanoligosaccharide) on growth and non-specific immunity of grass carp (*Ctenopharyngodon idella*), *Environ. Sci. Pollut. Control Ser.* 22 (22) (2015) 17663–17671, <https://link.springer.com/article/10.1007/s11356-015-4971-z>.
- [79] S. Syed Raffiq Ali, K. Ambasankar, P.E. Praveena, S. Nandakumar, M. Saiyad Musthafa, Effect of dietary prebiotic inulin on histology, immuno-haematological and biochemical parameters of Asian seabass (*Lates calcarifer*), *Aquacult. Res.* 49 (8) (2018) 2732–2740, <https://doi.org/10.1111/are.13734>.
- [80] X. Zhang, L. Xie, J. Long, Q. Xie, Y. Zheng, K. Liu, X. Li, Salidroside: a review of its recent advances in synthetic pathways and pharmacological properties, *Chem. Biol. Interact.* 339 (2021), 109268, <https://doi.org/10.1016/j.cbi.2020.109268>.
- [81] M. Zare, M.K. Mirzakhani, V. Stejskal, Growth, body proximate composition and selected blood parameters of rainbow trout *Oncorhynchus mykiss* fingerlings fed on a diet supplemented with nettle *Urtica dioica* and tarragon *Artemisia dracunculoides*, *Aquacult. Res.* 52 (11) (2021) 5691–5702, <https://doi.org/10.1111/are.15444>.
- [82] J. Xie, H. Fang, X. He, S. Liao, Y. Liu, L. Tian, J. Niu, Study on mechanism of synthetic astaxanthin and *Haematococcus pluvialis* improving the growth performance and antioxidant capacity under acute hypoxia stress of golden pompano (*Trachinotus ovatus*) and enhancing anti-inflammatory by activating Nrf2-ARE pathway to antagonize the NF- κ B pathway, *Aquaculture* 518 (2020), 734657, <https://doi.org/10.1016/j.aquaculture.2019.734657>.
- [83] L. Chen, M. Li, Z. Yang, W. Tao, P. Wang, X. Tian, W. Wang, Gardenia jasminoides Ellis: ethnopharmacology, phytochemistry, and pharmacological and industrial applications of an important traditional Chinese medicine, *J. Ethnopharmacol.* 257 (2020), 112829, <https://doi.org/10.1016/j.jep.2020.112829>.
- [84] W. Sun, X. Li, H. Xu, J. Chen, X. Xu, X. Leng, Effects of dietary geniposide on growth, flesh quality, and lipid metabolism of grass carp, *Ctenopharyngodon idella*, *J. World Aquacult. Soc.* 48 (6) (2017) 927–937, <https://doi.org/10.1111/jwas.12412>.
- [85] A.G. Huang, X. Tu, X.Z. Qi, F. Ling, B. Zhu, G.X. Wang, Gardenia jasminoides Ellis inhibit white spot syndrome virus replication in red swamp crayfish *Procambarus clarkii*, *Aquaculture* 504 (2019) 239–247, <https://doi.org/10.1016/j.aquaculture.2019.02.008>.
- [86] E. Wang, X. Chen, K. Wang, J. Wang, D. Chen, Y. Geng, X. Wei, Plant polysaccharides used as immunostimulants enhance innate immune response and disease resistance against *Aeromonas hydrophila* infection in fish, *Fish Shellfish Immunol.* 59 (2016) 196–202, <https://doi.org/10.1016/j.fsi.2016.10.039>.
- [87] M. He, G. Liu, Y. Liu, K. Yang, X. Qi, A. Huang, E. Wang, Effects of geniposide as immunostimulant on the innate immune response and disease resistance in crucian carp, *Aquaculture* 529 (2020), 735713, <https://doi.org/10.1016/j.aquaculture.2020.735713>.
- [88] M. Tsaneva, E.J. Van Damme, 130 years of plant lectin research, *Glycoconj. J.* 37 (5) (2020) 533–551, <https://doi.org/10.1007/s10719-020-09942-y>.
- [89] Y. Huang, F. Jiang, R. Wang, Y. Shi, S. Hu, T. Wu, Z. Zhao, In silico characterization and expression analysis of eight C-type lectins in obscure puffer *Takifugu obscurus*, *Vet. Immunol. Immunopathol.* 234 (2021), 110200, <https://doi.org/10.1016/j.vetimm.2021.110200>.
- [90] Y. Huang, Y. Shi, S. Hu, T. Wu, Z. Zhao, Characterization and functional analysis of two transmembrane C-type lectins in obscure puffer (*Takifugu obscurus*), *Front. Immunol.* 11 (2020) 436, <https://doi.org/10.3389/fimmu.2020.00436>.
- [91] D.L. Lindenwald, B. Lepenies, C-Type lectins in veterinary species: recent advancements and applications, *Int. J. Mol. Sci.* 21 (14) (2020) 5122, <https://doi.org/10.3390/ijms21145122>.
- [92] S. Narla, M. Azzam, S. Townsend, G. Vellaichamy, A.V. Marzano, A. Alavi, I. H. Hamzavi, Identifying key components and therapeutic targets of the immune system in hidradenitis suppurativa with an emphasis on neutrophils, *Br. J. Dermatol.* 184 (6) (2021) 1004–1013, <https://doi.org/10.1111/bjd.19538>.
- [93] V. Morales-Moya, F. Stambuk, R. Farlora, F. Guzmán, L. Mercado, K. Brokordt, P. Schmitt, Molecular characterization and cellular localization of a transmembrane C-type lectin receptor in hemocytes from the scallop *Argopecten purpuratus*, *Aquaculture* 546 (2022), 737293, <https://doi.org/10.1016/j.aquaculture.2021.737293>.
- [94] K. Zhang, X. Liu, X. Li, Y. Liu, H. Yu, J. Liu, Q. Zhang, Antibacterial functions of a novel fish-egg lectin from spotted knifejaw (*Oplegnathus punctatus*) during host defense immune responses, *Dev. Comp. Immunol.* 111 (2020), 103758, <https://doi.org/10.1016/j.dci.2020.103758>.
- [95] E. Catanzaro, C. Calcabrini, A. Bishayee, C. Fimognari, Antitumor potential of marine and freshwater lectins, *Mar. Drugs* 18 (1) (2019) 11, <https://doi.org/10.3390/md18010011>.
- [96] W. Liang, R. Wu, T. Yang, H. Shen, Z. Hu, Effect of pathogenic bacteria on a novel C-type lectin, hemocyte and superoxide dismutase/alkaline phosphatase activity in *Onchidium reevesii*, *Fish Shellfish Immunol.* 102 (2020) 185–194, <https://doi.org/10.1016/j.fsi.2020.04.001>.
- [97] Z. Meng, B. Zhang, B. Liu, H. Li, S. Fan, D. Yu, High carotenoids content can enhance resistance of selected Pinctada fucata families to high temperature stress, *Fish Shellfish Immunol.* 61 (2017) 211–218, <https://doi.org/10.1016/j.fsi.2016.12.032>.
- [98] R. Sowmya, N.M. Sachindra, Enhancement of non-specific immune responses in common carp, *Cyprinus carpio*, by dietary carotenoids obtained from shrimp exoskeleton, *Aquacult. Res.* 46 (7) (2015) 1562–1572, <https://doi.org/10.1111/are.12310>.
- [99] T. Nakano, G. Wiegertjes, Properties of carotenoids in fish fitness: a review, *Mar. Drugs* 18 (11) (2020) 568, <https://doi.org/10.3390/md18110568>.
- [100] S.K. Singh, M.P. Bhandari, P. Timalina, Effect of dietary supplementation of carotenoids on growth performance of rainbow trout (*Oncorhynchus mykiss*, Walbaum) in Rasuwa, Nepal, *Nepalese J. Agric. Sci.* 75, https://hcast.edu.np/wp-content/uploads/2021/08/NJAS_2021_V21_July.pdf#page=76.
- [101] J.M. Joy, A. Joseph, R. Anandan, The Role of Carotenoids in Enhancing the Health of Aquatic Organisms, 2021, <http://krishi.icar.gov.in/jspui/handle/123456789/60365>.
- [102] M.S. Hassan, E.Y. Mohammady, M.R. Soady, S.A. Sabae, A.M. Mahmoud, E. R. El-Haroun, Comparative study on the effect of dietary β -carotene and phycocyanin extracted from *Spirulina platensis* on immune-oxidative stress biomarkers, genes expression and intestinal enzymes, serum biochemical in Nile tilapia, *Oreochromis niloticus*, *Fish Shellfish Immunol.* 108 (2021) 63–72, <https://doi.org/10.1016/j.fsi.2020.11.012>.
- [103] A. Babin, C. Siciat, M. Teixeira, J.P. Troussard, S. Motreuil, J. Moreau, Y. Moret, Limiting immunopathology: interaction between carotenoids and enzymatic antioxidant defences, *Dev. Comp. Immunol.* 49 (2) (2015) 278–281, <https://doi.org/10.1016/j.dci.2014.12.007>.
- [104] X. Yuan, J. Zheng, S. Jiao, G. Cheng, C. Feng, Y. Du, H. Liu, A review on the preparation of chitosan oligosaccharides and application to human health, animal husbandry and agricultural production, *Carbohydr. Polym.* 220 (2019) 60–70, <https://doi.org/10.1016/j.carbpol.2019.05.050>.
- [105] D. Kamilya, M.I.R. Khan, Chitin and chitosan as promising immunostimulant for aquaculture, in: *Handbook of Chitin and Chitosan*, Elsevier, 2020, pp. 761–771, <https://doi.org/10.1016/B978-0-12-817966-6.00024-8>.
- [106] I.V. Zadine, L.D. dos Santos, H.J. Alves, A. de Marco Viott, L. de Souza Neves Ellendersen, G.I.B. de Muniz, R.A. Bombardelli, Chitosan foam-based filter: maintenance of water quality for Nile Tilapia cultivation, *Water, Air, Soil Pollut.* 231 (10) (2020) 1–20, <https://doi.org/10.1007/s11270-020-04905-3>.
- [107] W. Wang, C. Xue, X. Mao, Chitosan: structural modification, biological activity and application, *Int. J. Biol. Macromol.* 164 (2020) 4532–4546, <https://doi.org/10.1016/j.jbiomac.2020.09.042>.
- [108] S. Saini, A. Dhiman, S. Nanda, Immunomodulatory properties of chitosan: impact on wound healing and tissue repair, *Endocr. Metab. Immune Disord. - Drug Targets* 20 (10) (2020) 1611–1623, <https://doi.org/10.2174/1871530320666200503054605>.
- [109] D.M. Gatlin III, F.Y. Yamamoto, Nutritional supplements and fish health, in: *Fish Nutrition*, Academic Press, 2022, pp. 745–773, <https://doi.org/10.1016/B978-0-12-819587-1.00004-5>.
- [110] H.M. Abdel-Ghany, M.E.S. Salem, Effects of dietary chitosan supplementation on farmed fish: a review, *Rev. Aquacult.* 12 (1) (2020) 438–452, <https://doi.org/10.1111/raq.12326>.
- [111] A. Gupta, S.K. Gupta, M. Priyam, M.A. Siddik, N. Kumar, P.K. Mishra, A. Pattanayak, Immunomodulation by dietary supplements: a preventive health strategy for sustainable aquaculture of tropical freshwater fish, *Labeo rohita* (Hamilton, 1822), *Rev. Aquacult.* 13 (4) (2021) 2364–2394, <https://doi.org/10.1111/raq.12581>.
- [112] A.K. Oushani, M. Soltani, N. Sheikhzadeh, M.S. Mehrhan, H.R. Islami, Effects of dietary chitosan and nano-chitosan loaded cinnolitolite on growth and immune responses of rainbow trout (*Oncorhynchus mykiss*), *Fish Shellfish Immunol.* 98 (2020) 210–217, <https://doi.org/10.1016/j.fsi.2020.01.016>.
- [113] Y. Wu, A. Rashidpour, M.P. Almajano, I. Metón, Chitosan-based drug delivery system: applications in fish biotechnology, *Polymers* 12 (5) (2020) 1177, <https://doi.org/10.3390/polym12051177>.
- [114] H. Mondal, N. Chandrasekaran, A. Mukherjee, J. Thomas, Viral infections in cultured fish and shrimps: current status and treatment methods, *Aquacult. Int.* (2021) 1–36, <https://link.springer.com/article/10.1007/s10499-021-00795-2>.
- [115] N.M. Abu-Elala, S.H. Mohamed, M.M. Zaki, A.E. Eissa, Assessment of the immunomodulatory and antimicrobial effects of dietary chitosan on Nile tilapia (*Oreochromis niloticus*) with special emphasis to its bio-remediating impacts, *Fish Shellfish Immunol.* 46 (2) (2015) 678–685, <https://doi.org/10.1016/j.fsi.2015.08.004>.
- [116] B. Wang, Y.P. Timilsena, E. Blanch, B. Adhikari, Lactoferrin: structure, function, denaturation and digestion, *Crit. Rev. Food Sci. Nutr.* 59 (4) (2019) 580–596, <https://doi.org/10.1080/10408398.2017.1381583>.
- [117] P. Elumalai, A. Kurian, S. Lakshmi, C. Faggio, M.A. Esteban, E. Ringo, Herbal immunomodulators in aquaculture, *Rev. Fish. Sci. Aquacult.* 29 (1) (2020) 33–57, <https://doi.org/10.1080/23308249.2020.1779651>.
- [118] V. Morshedi, N. Agh, F. Noori, F. Jafari, A. Ghasemi, M.T. Mozanadeh, Effects of single and combined supplementation of dietary probiotic with bovine lactoferrin and xylooligosaccharide on hemato-immunological and digestive enzymes of silvery-black porgy fingerlings, *Ann. Anim. Sci.* 20 (1) (2020) 137–155, <https://sciendo.com/downloadpdf/journals/aoas/20/1/article-p137.xml>.
- [119] K.E. Fernandes, D.A. Carter, The antifungal activity of lactoferrin and its derived peptides: mechanisms of action and synergy with drugs against fungal pathogens, *Front. Microbiol.* 8 (2017) 2, <https://doi.org/10.3389/fmicb.2017.00002>.
- [120] V. Morshedi, B. Bojarski, S. Hamed, H. Torahi, G. Hashemi, C. Faggio, Effects of dietary bovine lactoferrin on growth performance and immuno-physiological responses of Asian sea bass (*Lates calcarifer*) fingerlings, *Probiot. Antimicrob. Protein* 13 (6) (2021) 1790–1797, <https://link.springer.com/article/10.1007/s12602-021-09805-4>.

- [121] H.L. Xia, J. Cheng, D.P. Yu, W.J. Chen, Y.S. Lu, Research progress on peptidoglycan recognition proteins in fish, *Biotechnol. Bull.* 34 (8) (2018) 58, <https://doi.org/10.13560/j.cnki.biotech.bull.1985.2018-0218>.
- [122] S. Roy, P. Bossier, P. Norouzitalab, D. Vanrompay, Trained immunity and perspectives for shrimp aquaculture, *Rev. Aquacult.* 12 (4) (2020) 2351–2370, <https://doi.org/10.1111/raq.12438>.
- [123] H.L. Yang, X. Hu, J.D. Ye, V. Seerengaraj, W. Yang, C.X. Ai, Y.Z. Sun, Cell wall components of *Bacillus pumilus* SE5 improved the growth, digestive and immunity of grouper (*Epinephelus coioides*), *Curr. Chin. Sci.* 1 (2) (2021) 231–239. <https://www.eurekaselect.com/node/189018/article/cell-wall-components-of-bacillus-pumilus-se5-improved-the-growth-digestive-and-immunity-of-grouper-epinephelus-coioides>.
- [124] H. Kondo, T. Kikumoto, K. Yoshii, N. Murase, H. Yamada, Y. Fukuda, I. Hirono, Effects of peptidoglycan and polyinosinic: polycytidylic acid on the recombinant subunit vaccine efficacy against *Edwardsiella tarda* in Japanese flounder *Paralichthys olivaceus*, *Fish Pathol.* 56 (3) (2021) 149–155, <https://doi.org/10.3147/jstfp.56.149>.
- [125] C.Y. Ang, M. Sano, S. Dan, M. Leelakriangsak, T.M. Lal, Postbiotics applications as infectious disease control agent in aquaculture, *Biocontrol Sci.* 25 (1) (2020) 1–7, <https://doi.org/10.4265/bio.25.1>.
- [126] Z. Hu, X. Cao, M. Guo, C. Li, Identification and characterization of a novel short-type peptidoglycan recognition protein in *Apostichopus japonicus*, *Fish Shellfish Immunol.* 99 (2020) 257–266, <https://doi.org/10.1016/j.fsi.2020.02.013>.
- [127] C.R. de Cruz, F.Y. Yamamoto, M. Ju, K. Chen, A. Velasquez, D.M. Gatlin III, Efficacy of purified nucleotide supplements on the growth performance and immunity of hybrid striped bass *Morone chrysops* x *Morone saxatilis*, *Fish Shellfish Immunol.* 98 (2020) 868–874, <https://doi.org/10.1016/j.fsi.2019.11.046>.
- [128] A.P.D.C. Alves, R.R. Paulino, R.T. Pereira, D.V. da Costa, P.V. e Rosa, Nile tilapia fed insect meal: growth and innate immune response in different times under lipopolysaccharide challenge, *Aquacult. Res.* 52 (2) (2021) 529–540, <https://doi.org/10.1111/are.14911>.
- [129] S.W. Luo, N.X. Xiong, Z.Y. Luo, K.K. Luo, S.J. Liu, C. Wu, M. Wen, Effect of Lipopolysaccharide (LPS) stimulation on apoptotic process and oxidative stress in fibroblast cell of hybrid crucian carp compared with those of *Carassius cuvieri* and *Carassius auratus* red var, *Comp. Biochem. Physiol. C Toxicol. Pharmacol.* 248 (2021), 109085, <https://doi.org/10.1016/j.cbpc.2021.109085>.
- [130] M.F. Abd El-Kader, M. Shukry, M.A. Dawood, M.M. Hassan, F. Farrag, A. Omar, E. M. Moustafa, Ameliorative effect of dietary lipopolysaccharides on *Oreochromis niloticus* juveniles submitted to aflatoxin B1-induced oxidative stress and bacterial challenge, *Aquacult. Res.* 52 (8) (2021) 3660–3676, <https://doi.org/10.1111/are.15211>.
- [131] E. Holen, M.H. Austgulen, M. Espe, RNA form baker's yeast cultured with and without lipopolysaccharide (LPS) modulates gene transcription in an intestinal epithelial cell model, RTgutGC from rainbow trout (*Oncorhynchus mykiss*), *Fish Shellfish Immunol.* 119 (2021) 397–408, <https://doi.org/10.1016/j.fsi.2021.10.018>.
- [132] G. Mohammadi, T.J. Adorian, G. Rafiee, Beneficial effects of *Bacillus subtilis* on water quality, growth, immune responses, endotoxemia and protection against lipopolysaccharide-induced damages in *Oreochromis niloticus* under biofloc technology system, *Aquacult. Nutr.* 26 (5) (2020) 1476–1492, <https://doi.org/10.1111/anu.13096>.
- [133] J. Xue, K. Shen, Y. Hu, Y. Hu, Y. Kumar, G. Yang, C. Wen, Effects of dietary *Bacillus cereus*, *B. subtilis*, *Paracoccus marcusii*, and *Lactobacillus plantarum* supplementation on the growth, immune response, antioxidant capacity, and intestinal health of juvenile grass carp (*Ctenopharyngodon idellus*), *Aquacult. Rep.* 17 (2020), 100387, <https://doi.org/10.1016/j.aqrep.2020.100387>.
- [134] H.L. Yang, Y.Z. Sun, X. Hu, J.D. Ye, K.L. Lu, L.H. Hu, J.J. Zhang, *Bacillus pumilus* SE5 originated PG and LTA tuned the intestinal TLRs/MyD88 signaling and microbiota in grouper (*Epinephelus coioides*), *Fish Shellfish Immunol.* 88 (2019) 266–271, <https://doi.org/10.1016/j.fsi.2019.03.005>.
- [135] Y. Yang, J. Lim, C. Li, S. Lee, S. Hong, Effects of sulfated polysaccharides isolated from *Codium fragile* on inflammatory cytokine gene expression and *Edwardsiella tarda* infection in rockfish, *Sebastes schlegelii*, *Fish Shellfish Immunol.* 112 (2021) 125–134, <https://doi.org/10.1016/j.fsi.2021.03.001>.
- [136] C. Jiang, M. Xu, X. Kuang, J. Xiao, M. Tan, Y. Xie, Y. Wu, *Treponema pallidum* flagellins stimulate MMP-9 and MMP-13 expression via TLR5 and MAPK/NF- κ B signaling pathways in human epidermal keratinocytes, *Exp. Cell Res.* 361 (1) (2017) 46–55, <https://doi.org/10.1016/j.yexcr.2017.09.040>.
- [137] Q. Gao, Y. Xiao, C. Zhang, M. Min, S. Peng, Z. Shi, Molecular characterization and expression analysis of toll-like receptor 2 in response to bacteria in silvery pomfret intestinal epithelial cells, *Fish Shellfish Immunol.* 58 (2016) 1–9, <https://doi.org/10.1016/j.fsi.2016.08.057>.
- [138] Q. Gao, Y. Yue, M. Min, S. Peng, Z. Shi, W. Sheng, T. Zhang, Characterization of TLR5 and TLR9 from silver pomfret (*Pampus argenteus*) and expression profiling in response to bacterial components, *Fish Shellfish Immunol.* 80 (2018) 241–249, <https://doi.org/10.1016/j.fsi.2018.06.014>.
- [139] Á. González-Garcinúño, A. Tabernero, Á. Domínguez, M.A. Galán, E.M. Martín del Valle, Levan and levansucrases: polymer, enzyme, micro-organisms and biomedical applications, *Biocatal. Biotransform.* 36 (3) (2018) 233–244, <https://doi.org/10.1080/10242422.2017.1314467>.
- [140] M.Y. Huang, C.I. Chang, C.C. Chang, L.W. Tseng, C.L. Pan, Effects of dietary levan on growth performance, nonspecific immunity, pathogen resistance and body composition of orange-spotted grouper (*Epinephelus coioides* H.), *Aquacult. Res.* 46 (11) (2015) 2752–2767. <https://onlinelibrary.wiley.com/doi/abs/10.1111/are.12430>.
- [141] S. Yilmaz, E. Yilmaz, M.A. Dawood, E. Ringø, E. Ahmadifar, H.M. Abdel-Latif, Probiotics, prebiotics, and synbiotics used to control vibriosis in fish: a review, *Aquaculture* 547 (2022), 737514, <https://doi.org/10.1016/j.aquaculture.2021.737514>.
- [142] K. Mohan, T. Muralisankar, V. Uthayakumar, R. Chandrasekar, N. Revathi, A. R. Ganesan, P. Seedeivi, Trends in the extraction, purification, characterisation and biological activities of polysaccharides from tropical and sub-tropical fruits—A comprehensive review, *Carbohydr. Polym.* 238 (2020), 116185, <https://doi.org/10.1016/j.carbpol.2020.116185>.
- [143] F. Motta, M.E. Gershwin, C. Selmi, Mushrooms and immunity, *J. Autoimmun.* 117 (2021), 102576, <https://doi.org/10.1016/j.jaut.2020.102576>.
- [144] B. Deng, Z.P. Wang, W.J. Tao, W.F. Li, C. Wang, M.Q. Wang, D. Wu, Effects of polysaccharides from mycelia of *Cordyceps sinensis* on growth performance, immunity and antioxidant indicators of the white shrimp *Litopenaeus vannamei*, *Aquacult. Nutr.* 21 (2) (2015) 173–179, <https://doi.org/10.1111/anu.12147>.
- [145] K. Mohan, A.M. Padmanaban, V. Uthayakumar, R. Chandrasekar, T. Muralisankar, P. Santhanam, Effect of dietary *Ganoderma lucidum* polysaccharides on biological and physiological responses of the giant freshwater prawn *Macrobrachium rosenbergii*, *Aquaculture* 464 (2016) 42–49, <https://doi.org/10.1016/j.aquaculture.2016.05.046>.
- [146] G. Mohammadi, M. Hafezieh, A.A. Karimi, M.N. Azra, H. Van Doan, W. Tapingkae, M.A. Dawood, The synergistic effects of plant polysaccharide and *Pediococcus acidilactici* as a synbiotic additive on growth, antioxidant status, immune response, and resistance of Nile tilapia (*Oreochromis niloticus*) against *Aeromonas hydrophila*, *Fish Shellfish Immunol.* 120 (2022) 304–313, <https://doi.org/10.1016/j.fsi.2021.11.028>.
- [147] H. Do Huu, H.M. Sang, N.T.T. Thuy, Dietary β -glucan improved growth performance, Vibrio counts, haematological parameters and stress resistance of pompano fish, *Trachinotus ovatus* Linnaeus, 1758, *Fish Shellfish Immunol.* 54 (2016) 402–410, <https://doi.org/10.1016/j.fsi.2016.03.161>.
- [148] X. Xu, X. Zhang, Lentinula edodes-derived polysaccharide alters the spatial structure of gut microbiota in mice, *PLoS One* 10 (1) (2015), e0115037, <https://doi.org/10.1371/journal.pone.0115037>.
- [149] G. Ren, M. Yu, K. Li, Y. Hu, Y. Wang, X. Xu, J. Qu, Seleno-lentinan prevents chronic pancreatitis development and modulates gut microbiota in mice, *J. Funct. Foods* 22 (2016) 177–188, <https://doi.org/10.1016/j.jff.2016.01.035>.
- [150] S. Shahbazi, A. Bolhassani, Immunostimulants: types and functions, *J. Med. Microbiol. Infect. Dis.* 4 (3) (2016) 45–51. <https://jommid.pasteur.ac.ir/article-1-131-en.html>.
- [151] S. Xiang, X. Chen, J. Wang, H. Peng, X. Lv, L. Jiang, L. Ma, A lentinan-loaded hydrogel with core-shell structure induces broad-spectrum resistance against plant virus by activating the expression of CML30, *Author Preprints*, <https://www.authorea.com/doi/full/10.22541/au.160503099.94865844>, 2020.
- [152] W. Yao, X. Li, C. Zhang, J. Wang, Y. Cai, X. Leng, Effects of dietary synbiotics supplementation methods on growth, intestinal health, non-specific immunity and disease resistance of Pacific white shrimp, *Litopenaeus vannamei*, *Fish Shellfish Immunol.* 112 (2021) 46–55, <https://doi.org/10.1016/j.fsi.2021.02.011>.
- [153] F. Li, F. Gao, J. Tan, C. Fan, H. Sun, J. Yan, X. Wang, Characterization and identification of enzyme-producing microflora isolated from the gut of sea cucumber *Apostichopus japonicus*, *Chin. J. Oceanol. Limnol.* 34 (1) (2016) 153–162, <https://doi.org/10.1007/s00343-015-4149-z>.
- [154] R. Yuji-Sado, F. Raulino-Domanski, P.F. de Freitas, F. Baioco-Sales, Growth, immune status and intestinal morphology of Nile tilapia fed dietary prebiotics (mannan oligosaccharides-MOS), *Lat. Am. J. Aquat. Res.* 43 (5) (2015) 944–952. <https://www.redalyc.org/pdf/1750/175042668014.pdf>.
- [155] Z.Y. Lu, W.D. Jiang, P. Wu, Y. Liu, S.Y. Kuang, L. Tang, L. Feng, Mannan oligosaccharides supplementation enhanced head-kidney and spleen immune function in on-growing grass carp (*Ctenopharyngodon idella*), *Fish Shellfish Immunol.* 106 (2020) 596–608, <https://doi.org/10.1016/j.fsi.2020.08.035>.
- [156] M. Abdel Gayed, H. Elabd, M. Shehabeldin, A. Shaheen, A. El-Asely, A. El-Gawad, A. Abbas, A Review of some prebiotics and probiotics supplementation effects on farmed fishes: with special reference to Mannan oligosaccharides (MOS), *Benha Vet. Med. J.* 40 (1) (2021) 141–145, <https://doi.org/10.21608/bvmj.2021.62545.1342>.
- [157] M.A. Dawood, S. Koshio, S.E. Fadl, H.A. Ahmed, A. El Asely, M.M. Abdel-Daim, S. Alkahtani, The modulatory effect of mannanoligosaccharide on oxidative status, selected immune parameters and tolerance against low salinity stress in red sea bream (*Pagrus major*), *Aquacult. Rep.* 16 (2020), 100278, <https://doi.org/10.1016/j.aqrep.2020.100278>.
- [158] M.N. Akter, K. Zahan, M.A. Zafar, N. Khatun, M.S. Rana, M.I. Mursalin, Effects of dietary mannan oligosaccharide on growth performance, feed utilization, body composition and haematological parameters in asian catfish (*Clarias batrachus*) juveniles, *Turk. J. Fish. Aquat. Sci.* 21 (11) (2021) 559–567, https://doi.org/10.4194/1303-2712-v21_11_04.
- [159] S. Ayiku, J. Shen, B.P. Tan, X.H. Dong, H.Y. Liu, Effects of reducing dietary fishmeal with yeast supplementations on *Litopenaeus vannamei* growth, immune response and disease resistance against *Vibrio harveyi*, *Microbiol. Res.* 239 (2020), 126554, <https://doi.org/10.1016/j.micres.2020.126554>.
- [160] E. De Marco Castro, P.C. Calder, H.M. Roche, β -1, 3/1, 6-glucans and immunity: state of the art and future directions, *Mol. Nutr. Food Res.* 65 (1) (2021), 1901071, <https://doi.org/10.1002/mnfr.201901071>.
- [161] M.H. Khanjani, G. Ghaedi, M. Sharifnia, Effects of diets containing β -glucan on survival, growth performance, haematological, immunity and biochemical parameters of rainbow trout (*Oncorhynchus mykiss*) fingerlings, *Aquacult. Res.* 53 (5) (2022) 1842–1850, <https://doi.org/10.1111/are.15712>.

- [162] A. Chhetri, The role of (1, 3)- β -D-Glucan synthase, in: *Fungal Cell Wall Biosynthesis* (Doctoral Dissertation, Duke University, 2020). <https://www.proquest.com/ope/nview/7ad737cf781564ac5c8d4b24e57de6f6/1?pq-origsite=gscholar&cbl=18750&diss=y>.
- [163] R. Pogue, E.J. Murphy, G.W. Fehrenbach, E. Rezoagli, N.J. Rowan, Exploiting immunomodulatory properties of β -glucans derived from natural products for improving health and sustainability in aquaculture-farmed organisms: concise review of existing knowledge, innovation and future opportunities, *Curr. Opin. Environ. Sci. Health* 21 (2021), 100248, <https://doi.org/10.1016/j.coesh.2021.100248>.
- [164] M.A. Dawood, S. Koshio, M. Ishikawa, S. Yokoyama, Interaction effects of dietary supplementation of heat-killed *Lactobacillus plantarum* and β -glucan on growth performance, digestibility and immune response of juvenile red sea bream, *Pagrus major*, *Fish Shellfish Immunol.* 45 (1) (2015) 33–42, <https://doi.org/10.1016/j.fsi.2015.01.033>.
- [165] M. Marcos-López, H.D. Rodger, Amoebic gill disease and host response in Atlantic salmon (*Salmo salar* L.): a review, *Parasite Immunol.* 42 (8) (2020), e12766, <https://doi.org/10.1111/pim.12766>.
- [166] J.Y. Loh, K.S. Lai, P.T. Lee, H.J. Liew, A.S.Y. Ting, Effects of Artemia nauplii bioencapsulated with *Lactococcus lactis* subsp. *lactis* CF4MRS and sodium alginate on Edwardsiella protection and digestive enzyme production in climbing perch larvae, *Anabas testudineus* (Bloch, 1792), *J. Appl. Aquacult.* (2021) 1–18, <https://doi.org/10.1080/10454438.2021.1935389>.
- [167] J.Y. Loh, H.K. Chan, H.C. Yam, L.L.A. In, C.S.Y. Lim, An overview of the immunomodulatory effects exerted by probiotics and prebiotics in grouper fish, *Aquacult. Int.* 28 (2) (2020) 729–750. <https://link.springer.com/article/10.1007/s10499-019-00491-2>.
- [168] M. Chen, X.Q. Chen, L.X. Tian, Y.J. Liu, J. Niu, Beneficial impacts on growth, intestinal health, immune responses and ammonia resistance of Pacific white shrimp (*Litopenaeus vannamei*) fed dietary synbiotic (mannan oligosaccharide and *Bacillus licheniformis*), *Aquacult. Rep.* 17 (2020), 100408, <https://doi.org/10.1016/j.aqrep.2020.100408>.
- [169] V.T. Ojerio, V.L. Corre, N.A. Toledo, K.G.S. Andriano-Felarcia, L.M. Nievaes, R.F. M. Traifalgar, Alginic acid as immunostimulant: effects of dose and frequency on growth performance, immune responses, and white spot syndrome virus resistance in tiger shrimp *Penaeus monodon* (Fabricius, 1798), *Aquacult. Int.* 26 (1) (2018) 267–278. <https://link.springer.com/article/10.1007/s10499-017-0212-4>.
- [170] H. Tayefi Nasrabadi, M. Heidarieh, N. Sheikhzadeh, S. Niksefat Zende, P. Karimi, Effects of irradiated Ergosan on growth performance, serum biochemical parameters and skin mucosal immune responses in crucian carp (*Carassius carassius*), *Aquat. Physiol. Biotechnol.* 9 (2) (2021) 107–124, <https://doi.org/10.22124/japb.2021.17238.1389>.
- [171] F.S. Farooqi, W.U.H. Qureshi, Immunostimulants for Aquaculture. health management, *J. Pharmacogn. Phytochem.* 7 (2018) 1441–1447. <https://www.phytojournal.com/archives/2018/vol7issue6/PartY/7-5-684-797.pdf>.
- [172] E. Valjeos-Vidal, F. Reyes-López, M. Teles, S. MacKenzie, The response of fish to immunostimulant diets, *Fish Shellfish Immunol.* 56 (2016) 34–69, <https://doi.org/10.1016/j.fsi.2016.06.028>.
- [173] D.L. Prabu, N.P. Sahu, A.K. Pal, S. Dasgupta, A. Narendra, Immunomodulation and interferon gamma gene expression in sutchi cat fish, *Pangasianodon hypophthalmus*: effect of dietary fucoidan rich seaweed extract (FRSE) on pre and post challenge period, *Aquacult. Res.* 47 (1) (2016) 199–218, <https://doi.org/10.1111/are.12482>.
- [174] M. Saeed, M.A. Arain, S. Ali Fazlani, I.B. Marghazani, M. Umar, J. Soomro, M. Alagawany, A comprehensive review on the health benefits and nutritional significance of fucoidan polysaccharide derived from brown seaweeds in human, animals and aquatic organisms, *Aquacult. Nutr.* 27 (3) (2021) 633–654, <https://doi.org/10.1111/anu.13233>.
- [175] C.H. Hsu, J.C. Chen, Y.C. Lin, Y.Y. Chen, P.C. Liu, B.W. Lin, J.F. Hsieh, White shrimp *Litopenaeus vannamei* that have received mixtures of heat-killed and formalin-inactivated *Vibrio alginolyticus* and *V. harveyi* exhibit recall memory and show increased phagocytosis and resistance to *Vibrio* infection, *Fish Shellfish Immunol.* 112 (2021) 151–158, <https://doi.org/10.1016/j.fsi.2020.11.013>.
- [176] R. Salehpour, N.A. Biuki, M. Mohammadi, A. Dashtiannasab, P. Ebrahimnejad, The dietary effect of fucoidan extracted from brown seaweed, *Cystoseira trindis* (C. Agardh) on growth and disease resistance to WSSV in shrimp *Litopenaeus vannamei*, *Fish Shellfish Immunol.* 119 (2021) 84–95, <https://doi.org/10.1016/j.fsi.2021.09.005>.
- [177] S.U. Kadam, C.P. O'Donnell, D.K. Rai, M.B. Hossain, C.M. Burgess, D. Walsh, B. K. Tiwari, Laminarin from Irish brown seaweeds *Ascophyllum nodosum* and *Laminaria hyperborea*: ultrasound assisted extraction, characterization and bioactivity, *Mar. Drugs* 13 (7) (2015) 4270–4280, <https://doi.org/10.3390/md13074270>.
- [178] S.U. Kadam, B.K. Tiwari, C.P. O'Donnell, Extraction, structure and biofunctional activities of laminarin from brown algae, *Int. J. Food Sci. Technol.* 50 (1) (2015) 24–31, <https://doi.org/10.1111/ijfs.12692>.
- [179] Y. Meng, F. Lyu, X. Xu, L. Zhang, Recent advances in chain conformation and bioactivities of triple-helix polysaccharides, *Biomacromolecules* 21 (5) (2020) 1653–1677, <https://doi.org/10.1021/acs.biomac.9b01644>.
- [180] C. Cui, L. Zhu, X. Tang, J. Xing, X. Sheng, H. Chi, W. Zhan, Differential white spot syndrome virus-binding proteins in two hemocyte subpopulations of Chinese shrimp (*Penaeus chinensis*), *Dev. Comp. Immunol.* 125 (2021), 104215, <https://doi.org/10.1016/j.dci.2021.104215>.
- [181] G. Shanmugavel, G. Krishnamoorthy, Mucuna pruriens-seed extract stimulating immune system in *Litopenaeus vannamei* against *Vibrio harveyi*, *World News Nat. Sci.* 39 (2021) 95–112. <https://bibliotekanauki.pl/articles/1839790>.
- [182] D.S. Lakshmi, S. Sankaranarayanan, T.K. Gajaria, G. Li, W. Kujawski, J. Kujawa, R. Navia, A short review on the valorization of green seaweeds and ulvan: feedstock for chemicals and biomaterials, *Biomolecules* 10 (7) (2020) 991, <https://doi.org/10.3390/biom10070991>.
- [183] P. del Rocio Quezada-Rodríguez, E.J. Fajer-Ávila, The dietary effect of ulvan from *Ulva clathrata* on hematological-immunological parameters and growth of tilapia (*Oreochromis niloticus*), *J. Appl. Phycol.* 29 (1) (2017) 423–431, <https://doi.org/10.1007/s10811-016-0903-7>.
- [184] M. Ponce, E. Zuasti, V. Anguís, C. Fernández-Díaz, Effects of the sulfated polysaccharide ulvan from *Ulva ohnoi* on the modulation of the immune response in Senegalese sole (*Solea senegalensis*), *Fish Shellfish Immunol.* 100 (2020) 27–40, <https://doi.org/10.1016/j.fsi.2020.02.054>.
- [185] R. Harikrishnan, G. Devi, H. Van Doan, C. Balasundaram, J. Arockiaraj, C. Jagruthi, Efficacy of ulvan on immune response and immuno-antioxidant gene modulation in *Labeo rohita* against columnaris disease, *Fish Shellfish Immunol.* 117 (2021) 262–273, <https://doi.org/10.1016/j.fsi.2021.08.004>.
- [186] Q.D. Lauzon, A.E. Serrano Jr., Ulvan extract from *Enteromorpha intestinalis* enhances immune responses in *Litopenaeus vannamei* and *Penaeus monodon* juveniles, *Anim. Biol. Anim. Husband.* 7 (1) (2015) 1–10. <https://doi.org/10.1016/j.dci.2015.07.007>.
- [187] M. He, G. Liu, Y. Liu, K. Yang, X. Qi, A. Huang, E. Wang, Effects of geniposide as immunostimulant on the innate immune response and disease resistance in crucian carp, *Aquaculture* 529 (2020), 735713, <https://doi.org/10.1016/j.aquaculture.2020.735713>.
- [188] C. Qin, Z. Zhang, Y. Wang, S. Li, C. Ran, J. Hu, Z. Zhou, EPSP of L. Casei BL23 protected against the infection caused by *Aeromonas veronii* via enhancement of immune response in zebrafish, *Front. Microbiol.* 8 (2017) 2406. <https://www.frontiersin.org/articles/10.3389/fmicb.2017.02406/full>.
- [189] C.Y. Wang, Z.B. Li, Growth performance, digestive enzyme activity and immune response of Japanese sea bass, *Lateolabrax japonicus* fed with fructooligosaccharide, *Aquacult. Nutr.* 26 (2) (2020) 296–305, <https://doi.org/10.1111/anu.12990>.
- [190] A. de Lima Paz, J.M. da Silva, K.M.M. da Silva, A.L. Val, Protective effects of the fructooligosaccharide on the growth performance, hematology, immunology indicators and survival of tambaqui (*Colossoma macropomum*, Characiformes: serrasalmidae) infected by *Aeromonas hydrophila*, *Aquacult. Rep.* 15 (2019), 100222, <https://doi.org/10.1016/j.aqrep.2019.100222>.
- [191] L. Hu, H. Yang, Y. Zhao, J. Ye, K. Lu, Y. Sun, Effects of fructooligosaccharide on immune function and intestinal morphology of grouper *Epinephelus coioides*, *Fish. Sci.* 38 (5) (2019) 589–594. <https://www.cabdirect.org/cabdirect/abstract/20193471302>.
- [192] E. Pietrzak, J. Mazurkiewicz, A. Slawinska, Innate immune responses of skin mucosa in common carp (*Cyprinus carpio*) fed a diet supplemented with galactooligosaccharides, *Animals* 10 (3) (2020) 438. <https://www.mdpi.com/2076-2615/10/3/438>.
- [193] A. Mustafa, A. Buentello, D. Gatlin, D. Lightner, M. Hume, A. Lawrence, Dietary supplementation of galactooligosaccharides (GOS) in Pacific white shrimp, *Litopenaeus vannamei*, cultured in a recirculating system and its effects on gut microflora, growth, stress, and immune response, *J. Immunoassay Immunochem.* 40 (6) (2019) 662–675, <https://doi.org/10.1080/15321819.2019.1675694>.
- [194] A. Ajdari, H. Ghafarifarani, S.H. Hoseinifar, S. Javahery, F. Narimanizad, K. Gatphayak, H. Van Doan, Effects of dietary supplementation of PrimaLac, inulin, and biomin imbo on growth performance, antioxidant, and innate immune responses of common carp (*Cyprinus carpio*), *Aquacult. Nutr.* (2022), <https://doi.org/10.1155/2022/8297479>, 2022.
- [195] L. Zhou, J. Zhang, M. Yan, S. Tang, X. Wang, J.G. Qin, E. Li, Inulin alleviates hypersaline-stress induced oxidative stress and dysbiosis of gut microbiota in Nile tilapia (*Oreochromis niloticus*), *Aquaculture* 529 (2020), 735681, <https://doi.org/10.1016/j.aquaculture.2020.735681>.
- [196] A.G. Huang, W.H. He, F.L. Zhang, C.S. Wei, Y.H. Wang, Natural component geniposide enhances survival rate of crayfish *Procambarus clarkii* infected with white spot syndrome virus, *Fish Shellfish Immunol.* (2022), <https://doi.org/10.1016/j.fsi.2022.05.037>.
- [197] S.H. Hoseinifar, M.A. Jahazi, N. Nikdehghan, H. Van Doan, M.G. Volpe, M. Paolucci, Effects of dietary polyphenols from agricultural by-products on mucosal and humoral immune and antioxidant responses of convict cichlid (*Amatitlania nigrofasciata*), *Aquaculture* 517 (2020), 734790, <https://doi.org/10.1016/j.aquaculture.2019.734790>.
- [198] A. Ahmadi, D. Bagheri, S.H. Hoseinifar, V. Morshedi, M. Paolucci, Beneficial role of polyphenols as feed additives on growth performances, immune response and antioxidant status of Lates Calcarifer (Bloch, 1790) juveniles, *Aquaculture* 552 (2022), 737955, <https://doi.org/10.1016/j.aquaculture.2022.737955>.
- [199] L. Zhao, B.D. Yuan, J.L. Zhao, N. Jiang, A.Z. Zhang, G.Q. Wang, M.Y. Li, Amelioration of hexavalent chromium-induced bioaccumulation, oxidative stress, tight junction proteins and immune-related signaling factors by Allium mongolicum Regel flavonoids in *Ctenopharyngodon idella*, *Fish Shellfish Immunol.* 106 (2020) 993–1003, <https://doi.org/10.1016/j.fsi.2020.09.005>.
- [200] P. Bussabong, T. Rairat, N. Chuchird, A. Keetanon, P. Phansawat, K. Cherdkeatipol, W. Kraitavin, Effects of isoquinoline alkaloids from *Macleaya cordata* on growth performance, survival, immune response, and resistance to *Vibrio parahaemolyticus* infection of Pacific white shrimp (*Litopenaeus vannamei*), *PLoS One* 16 (5) (2021), e0251343. <https://journals.plos.org/plosone/article?id=10.1371/journal.pone.0251343>.

- [202] J. Jiang, S. Gao, X. Wang, X. Guan, B. Wang, Z. Chen, Z. Zhou, The role of a novel secretory peptidoglycan recognition protein from the sea cucumber *Apostichopus japonicus* in innate immunity, *Aquaculture* 546 (2022), 737339, <https://doi.org/10.1016/j.aquaculture.2021.737339>.
- [203] X. Li, S. Yuan, Z. Sun, L. Lei, S. Wan, J. Wang, Q. Gao, Gene identification and functional analysis of peptidoglycan recognition protein from the spotted sea bass, *Fish Shellfish Immunol.* 106 (2020) 1014–1024. <https://www.sciencedirect.com/science/article/abs/pii/S1050464820306057>.
- [204] M.Y. Li, X.M. Zhu, X.T. Niu, X.M. Chen, J.X. Tian, Y.D. Kong, G.Q. Wang, Effects of dietary *Allium mongolicum* Regel polysaccharide on growth, lipopolysaccharide-induced antioxidant responses and immune responses in *Channa argus*, *Mol. Biol. Rep.* 46 (2) (2019) 2221–2230. <https://link.springer.com/article/10.1007/s11033-019-04677-y>.
- [205] Y. Risjani, N. Mutmainnah, P. Manurung, S.N. Wulan, Exopolysaccharide from *Porphyridium cruentum* (purpureum) is not toxic and stimulates immune response against vibriosis: the assessment using zebrafish and white shrimp *Litopenaeus vannamei*, *Mar. Drugs* 19 (3) (2021) 133. <https://www.mdpi.com/1660-3397/19/3/133>.
- [206] C. Ma, H. Guo, H. Chang, S. Huang, S. Jiang, D. Huo, X. Zhu, The effects of exopolysaccharides and exopolysaccharide-producing *Lactobacillus* on the intestinal microbiome of zebrafish (*Danio rerio*), *BMC Microbiol.* 20 (1) (2020) 1–11. <https://bmcmicrobiol.biomedcentral.com/articles/10.1186/s12866-020-01990-6>.
- [207] S.K. Gupta, Ameliorative and protective effects of prebiotic, microbial levan in common carp, (*Cyprinus carpio*) fry under experimental exposure to fipronil, *Int. J. Aquat. Biol.* 9 (2) (2021) 134–147. <https://ij-aquaticbiology.com/index.php/ijab/article/view/866>.
- [208] S. Anusha, T. Neeraja, P. Haribabu, P. Akshaya, Effect of different biofloc based culture systems on the growth and immune response of *Tilapia* (*Oreochromis niloticus*), *Survival* 2 (2020) 63. <https://www.fisheriesjournal.com/archives/2020/vol8issue6/PartC/8-5-67-508.pdf>.
- [209] C. Carballo, P.I. Pinto, A.P. Mateus, C. Berbel, C.C. Guerreiro, J.F. Martinez-Blanch, M. Manchado, Yeast β -glucans and microalgal extracts modulate the immune response and gut microbiome in Senegalese sole (*Solea senegalensis*), *Fish Shellfish Immunol.* 92 (2019) 31–39, <https://doi.org/10.1016/j.fsi.2019.05.044>.