

Lower GI

Median ID50 (log10)					Culture
1.2	>7.4	2.5	2	6.5	
4.1	1.2	<1.2	1.4	1.7	
6.5	>7.4	7.4	7.4	7.1	
					Total RNA
<1.2	<1.2	<1.2	<1.2	<1.2	
<1.2	<1.2	<1.2	<1.2	<1.2	
<1.2	<1.2	1.2	1.4	<1.2	
IN	IT	IN+IT	AE	IG	

	Diff. between Max & Min Dose					
IN	+38	-0.4		+1.4		
IT	+2					
IN+IT	+46	-0.4	-0.6	+1.6		+1.4
AE	+50	-0.6		+2.6		
IG	+13					
IN	+9		-1.2	+1.7		
IT	+13	-0.2				
IN+IT	+19		-1.6	+1.4		
AE	+36		-2.3	+2.9		
IG	+8					
IN	+21		+1.8	+1.4	+2.8	+2
IT	+5					
IN+IT	+25			+1.8	+2.5	+2.3
AE	+30		+3.1	+2.7	+7.7	+5.1
IG	+24	-0.7		+1.2		

Figure 2 displays three stacked line graphs showing viral titers (log₁₀ pfu) over 12 days post-infection for three different viral strains: purple (top), yellow (middle), and red (bottom). The x-axis for all graphs is 'Days post infection' (0 to 12). The y-axis for all graphs is 'Viral titer (log₁₀ pfu)'. Each graph shows a peak around day 4.

Days post infection	Purple Viral Titer (log ₁₀ pfu)	Yellow Viral Titer (log ₁₀ pfu)	Red Viral Titer (log ₁₀ pfu)
0	0	0	0
2	~3.5	~1.5	~1.5
4	~3.5	~3.5	~4.5
6	~1.5	~1.5	~1.5
8	0	0	0