

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/248018135>

# The solution of the linear bilevel programming problem by using the linear complementarity problem

Article

---

CITATIONS

25

---

READS

22

2 authors, including:



Ana M. Faustino

University of Porto

16 PUBLICATIONS 234 CITATIONS

SEE PROFILE

## THE SOLUTION OF THE LINEAR BILEVEL PROGRAMMING PROBLEM BY USING THE LINEAR COMPLEMENTARITY PROBLEM

J. J. Júdice\*  
A. M. Faustino\*\*

\*Dept. de Matemática, Universidade de Coimbra  
\*\*Dept. de Matemática, Universidade do Porto

### Resumo

O Problema Linear de 2 níveis (LBP) é um problema de optimização não convexa que aparece muito frequentemente em aplicações. Bialas e Karwan desenvolveram um algoritmo SLCP para a solução do LBP que consiste em resolver uma sucessão de Problemas Lineares Complementares. Contudo, o algoritmo não é sempre convergente. Neste artigo é proposta uma versão modificada e convergente desse algoritmo SLCP. Experiência computacional com o algoritmo SLCP modificado na solução de LBPs de média dimensão e estrutura esparsa é também incluída e mostra que o processo é eficiente.

### Abstract

The Linear Bilevel Programming Problem (LBP) is a nonconvex optimization problem that occurs quite frequently in applications. Bialas and Karwan have designed a SLCP algorithm for the solution of the LBP, which consists of solving a sequence of linear complementarity problems. However, the method is not convergent in general. In this paper we describe a modified convergent version of the SLCP algorithm. Computational experience with the modified SLCP algorithm on the solution of medium-scale LBPs is also included and shows the efficiency of the approach.

**Keywords:** Bilevel Programming, Nonconvex Optimization, Linear Complementarity Problem, SLCP algorithms, sparse matrices.

### 1 - INTRODUCTION

The Bilevel Programming Problem (BLP) can be defined as

$$\left. \begin{array}{l} \text{Minimize } f_1(x,y) \\ y \in R^m \\ \text{subject to } g_i(x,y) \leq 0, \quad i = 1, \dots, l \end{array} \right\} \quad (1)$$

where  $x \in R^n$  is implicitly defined as the optimal solution of the program

$$\left. \begin{array}{l} \text{Minimize } f_2(x,y) \\ x \in R^n \\ \text{subject to } h_i(x,y) \leq 0, \quad i = 1, \dots, r \end{array} \right\} \quad (2)$$

and  $f_1, f_2, g_1$  and  $h_1$  are real functions of the variables  $x_j$  and  $y_j$ .

The BLP is nowadays a well-known nonconvex optimization problem. A large number of applications of the BLP have been proposed, primarily in economic planning [6, 11, 12]. A number of algorithms [2, 3, 4, 5, 6, 7, 9, 12] have been developed for finding a global or a local minimum for the BLP.

In [7], Biales and Karwan study the linear BLP (LBLP), that is, the BLP in which all the functions are linear. They discuss a number of algorithms and recommend a Sequential Linear Complementarity Problem (SLCP) algorithm for the solution of LBLPs of small dimension ( $n \leq 50, m \leq 50$ ). This algorithm consists of solving a sequence of linear complementarity problems (LCP) by using a Restricted Basis Entry Simplex method similar to the Wolfe-Markowitz method [13, 17] for quadratic programming. However, such scheme is not convergent in general [5].

In this paper we describe a modification of the SLCP algorithm capable of finding a global minimum for the LBLP. In this procedure, the LCPs required by the sequential algorithm are solved by an extension of Al-Khayyal's enumerative method [1]. We show that the computational effort of this hybrid enumerative method is highly reduced because of the special structure of the LCPs. We also present some computational experience with medium-scale LBLPs ( $n \leq 150$  and  $m \leq 250$ ) which shows the efficiency of the modified SLCP algorithm.

The organization of the paper is as follows. In Section 2 Biales and Karwan SLCP algorithm for the LBLP is described. In Section 3 the incorporation of the hybrid enumerative method in the SLCP algorithm is discussed. Finally computational experience on the solution of medium-scale LBLPs by the modified SLCP method is presented in Section 4.

## 2 - SLCP ALGORITHM FOR THE LBLP

Consider the LBLP as stated in [7]

$$\left. \begin{array}{l} \text{Minimize } c^T x + d^T y \\ y \in R_+^m \end{array} \right\} \quad (3)$$

where  $R_+^m = \{y \in R^m : y \geq 0\}$  and  $x \in R^n$  solves the linear program

$$\left. \begin{array}{l} \text{Minimize } a^T x \\ x \in R^n \\ \text{subject to } A_1 x + A_2 y \geq b, x \geq 0 \end{array} \right\} \quad (4)$$

where  $A_1 \in R^{m \times n}$  and  $A_2 \in R^{m \times m}$ . As in [7], we assume that the set

$$H = \{(x, y) \in R^{n+m} : A_1 x + A_2 y \geq b, x \geq 0, y \geq 0\} \quad (5)$$

is bounded, so that the LBLP has an optimal solution [7].

Consider the dual of the linear program (4)

$$\left. \begin{array}{l} \text{Maximize } (b - A_2 y)^T u \\ \text{subject to } A_1^T u \leq a \\ u \geq 0 \end{array} \right\} \quad (6)$$

If  $\alpha$  and  $\beta$  are the vectors of the slack variables corresponding to the primal and dual respectively, then by the complementarity slackness property [14, pages 166-167], the LBLP is equivalent to the following nonconvex optimization problem

$$\left. \begin{array}{l} \text{Minimize } c^T x + d^T y \\ \text{subject to } \alpha = -b + A_1 x + A_2 y \\ \beta = a - A_1^T u \\ x, y, u, \alpha, \beta \geq 0, x^T \beta = u^T \alpha = 0 \end{array} \right\} \quad (7)$$

Hence a global minimum  $(\bar{x}, \bar{y})$  of the LBLP can be found by solving the nonconvex program (7). Biales and Karwan [7] propose a SLCP algorithm for such purpose. In this method a parameter  $\lambda$  is introduced and the objective function is replaced by the constraint  $c^T x + d^T y \leq \lambda$  to obtain the following parametric LCP

LCP( $\lambda$ ):

$$\left. \begin{array}{l} \left[ \begin{array}{c} \alpha \\ \beta \\ v_0 \end{array} \right] = \left[ \begin{array}{c} -b \\ a \\ 0 \end{array} \right] + \left[ \begin{array}{c} 0 \\ 0 \\ 1 \end{array} \right] \lambda + \left[ \begin{array}{ccc} 0 & A_1 & A_2 \\ -A_1^T & 0 & 0 \\ 0 & -c^T & -d^T \end{array} \right] \left[ \begin{array}{c} u \\ x \\ y \end{array} \right] \\ x, y, u, \alpha, \beta, v_0 \geq 0, x^T \beta = u^T \alpha = 0 \end{array} \right\} \quad (8)$$

The SLCP algorithm consists of finding the solution of the LCP( $\bar{\lambda}$ ), where  $\bar{\lambda}$  is the smallest value of  $\lambda$  such that LCP( $\lambda$ ) has a solution. To achieve this, the method solves a sequence of LCP( $\lambda_k$ ), where  $(\lambda_k)$  is a decreasing sequence defined by

$$\left. \begin{aligned} \lambda_0 &= \text{upper-bound of } c^T x + d^T y \text{ on } H \\ \lambda_k &= c^T x^{k-1} + d^T y^{k-1} - \left\{ \gamma_k (c^T x^{k-1} + d^T y^{k-1}) \right\} \end{aligned} \right\} \quad (9)$$

with  $(x^{k-1}, y^{k-1})$  the solution of  $LCP(\lambda_{k-1})$  and  $\gamma_k$  a small positive number. The method terminates in a iteration  $k$  such that  $LCP(\lambda_k)$  has no solution. When this occurs, the solution  $(x^{k-1}, y^{k-1})$  of the  $LCP(\lambda_{k-1})$  satisfies

$$0 \leq c^T x^{k-1} + d^T y^{k-1} - \text{VAL} \leq \left\{ \gamma_k (c^T x^{k-1} + d^T y^{k-1}) \right\} \quad (10)$$

where VAL is the value of the objective function at the optimal solution. Hence, if the set  $H$  defined by (5) is nonempty and bounded, the algorithm finds an  $\epsilon$ -optimal solution of the LBLP, where

$$\epsilon = \left\{ \gamma_k (c^T x^{k-1} + d^T y^{k-1}) \right\} \quad (11)$$

In practice, if  $\gamma_k$  is quite small, the solution  $(x^{k-1}, y^{k-1})$  of the last  $LCP(\lambda_{k-1})$  is usually the optimal solution of the LBLP. Biales and Karwan [7] recommend the choice of  $\gamma_k = 0.01$  in each iteration  $k$  and our experience has confirmed that this choice is usually the most suitable.

The steps of the SLCP algorithm are presented below.

#### SLCP Algorithm

STEP 0 - Let  $k = 0$  and  $\lambda_0$  be an upper-bound of  $c^T x + d^T y$  on  $H$ .

General Step - Solve the  $LCP(\lambda_k)$ . If  $LCP(\lambda_k)$  has no solution, go to Exit.

Otherwise let  $(x^k, y^k)$  be the solution of this LCP. Set

$$\lambda_{k+1} = c^T x^k + d^T y^k - \left\{ \gamma_k (c^T x^k + d^T y^k) \right\} \quad (12)$$

where  $\gamma_{k+1}$  is a fixed value.

Set  $k = k + 1$  and repeat.

EXIT - If  $k = 0$  the LBLP is infeasible. Otherwise  $(x^{k-1}, y^{k-1})$  is an  $\epsilon$ -optimal solution of the LBLP, where  $\epsilon$  is given by (11).

The efficiency of the SLCP algorithm depends essentially on the procedure to solve the  $LCP(\lambda_k)$ . Biales and Karwan [7] recommend the use of a Basis Restricted Entry Simplex (BRES) method for this purpose. They do not provide sufficient details for the complete understanding of the algorithm, but this kind of method can not always be used to process the  $LCP(\lambda_k)$  [5]. Next we describe a BRES algorithm with exactly one artificial variable. Consider the  $LCP(\lambda_k)$  in the following form

$$\left. \begin{aligned} w &= q + Mz + Ny, \quad w \geq 0, z \geq 0, y \geq 0 \\ z_1 w_1 &= 0, \quad i = 1, \dots, r+n \end{aligned} \right\} \quad (13)$$

where

$$w = \begin{bmatrix} \alpha \\ \beta \\ v_0 \end{bmatrix} \in R^{r+n+1}, \quad z = \begin{bmatrix} u \\ x \end{bmatrix} \in R^{r+n}, \quad q = \begin{bmatrix} -b \\ a \\ \lambda_k \end{bmatrix} \in R^{r+n+1}$$

$$M = \begin{bmatrix} 0 & A_1 \\ -A_1^T & 0 \\ 0 & -c^T \end{bmatrix} \in R^{(r+n+1) \times (r+n)}, \quad N = \begin{bmatrix} A_2 \\ 0 \\ -d^T \end{bmatrix} \in R^{(r+n+1) \times m}$$

Let  $z_0$  be an artificial variable and  $p$  be a nonnegative vector satisfying  $p_i > 0$  for all  $i$  such that  $q_i < 0$ . Consider the following program

$$\left. \begin{aligned} \text{Minimize } & z_0 \\ \text{subject to } & w = q + z_0 p + Mz + Ny \\ & z, w, y, z_0 \geq 0 \\ & z_1 w_1 = 0, \quad i = 1, \dots, r+n \end{aligned} \right\} \quad (14)$$

The BRES algorithm is an extension of the Phase 1 method with a single artificial variable [14, pages 136-138], and seeks the optimal solution of the program (14) by only using complementary feasible solutions (that is, solutions satisfying the constraints of (14)). To assure this condition, a nonbasic variable  $z_i$  (or  $w_i$ ) with a positive reduced cost coefficient can only be candidate to be an entering variable if its complementary  $w_i$  (or  $z_i$ ) is also nonbasic or becomes nonbasic in this iteration. Because of this additional criterion, the BRES algorithm may have three possible terminations:

TERM = 1 - an optimal solution  $(\bar{w}, \bar{z}, \bar{y})$  of (14) with  $\bar{z}_0 = 0$ .

TERM = 2 - an optimal solution  $(\bar{w}, \bar{z}, \bar{y})$  of (14) with  $\bar{z}_0 > 0$ .

TERM = 3 - a basic solution which is not optimal for (14) but there are no candidates for an entering variable.

In the first case (TERM = 1)  $(\bar{w}, \bar{z}, \bar{y})$  is a solution of the LCP (13). If TERM = 2 this LCP has no solution. In the latter case (TERM = 3) no conclusion can be made about the existence of a solution to the LCP(13). In appendix we present an example of a LCP of the form (13) for which there is a complementary solution but the BRES algorithm terminates with TERM = 3. This shows that the BRES method can not be used in general to solve the LCPs( $\lambda_k$ ) that the SLCP algorithm requires. However, the BRES method can be incorporated in a hybrid enumerative algorithm and this usually improves the efficiency of such procedure. This is discussed in the next section.

### 3 - A HYBRID ENUMERATIVE METHOD FOR THE LCP( $\lambda_k$ )


In this section we describe a hybrid enumerative method capable of solving any LCP( $\lambda_k$ ) required by the SLCP algorithm. Consider a LCP( $\lambda_k$ ) in the form

$$w = q + Mz + Ny \quad (15)$$

$$w \geq 0, z \geq 0, y \geq 0 \quad (16)$$

$$z_i w_i = 0, i = 1, \dots, r+n \quad (17)$$

where  $w \in R^{r+n+1}$ ,  $z \in R^{r+n}$  and  $y \in R^m$ . As in linear programming, a solution  $(z, w, y)$  satisfying the linear constraints (15) and (16) is called Feasible. A solution is Complementary if the variables  $z_i$  and  $w_i$  satisfy (17). An enumerative method attempts to find a complementary solution by only using basic feasible solutions of the system (15). To achieve this, the following tree is explored



where  $i_1, i_2, \dots$  are integer numbers of  $\{1, \dots, r+n\}$ . In node 1 an initial feasible solution is obtained by a variant of the Phase 1 algorithm with a single artificial variable, as is discussed later. Each one of the other nodes is generated by solving a subproblem, which consists of minimizing a variable  $z_i$  or  $w_i$  subject to the linear constraints of the LCP and some constraints  $z_j = 0$  or  $w_j = 0$ . For instance, to generate node 4 of the tree (18) it is necessary to solve the linear program

$$\left. \begin{array}{ll} \text{Minimize} & z_{i_2} \\ \text{subject to} & w = q + Mz + Ny, z \geq 0, w \geq 0, y \geq 0, \\ & z_{i_1} = 0 \end{array} \right\} \quad (19)$$

Such linear program is solved by a modification of the Phase 2 of the simplex method and two cases may occur:

- (i) if the variable minimized has value equal to zero, then it is fixed at zero in all descendant paths of the tree (the variable is said to be starred).
- (ii) if the minimum value of the variable is positive, then the branch is pruned and the node is fathomed.

The enumerative method attempts to solve the LCP by generating successive nodes of the tree according to the process explained above. The algorithm either finds a solution of the LCP (it is the first complementary feasible solution) or establishes that the LCP has no solutions (all the nodes of the tree are fathomed).

The enumerative method is efficient if there are only few nodes to be visited before a complementary solution is found or it is verified that none exists. There exist some heuristic rules and some procedures that usually improve the efficiency of the algorithm. The heuristic rules are related with the choice of the node of the tree and of the pair  $(z_{i_k}, w_{i_k})$  and are presented later. Next we describe the extensions for the LCP (13) of two efficient algorithms [10] that are usually incorporated in the enumerative method [10].

#### (i) A MODIFIED BRES ALGORITHM

Consider the problem of generating the node 1 of the tree (18), that is, finding a first feasible solution for the LCP (13). As stated before, this can be achieved by solving the linear program

$$\left. \begin{array}{l} \text{Minimize } z_0 \\ \text{subject to } w = q + z_0 p + Mz + Ny \\ z, w, y, z_0 \geq 0 \end{array} \right\} \quad (20)$$

where  $p$  is a nonnegative vector satisfying  $p_i > 0$  for all  $i$  such that  $q_i < 0$  and  $z_0$  is an artificial variable. Since our ultimate goal is to find a complementary feasible solution, then the BRES algorithm can be used to solve such linear program. If this algorithm terminates with  $\text{TERM} = 1$  or  $\text{TERM} = 2$ , then a solution of the LCP (13) has been found or LCP has no solutions respectively, and the enumerative method stops. If  $\text{TERM} = 3$ , the value of  $z_0$  can still be reduced if a simplex pivot step is performed, where the pivot belongs to a column with a positive reduced cost coefficient. The modified BRES algorithm consists of performing such pivot step when  $\text{TERM} = 3$  and reapplying the BRES method by starting with the basic solution obtained by this pivot step. The process is then repeated. This algorithm can also be used to generate any other node  $k$ , since, as discussed before, the generation of a node consists of minimizing a variable  $z_i$  or  $w_i$ .

To describe the modified BRES algorithm, we associate to each basic solution a quantity NCP defined as the number of pairs of complementary variables  $(z_i, w_i)$  such that  $z_i$  and  $w_i$  are both basic in this solution. If  $\text{NCP} = 0$  the basic solution is complementary. The converse is true for nondegenerate basic solutions, but a degenerate basic solution may be complementary with  $\text{NCP} > 0$ . Hence the value of NCP is increased by one whenever  $\text{TERM} = 3$  occurs in the BRES algorithm. In addition, NCP never increases and may even be reduced during the BRES procedure. Hence a complementary feasible solution may occur during this algorithm. The steps of the modified BRES algorithm are presented below.

#### A Modified BRES Algorithm for Generating a node $k$

STEP 0 - Let NCP be the number of pairs of complementary variables  $(z_i, w_i)$  such that  $z_i$  and  $w_i$  are both basic in the basic solution ( $\text{NCP} = 0$  if  $k = 1$ ).

STEP 1 - Apply the BRES algorithm. If a complementary feasible solution is found when the BRES algorithm is in use, set  $\text{TERM} = 1$  and  $\text{NCP} = 0$  and go to Exit. If the BRES algorithm terminates with  $\text{TERM} = 1$  or  $\text{TERM} = 2$ , go to Exit. Otherwise ( $\text{TERM} = 3$ ) go to step 2.

STEP 2 - Set  $\text{NCP} = \text{NCP} + 1$  and perform a simplex pivot step, where the pivot belongs to a column with a positive reduced cost coefficient. Go to step 1.

EXIT - If  $\text{TERM} = 1$  and  $\text{NCP} = 0$ , a solution of the LCP has been found and the enumerative method stops. If  $\text{TERM} = 1$  and  $\text{NCP} > 0$ , the node  $k$  has been generated. If  $\text{TERM} = 2$ , the node  $k$  can not be generated and is fathomed.

#### (II) A MODIFIED REDUCED-GRADIENT (MRG) ALGORITHM

This algorithm is described in [1] and consists of finding a local star minimum of the function

$$f(z, w, y) = \sum_{i=1}^{r+n} z_i w_i \quad (21)$$

that is, a basic feasible solution  $(\bar{z}, \bar{w}, \bar{y})$  such that

$$f(\bar{z}, \bar{w}, \bar{y}) \leq f(z, w, y) \quad (22)$$

for all its adjacent basic feasible solutions  $(z, w, y)$ . To describe the algorithm for the case of the LCP(13), consider a basic feasible solution  $(\bar{z}, \bar{w}, \bar{y})$  with Basis  $B$ . If  $(\bar{z}, \bar{w}, \bar{y})$  is an adjacent basic solution of  $(\bar{z}, \bar{w}, \bar{y})$ , then

$$(\bar{z}, \bar{w}, \bar{y}) = (\bar{z}, \bar{w}, \bar{y}) + \mu_0 (d^z, d^w, d^y) \quad (23)$$

where  $\mu_0$  is the minimum ratio of the simplex method and  $d = (d^z, d^w, d^y)$  is a feasible direction such that  $d^z, d^w$  and  $d^y$  are the vectors of the components of  $d$  corresponding to the variables  $z, w$  and  $y$  respectively. To define this direction, let  $F$  and  $T$  be the sets of the indices of the basic and nonbasic variables respectively and  $s$  be the index of the nonbasic entering variable that is increased to generate the basic feasible solution  $(\bar{z}, \bar{w}, \bar{y})$ . Then

$$\begin{aligned} d_s &= 1 \\ d_j &= 0 \quad \text{for all } j \in T - \{s\} \end{aligned}$$



$$d_j = \begin{cases} -B^{-1}M_s & \text{if } s \text{ is a column of a } z \text{ variable} \\ B^{-1}I_s & \text{if } s \text{ is a column of a } w \text{ variable} \\ -B^{-1}N_s & \text{if } s \text{ is a column of a } y \text{ variable} \end{cases}$$

where  $M_s$ ,  $N_s$  and  $I_s$  represent the column  $s$  of the matrices  $M$ ,  $N$  and identity matrix  $I$  respectively. By the structures of the matrices  $M$  and  $N$  of the LCP(13), we can write

$$P(I - M - N)Q = \begin{bmatrix} E & 0 \\ 0 & G \end{bmatrix} \quad (24)$$

where  $P$  and  $Q$  are permutation matrices and  $E = R(r-1) \times (r+n-m+1)$ ,  $G = R^n \times (r-n)$  are matrices such that  $E$  contains a column of the variable  $z_i(w_i)$  if and only if  $G$  contains a column of the variable  $w_i(z_i)$ . Hence

$$PBQ = \begin{bmatrix} B_1 & 0 \\ 0 & B_2 \end{bmatrix} \quad (25)$$

where  $B_1 \in R^{(r-1) \times (r-1)}$  and  $B_2 \in R^{n \times n}$  are nonsingular matrices, and this implies that

$$d = \begin{bmatrix} 0 \\ 1 \\ B_2^{-1}G_s \end{bmatrix} \quad \text{or} \quad d = \begin{bmatrix} 1 \\ B_1^{-1}E_s \\ 0 \end{bmatrix} \quad (26)$$

Therefore

$$d_i^w = 0 \quad \text{or} \quad d_i^z = 0 \quad \text{for each } i = 1, \dots, r+n \quad (27)$$

Computing the value of the function (21) at the adjacent basic feasible solution given by (23), we have

$$f(\bar{z}, \bar{w}, \bar{y}) = \sum_{i=1}^{r+n} \bar{z}_i \bar{w}_i = \sum_{i=1}^{r+n} (\bar{z}_i + \mu_0 d_i^z) (\bar{w}_i + \mu_0 d_i^w)$$

$$= \sum_{i=1}^{r+n} \bar{z}_i \bar{w}_i + \mu_0 \sum_{i=1}^{r+n} (\bar{z}_i d_i^w + \bar{w}_i d_i^z) + \mu_0^2 \sum_{i=1}^{r+n} d_i^z d_i^w$$

Hence by (27) we have

$$f(\bar{z}, \bar{w}, \bar{y}) - f(\bar{z}, \bar{w}, \bar{y}) = \mu_0 \sum_{i=1}^{r+n} (\bar{z}_i d_i^w + \bar{w}_i d_i^z) \quad (28)$$

But

$$\sum_{i=1}^{r+n} (\bar{z}_i d_i^w + \bar{w}_i d_i^z) = \sum_{i=1}^{r+n} (\bar{w}_i z_i + \bar{z}_i w_i) \quad (29)$$

is exactly the symmetric of the reduced-cost coefficient  $\bar{\sigma}_j$  of the nonbasic variable of index  $s$  associated with the linear function

$$\sum_{i=1}^{r+n} (\bar{w}_i z_i + \bar{z}_i w_i) \quad (29)$$

Therefore we have shown that for the LCP( $\lambda_0$ ), Al-Khayyal's MRG Algorithm reduces to a simplex-type method in which the reduced-cost coefficients  $\bar{\sigma}_j$  of the nonbasic variables are associated with the linear function (29). The algorithm also incorporates a modification of Bland's rule [8] to avoid the occurrence of cycling and is presented below.

#### MRG Algorithm

- STEP 0 - Let  $(\bar{z}, \bar{w}, \bar{y})$  be a basic feasible solution with Basis  $B$ .  
Set  $\text{deg} = 0$ .
- STEP 1 - Compute the reduced-cost coefficients  $\bar{\sigma}_j$  associated with the linear function (29). If  $\bar{\sigma}_j \leq 0$  for all  $j$ , then  $(\bar{z}, \bar{w}, \bar{y})$  is a local star minimum and stop. Otherwise go to step 2.
- STEP 2 - Find the entering variable as the nonbasic variable in the column  $s$  given by

$$s = \begin{cases} \min \{j : \bar{\sigma}_j = \max \bar{\sigma}_j\} & \text{if } \text{deg} = 0 \\ \min \{j : \bar{\sigma}_j > 0\} & \text{if } \text{deg} = 1 \end{cases} \quad (30)$$

STEP 3 - Compute the minimum ratio  $\mu_0$  for this entering variable and let the leaving variable be the first basic variable at which  $\mu_0$  is attained. If  $\mu_0 = 0$  ( $\mu_0 > 0$ ) set  $\text{deg} = 1$  ( $\text{deg} = 0$ ). Perform a simplex pivot step and obtain a basic feasible solution  $(\bar{z}, \bar{w}, \bar{y})$  with Basis B. Go to step 1.

As described above, the MRG algorithm always terminates in a local star minimum  $(\bar{z}, \bar{w}, \bar{y})$  of the function  $f(z, w, y)$  given by (21). The method is quite suitable to be incorporated in a hybrid enumerative method, since the value of the complementarity function  $\sum_{i=1}^{r+n} z_i w_i$  is usually reduced and may take the value zero, in which case a solution of the LCP(13) is found.

The hybrid enumerative method, incorporating the two procedures referred in this section and the most efficient heuristic rules, is stated below.

#### Hybrid Enumerative Method for Solving LCP(13)

STEP 0 - Set  $\text{NODE} = \text{NNODE} = 1$ , where  $\text{NNODE}$  is the total number of nodes to be investigated and  $\text{NODE}$  is the current node. Set  $\text{STZ}(\text{NODE}) = \text{STW}(\text{NODE}) = \emptyset$ , where  $\text{STZ}(\text{NODE})$  and  $\text{STW}(\text{NODE})$  are the sets of the indices of the  $z$  and  $w$  starred variables at node  $\text{NODE}$  respectively.

STEP 1 - Generation of node 1 (First feasible solution).

(i) Apply the modified BRES algorithm for solving the linear program (20). If  $\text{TERM} = 1$  and  $\text{NCP} = 0$ , go to Exit. If  $\text{TERM} = 2$  go to Exit. Otherwise ( $\text{TERM} = 1$  and  $\text{NCP} > 0$ ) go to (ii).

(ii) Apply the MRG algorithm to find a local star minimum  $(\bar{z}, \bar{w}, \bar{y})$  of the function  $f(z, w, y)$  given by (21). If  $f(\bar{z}, \bar{w}, \bar{y}) = 0$ , set  $\text{TERM} = 1$  and go to Exit. Otherwise set  $\text{NCP}(\text{NODE}) = \text{NCP}$  and go to step 2.

STEP 2 - Branching - Set  $\text{NCP} = \text{NCP}(\text{NODE})$ ,  $\text{STZ} = \text{STZ}(\text{NODE})$  and  $\text{STW} = \text{STW}(\text{NODE})$ . Choose the pair of complementary variables  $(z_s, w_s)$  in node  $\text{NODE}$  such that

$$\bar{z}_s \bar{w}_s = \max \{ \bar{z}_i \bar{w}_i : i = 1, \dots, \text{NCP} \} \quad (31)$$

Fathom node  $\text{NODE}$  and go to step 3.

STEP 3 - Generate  $\text{NNODE} + 1$  - Set  $\text{NODE} = \text{NNODE} = \text{NNODE} + 1$ .

(i) Apply the modified BRES algorithm for solving the linear program

$$\left. \begin{array}{l} \text{Minimize } z_s \\ \text{subject to } w = q + Mz + Ny, z, w, y \geq 0 \\ z_j = 0, w_t = 0, j \in \text{STZ}, t \in \text{STW} \end{array} \right\} \quad (32)$$

If  $\text{TERM} = 1$  and  $\text{NCP} = 0$ , go to Exit. If  $\text{TERM} = 2$ , fathom node  $\text{NODE}$  and go to step 4. Otherwise set  $\text{STZ}(\text{NODE}) = \text{STZ} \cup \{s\}$  and go to (ii).

(ii) Apply the MRG algorithm to find a local star minimum  $(\bar{z}, \bar{w}, \bar{y})$  of the function  $f(z, w, y)$  given by (21). If  $f(\bar{z}, \bar{w}, \bar{y}) = 0$ , set  $\text{TERM} = 1$  and go to Exit. Otherwise set  $\text{NCP}(\text{NODE}) = \text{NCP}$ ,  $R(\text{NODE}) = \sum_{i=1}^{\text{NCP}} \bar{w}_i \bar{z}_i$  and go to step 4.

STEP 4 - Generate node  $\text{NNODE} + 2$  - This is step 3 with  $z_s$ ,  $\text{STZ}$  and step 4 replaced by  $w_s$ ,  $\text{STW}$  and step 5, respectively.

STEP 5 - Choice of the node - If all the nodes  $1, \dots, \text{NNODE}$  are fathomed, set  $\text{TERM} = 2$  and go to Exit. Otherwise choose the node  $\text{NODE}$  by

$$\text{NCP}(\text{NODE}) + \frac{R(\text{NODE})}{10} = \min \left\{ \text{NCP}(j) + \frac{R(j)}{10} : j = 1, \dots, \text{NNODE} \text{ and } j \text{ is unfathomed} \right\} \quad (33)$$

and go to step 2.

EXIT - If  $\text{TERM} = 1$ , the complementary basic feasible solution  $(\bar{z}, \bar{w}, \bar{y})$  is a solution of the LCP(13). If  $\text{TERM} = 2$ , the LCP(13) has no solution.

We note that (31) and (33) are heuristic rules that we have found efficient to reduce the computational effort of the algorithm [10]. The hybrid enumerative method can be implemented for solving large and sparse LCPs. The implementation uses reinversion and updating techniques for the LU decomposition of the Basis matrices [16] and special data structures for the representation of the sets discussed above. We suggest [10] for a description of this procedure.

As discussed in section 2, the hybrid enumerative method is used to solve the



LCPs ( $\lambda_k$ ) required by the SLCP algorithm. For any two values  $\lambda_k < \lambda_{k-1}$ , the LCP( $\lambda_{k-1}$ ) and LCP( $\lambda_k$ ) only differ in the  $r$ -th component of the vector  $q$ . Furthermore, if  $B$  is the Basis associated to the solution  $(\bar{x}, \bar{w}, \bar{y})$  of the LCP( $\lambda_{k-1}$ ) obtained by the hybrid enumerative method and  $q$  is the right-hand side vector of the LCP( $\lambda_k$ ), then the vector  $\bar{q} = B^{-1}q$  satisfies

$$\bar{q}_j \geq 0 \text{ for all } j = 1, \dots, r+n \text{ and } \bar{q}_{r+n+1} < 0 \quad (34)$$

Hence the solution  $(\bar{x}, \bar{w}, \bar{y})$  of the LCP( $\lambda_{k-1}$ ) can be used as the initial basic solution for the LCP( $\lambda_k$ ). Since the vector  $B^{-1}q$  satisfies (34), then only the last component of the vector  $B^{-1}p$  has to be positive. Hence we can choose in (20)

$$p = B \begin{bmatrix} 0 \\ 1 \end{bmatrix} \quad (35)$$

where  $0 \in \mathbb{R}^{r+n}$  is the null vector. The choice of this initial basic solution usually provides great reductions in the hybrid enumerative method. In fact, the algorithm requires quite often a unique node and a quite small number of pivot steps to find the solution of the LCP( $\lambda_k$ ). This is shown by the computational results presented in the next section.

In the original SLCP algorithm the LCP( $\lambda_0$ ) is solved in step 0, where  $\lambda_0$  is an upper-bound of  $c^T x + d^T y$  on  $H$ . The computation of this value of  $\lambda_0$  may be expensive in certain cases. Next, we present a modified step 0 of the SLCP algorithm, which does not require the computation of this upper-bound.

**Modified Step 0** - Set  $k = 0$  and solve the LCP(0) with  $v_0$  unrestricted in sign. If this LCP has no solution go to Exit. Otherwise let  $(x^0, y^0)$  be the solution of this LCP and  $\lambda_0 = c^T x^0 + d^T y^0$  be the value of the variable  $v_0$ . Compute  $\lambda_1$  by (12).

Since in this step the variable  $v_0$  is unrestricted in sign and is originally basic, then it remains basic during the solution of the LCP(0). Hence this LCP is exactly the LCP( $\lambda_0$ ) without the constraint  $c^T x + d^T y \leq \lambda_0$ . Since the value of  $\lambda_0$  obtained by the Modified Step 0 is usually smaller than the upper-bound chosen by the original SLCP algorithm, then this modification reduces the number of LCPs to be solved and improves the efficiency of the SLCP algorithm. This is another important reason for the use of the Modified Step 0.

#### 4. COMPUTATIONAL EXPERIENCE

In this section we present some computational experience with the SLCP

algorithm described in this paper on the solution of some medium-scale sparse LBLPs. The matrix  $A = [A_1, A_2]$  of the test problems has been constructed by an extension for sparse matrices of the technique described in [15]. This modification consists of introducing a quantity NZ, representing the number of nonzero elements of the matrix  $A$ , and randomly distributing these nonzero elements in a scheme similar to [15]. We have also considered in the matrix  $A$  a dense row of ones to make bounded the set  $H$  defined by (5). We associate to each matrix  $A$  two test problems, which differ in the cost coefficients. In the problems denoted by TPN, all the cost coefficients  $c, d$ , and  $a$  of the two linear functions are nonnegative. In the problems denoted by TPQ, the cost coefficients  $c$  and  $d$  of the function of the first level are nonnegative but the linear function of the second level has some negative cost coefficients. By doing this we simulate LBLPs with conflicting (problems TPQ) and nonconflicting (problems TPN) objectives.

The characteristics of the test problems and the results of the experiences of solving these problems by the SLCP algorithm are presented in Table 1. All the tests have been performed on a CDC CYBER 180-830 of the University of Porto. In these tables there are some parameters with the following meanings:

$n$  = number of second level variables  $x$

$m$  = number of first level variables  $y$

$r$  = number of constraints = number of rows of  $A = [A_1, A_2]$

$rsm$  = relative sparsity of  $A = [A_1, A_2] = \frac{\text{number of nonzeros of } A}{n+m}$

$nrow$  = number of rows of the LCP( $\lambda_k$ ) =  $r + n + 1$

$ncol$  = number of columns of the LCP( $\lambda_k$ ) =  $r + n + m$

NLCP = number of LCPs( $\lambda_k$ ) to be solved by the SLCP algorithm.

NI = Total number of simplex pivot steps required by the SLCP algorithm.

NIMIN = minimum number of pivot steps that a LCP( $\lambda_k$ ) has required.

NIMAX = maximum number of pivot steps that a LCP( $\lambda_k$ ) with a solution has required.

T = CPU time in seconds for the SLCP algorithm.

The results presented in Table 1 show that it is worthwhile to solve medium-scale LBLPs by the SLCP algorithm discussed in this paper. As expected the efficiency of the



A complementary basic feasible solution for the program (14) is obtained by performing the simplex pivot step whose pivot is circled in the tableau. The tableau associated with this basic solution is given below.

	$v_1$	$z_1$	$z_2$	$z_3$	$z_4$	$z_5$	$z_6$	$y_1$	
$z_0$	-1	0	0	0	0	0	2	1	10
$v_2$	0	0	0	0	0	0	-2	①	6
$v_3$	0	0	0	0	0	0	-1	2	21
$v_4$	0	0	0	0	0	0	2	1	38
$v_5$	0	0	0	0	0	0	2	-1	18
$v_6$	0	2	2	1	-2	-2	0	0	1
$v_7$	0	0	0	0	0	0	-1	0	0

Since we wish to minimize the value of the variable  $z_0$ , then the variables  $z_6$  and  $y_1$  are eligible to be the entering variable. However  $z_6$  can not be a candidate, since  $w_6$  is basic and does not become nonbasic if  $z_6$  is chosen as entering variable. Hence  $y_1$  must be the entering variable. After finding the pivot (entry circled in the tableau) by the usual minimum ratio criterion and performing the corresponding pivot step, we obtain another complementary basic feasible solution given by the following tableau

	$v_1$	$z_1$	$z_2$	$z_3$	$z_4$	$z_5$	$z_6$	$v_2$	
$z_0$	-1	0	0	0	0	0	4	-1	4
$y_1$	0	0	0	0	0	0	-2	1	6
$v_3$	0	0	0	0	0	0	3	-2	9
$v_4$	0	0	0	0	0	0	4	-1	32
$v_5$	0	0	0	0	0	0	0	1	24
$v_6$	0	2	2	1	-2	-2	0	0	1
$v_7$	0	0	0	0	0	0	-1	0	0

In this iteration only  $z_6$  is eligible to be the entering variable. As before this variable can not be a candidate, whence the termination TERM = 3 occurs. However, the LCP(0) has at least the solution  $x = \left(\frac{1}{2}, 0, 0, 0, 1\right)$ ,  $y_1 = 8$ .

## REFERENCES

- [1] F.A. Al-Khayyal, An implicit enumeration procedure for the general linear complementarity problem, *Mathematical Programming Studies* 81 (1987) 1-20.
- [2] J.F. Bard, An efficient point algorithm for a linear two-stage optimization problem, *Operations Research* 81 (1988) 670-684.
- [3] J.F. Bard, Convex two-level optimization, *Mathematical Programming* 40 (1988) 15-26.
- [4] J.F. Bard and J.E. Falk, An explicit solution to the multi-level programming problems, *Computers and Operations Research* 9 (1982) 77-100.
- [5] J.F. Bard and J.J. Moore, A branch-and-bound algorithm for the Bilevel linear programming problem, Technical Report, Department of Mechanical Engineering, University of Texas at Austin, 1988.
- [6] O. Ben-Ayed, C.E. Blair and D.E. Boyce, Solving a real world highway network design problem using bilevel linear programming, *IEBR Faculty Working Paper* 1463, University of Illinois at Urbana-Champaign, 1988.
- [7] W.F. Blais and M.H. Karwan, Two-level linear programming, *Management Science* 30 (1984) 1004-1020.
- [8] R.G. Bland, New finite pivoting rules for the simplex method, *Mathematics of Operations Research* 2 (1977) 103-107.
- [9] W. Candler and R. Townsley, A linear two-level programming problem, *Computers and Operations Research* 9 (1982) 59-78.
- [10] J.J. Júdice and A.M. Feustino, An experimental investigation of enumerative methods for the linear complementarity problem, *Computers and Operations Research* 15 (1988) 417-428.
- [11] C.D. Kolstad, A review of the literature on Bilevel Mathematical Programming, Technical Report LA-10284-MS, Los Alamos National Laboratory, 1985.
- [12] C.D. Kolstad and L.S. Lasdon, Derivative evaluation and computational experience with large Bilevel mathematical programs, Faculty Working Paper 1268, College of Commerce and Business Administration, University of Illinois at Urbana-Champaign, 1986.
- [13] H. M. Markowitz, The optimization of a quadratic function subject to linear constraints, *Naval Research Logistics Quarterly* 8 (1956) 111-183.
- [14] K.G. Murty, *Linear and Combinatorial Programming*, Wiley, 1978.
- [15] B. Ramareo and C.M. Shetty, Application of disjunctive programming to the linear complementarity problem, *Naval Research Logistics Quarterly* 81 (1984) 589-600.
- [16] J.K. Reid, A sparsity-exploitation variant of the Bartels-Golub decomposition for linear programming bases, *Mathematical Programming* 24 (1982) 55-69.
- [17] P. Wolfe, The simplex method for quadratic programming, *Econometrica* 27 (1959) 382-398.