Cell Tuition
1-1 PRIVATE TUTORS

ALGEBRA

The Quadratic Formula

Name _____ Date _____ Calculator allowed Version 1

Scan to practise this topic online

KEY WORDS quadratic formula · discriminant · coefficient *(tau whakarea)* · decimal places

A Naming a, b and c ●○○

Remember: Write the equation as $ax^2 + bx + c = 0$ first. Keep the sign with each number.

1. Write down the values of a, b and c.

a) $x^2 + 5x + 6 = 0$ _____

b) $2x^2 - 3x - 7 = 0$ _____

c) $x^2 - 9 = 0$ _____

d) $x^2 - 4x = 0$ _____

e) $5 - 2x - x^2 = 0$ _____

f) $3x^2 = 4x + 1$ _____

g) $x^2 + x = 12$ _____

h) $2x^2 = 5$ _____

B The discriminant ●●○

Remember: The discriminant is $b^2 - 4ac$. It is the part of the formula under the square root.

2. Work out $b^2 - 4ac$ for each equation.

a) $x^2 + 3x + 1 = 0$

b) $2x^2 - 3x - 5 = 0$

c) $x^2 - 6x + 9 = 0$

d) $3x^2 + 2x + 4 = 0$

e) $x^2 - 7 = 0$

f) $5x^2 - 4x - 2 = 0$

g) $-x^2 + 5x + 3 = 0$

h) $6x^2 + 7x + 2 = 0$

i) $2x^2 - x + 10 = 0$

C Solving with the formula ●●○

Remember: For $ax^2 + bx + c = 0$: $x = \dfrac{-b \pm \sqrt{b^2 - 4ac}}{2a}$

3. Solve with the quadratic formula. These have exact answers.

a) $x^2 + 5x + 6 = 0$

b) $x^2 - 7x + 10 = 0$

c) $2x^2 + 5x + 2 = 0$

d) $x^2 + 2x - 8 = 0$

e) $2x^2 - 7x + 3 = 0$

f) $4x^2 + 4x + 1 = 0$

g) $x^2 - x - 12 = 0$

h) $4x^2 - 9 = 0$

i) $5x^2 + 9x - 2 = 0$

© 2026 Cell Tuition. You may print and photocopy this worksheet for your own classes and children. It may not be changed, sold or posted on other websites.

D Answers to 2 decimal places ●●●

⚠ **Watch out:** Take care with negatives: if $b = -5$ then $-b = 5$ and $b^2 = 25$.

4. Solve with the quadratic formula. Give your answers to 2 decimal places.

a) $x^2 + 4x + 1 = 0$ **b)** $x^2 - 5x + 3 = 0$ **c)** $x^2 + 6x - 2 = 0$

d) $2x^2 + 3x - 4 = 0$ **e)** $3x^2 - 7x + 1 = 0$ **f)** $x^2 - 2x - 5 = 0$

g) $2x^2 - 8x + 3 = 0$ **h)** $5x^2 + 2x - 1 = 0$ **i)** $-x^2 + 4x + 3 = 0$

E How many solutions? ●●●

💡 **Remember:** Discriminant positive: two solutions. Zero: one solution. Negative: no solutions.

5. Use the discriminant to decide how many solutions each equation has.

a) $x^2 + 4x + 5 = 0$ **b)** $x^2 - 10x + 25 = 0$ **c)** $2x^2 + 7x + 1 = 0$


d) $3x^2 - 2x + 2 = 0$ **e)** $x^2 + x - 1 = 0$ **f)** $9x^2 + 6x + 1 = 0$

g) $2x^2 - 5x + 4 = 0$ **h)** $x^2 + 3 = 0$ **i)** $5x^2 + 3x - 2 = 0$

F Problems ●●●

6. A ball is thrown into the air. Its height h metres after t seconds is $h = -5t^2 + 12t + 2$. The ball lands when $h = 0$. After how many seconds does it land? Give your answer to 2 decimal places.

7. The rectangle has an area of 30 m². Write an equation and solve it to find x to 2 decimal places.



8. The square of a number plus the number itself equals 10. Find the two possible numbers, to 2 decimal places.

★ Extension

9. Solve $x^2 + 6x + 4 = 0$. Give your answers exactly, in the form $p \pm \sqrt{q}$.

10. The equation $x^2 + 6x + k = 0$ has exactly one solution. Find the value of k.

11. Solve $x(x + 3) = 5$. Give your answers to 2 decimal places.

© 2026 Cell Tuition. You may print and photocopy this worksheet for your own classes and children. It may not be changed, sold or posted on other websites.

The Quadratic Formula

Name _____ Date _____ Calculator allowed Version 2

Scan to practise this topic online

KEY WORDS quadratic formula · discriminant · coefficient *(tau whakarea)* · decimal places

A Naming a, b and c ●○○

Remember: Write the equation as $ax^2 + bx + c = 0$ first. Keep the sign with each number.

1. Write down the values of a, b and c.

a) $x^2 + 7x + 10 = 0$ _____

b) $3x^2 - 2x - 8 = 0$ _____

c) $x^2 - 16 = 0$ _____

d) $x^2 - 6x = 0$ _____

e) $7 - 3x - x^2 = 0$ _____

f) $2x^2 = 5x + 3$ _____

g) $x^2 + 2x = 15$ _____

h) $3x^2 = 7$ _____

B The discriminant ●●○

Remember: The discriminant is $b^2 - 4ac$. It is the part of the formula under the square root.

2. Work out $b^2 - 4ac$ for each equation.

a) $x^2 + 5x + 2 = 0$

b) $2x^2 - 5x - 3 = 0$

c) $x^2 - 8x + 16 = 0$

d) $2x^2 + 3x + 5 = 0$

e) $x^2 - 11 = 0$

f) $4x^2 - 6x - 1 = 0$

g) $-x^2 + 3x + 6 = 0$

h) $6x^2 + 11x + 3 = 0$

i) $3x^2 - x + 8 = 0$

C Solving with the formula ●●○

Remember: For $ax^2 + bx + c = 0$: $x = \dfrac{-b \pm \sqrt{b^2 - 4ac}}{2a}$

3. Solve with the quadratic formula. These have exact answers.

a) $x^2 + 7x + 12 = 0$

b) $x^2 - 8x + 15 = 0$

c) $2x^2 + 7x + 3 = 0$

d) $x^2 + 3x - 10 = 0$

e) $2x^2 - 9x + 4 = 0$

f) $4x^2 - 12x + 9 = 0$

g) $x^2 - x - 20 = 0$

h) $4x^2 - 25 = 0$

i) $5x^2 + 8x - 4 = 0$

© 2026 Cell Tuition. You may print and photocopy this worksheet for your own classes and children. It may not be changed, sold or posted on other websites.

D Answers to 2 decimal places ●●●

⚠ **Watch out:** Take care with negatives: if $b = -5$ then $-b = 5$ and $b^2 = 25$.

4. Solve with the quadratic formula. Give your answers to 2 decimal places.

a) $x^2 + 6x + 2 = 0$ **b)** $x^2 - 7x + 5 = 0$ **c)** $x^2 + 4x - 3 = 0$

d) $2x^2 + 5x - 1 = 0$ **e)** $3x^2 - 8x + 2 = 0$ **f)** $x^2 - 4x - 6 = 0$

g) $2x^2 - 6x + 1 = 0$ **h)** $5x^2 + 3x - 1 = 0$ **i)** $-x^2 + 6x + 2 = 0$

E How many solutions? ●●●

Remember: Discriminant positive: two solutions. Zero: one solution. Negative: no solutions.

5. Use the discriminant to decide how many solutions each equation has.

a) $x^2 + 2x + 6 = 0$ **b)** $x^2 - 12x + 36 = 0$ **c)** $2x^2 + 9x + 3 = 0$


d) $4x^2 - 3x + 1 = 0$ **e)** $x^2 + 3x - 2 = 0$ **f)** $4x^2 + 20x + 25 = 0$

g) $3x^2 - 4x + 2 = 0$ **h)** $x^2 + 5 = 0$ **i)** $5x^2 + 2x - 3 = 0$

F Problems ●●●

6. A ball is thrown into the air. Its height h metres after t seconds is $h = -5t^2 + 15t + 4$. The ball lands when $h = 0$. After how many seconds does it land? Give your answer to 2 decimal places.

7. The rectangle has an area of 20 m². Write an equation and solve it to find x to 2 decimal places.



8. The square of a number minus 3 times the number equals 5. Find the two possible numbers, to 2 decimal places.

★ Extension

9. Solve $x^2 + 8x + 9 = 0$. Give your answers exactly, in the form $p \pm \sqrt{q}$.

10. The equation $x^2 - 10x + k = 0$ has exactly one solution. Find the value of k.

11. Solve $x(x - 2) = 7$. Give your answers to 2 decimal places.

© 2026 Cell Tuition. You may print and photocopy this worksheet for your own classes and children. It may not be changed, sold or posted on other websites.