

# Discrete Mathematics

## Relations and Functions

H. Turgut Uyar Ayşegül Gençata Yayımlı Emre Harmancı

2001-2013

1 / 79

## License



©2001-2013 T. Uyar, A. Yayımlı, E. Harmancı

You are free:

- ▶ to Share – to copy, distribute and transmit the work
- ▶ to Remix – to adapt the work

Under the following conditions:

- ▶ Attribution – You must attribute the work in the manner specified by the author or licensor (but not in any way that suggests that they endorse you or your use of the work).
- ▶ Noncommercial – You may not use this work for commercial purposes.
- ▶ Share Alike – If you alter, transform, or build upon this work, you may distribute the resulting work only under the same or similar license to this one.

Legal code (the full license):

<http://creativecommons.org/licenses/by-nc-sa/3.0/>

2 / 79

## Topics

### Relations

Introduction  
Relation Properties  
Equivalence Relations

### Functions

Introduction  
Pigeonhole Principle  
Recursion

3 / 79

## Relation

### Definition

**relation:**  $\alpha \subseteq A \times B \times C \times \dots \times N$

- ▶ **tuple:** an element of a relation
- ▶  $\alpha \subseteq A \times B$ : *binary relation*
  - ▶  $a\alpha b$  is the same as  $(a, b) \in \alpha$
- ▶ representations:
  - ▶ by drawing
  - ▶ by matrix


4 / 79

## Relation Example

### Example

$A = \{a_1, a_2, a_3, a_4\}, B = \{b_1, b_2, b_3\}$

$\alpha = \{(a_1, b_1), (a_1, b_3), (a_2, b_2), (a_2, b_3), (a_3, b_1), (a_3, b_3), (a_4, b_1)\}$



	$b_1$	$b_2$	$b_3$
$a_1$	1	0	1
$a_2$	0	1	1
$a_3$	1	0	1
$a_4$	1	0	0

$$M_\alpha = \begin{bmatrix} 1 & 0 & 1 \\ 0 & 1 & 1 \\ 1 & 0 & 1 \\ 1 & 0 & 0 \end{bmatrix}$$

5 / 79

## Relation Composition

### Definition

**relation composition:**

let  $\alpha \subseteq A \times B, \beta \subseteq B \times C$


$\alpha\beta = \{(a, c) \mid a \in A, c \in C, \exists b \in B [a\alpha b \wedge b\beta c]\}$

- ▶  $M_{\alpha\beta} = M_\alpha \times M_\beta$ 
  - ▶ using logical operations:  
 $1 : T, 0 : F, \cdot : \wedge, + : \vee$

6 / 79

## Relation Composition Example

Example



7 / 79

## Relation Composition Matrix Example

Example

$$M_{\alpha} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 0 & 1 \end{bmatrix} \quad M_{\beta} = \begin{bmatrix} 1 & 1 & 0 & 0 \\ 0 & 0 & 1 & 1 \\ 0 & 1 & 1 & 0 \end{bmatrix} \quad M_{\alpha\beta} = \begin{bmatrix} 1 & 1 & 0 & 0 \\ 0 & 1 & 1 & 0 \\ 0 & 0 & 1 & 1 \\ 1 & 1 & 1 & 0 \end{bmatrix}$$

8 / 79

## Relation Composition Associativity

► relation composition is associative

$$(\alpha\beta)\gamma = \alpha(\beta\gamma).$$

$$\begin{aligned} & (a, d) \in (\alpha\beta)\gamma \\ \Leftrightarrow & \exists c [(a, c) \in \alpha\beta \wedge (c, d) \in \gamma] \\ \Leftrightarrow & \exists c [\exists b [(a, b) \in \alpha \wedge (b, c) \in \beta] \wedge (c, d) \in \gamma] \\ \Leftrightarrow & \exists b [(a, b) \in \alpha \wedge \exists c [(b, c) \in \beta \wedge (c, d) \in \gamma]] \\ \Leftrightarrow & \exists b [(a, b) \in \alpha \wedge (b, d) \in \beta\gamma] \\ \Leftrightarrow & (a, d) \in \alpha(\beta\gamma) \end{aligned}$$

□

9 / 79

## Relation Composition Theorems

- let  $\alpha, \delta \subseteq A \times B$ , and  
let  $\beta, \gamma \subseteq B \times C$
- $\alpha(\beta \cup \gamma) = \alpha\beta \cup \alpha\gamma$
- $\alpha(\beta \cap \gamma) \subseteq \alpha\beta \cap \alpha\gamma$
- $(\alpha \cup \delta)\beta = \alpha\beta \cup \delta\beta$
- $(\alpha \cap \delta)\beta \subseteq \alpha\beta \cap \delta\beta$
- $(\alpha \subseteq \delta \wedge \beta \subseteq \gamma) \Rightarrow \alpha\beta \subseteq \delta\gamma$

10 / 79

## Relation Composition Theorems

$$\alpha(\beta \cup \gamma) = \alpha\beta \cup \alpha\gamma.$$

$$\begin{aligned} & (a, c) \in \alpha(\beta \cup \gamma) \\ \Leftrightarrow & \exists b [(a, b) \in \alpha \wedge (b, c) \in (\beta \cup \gamma)] \\ \Leftrightarrow & \exists b [(a, b) \in \alpha \wedge ((b, c) \in \beta \vee (b, c) \in \gamma)] \\ \Leftrightarrow & \exists b [((a, b) \in \alpha \wedge (b, c) \in \beta) \\ & \vee ((a, b) \in \alpha \wedge (b, c) \in \gamma)] \\ \Leftrightarrow & (a, c) \in \alpha\beta \vee (a, c) \in \alpha\gamma \\ \Leftrightarrow & (a, c) \in \alpha\beta \cup \alpha\gamma \end{aligned}$$

□

11 / 79

## Converse Relation

Definition

$$\alpha^{-1} = \{(b, a) \mid (a, b) \in \alpha\}$$

$$\text{► } M_{\alpha^{-1}} = M_{\alpha}^T$$

12 / 79

## Converse Relation Theorems

- ▶  $(\alpha^{-1})^{-1} = \alpha$
- ▶  $(\alpha \cup \beta)^{-1} = \alpha^{-1} \cup \beta^{-1}$
- ▶  $(\alpha \cap \beta)^{-1} = \alpha^{-1} \cap \beta^{-1}$
- ▶  $\overline{\alpha}^{-1} = \overline{\alpha^{-1}}$
- ▶  $(\alpha - \beta)^{-1} = \alpha^{-1} - \beta^{-1}$
- ▶  $\alpha \subset \beta \Rightarrow \alpha^{-1} \subset \beta^{-1}$

13 / 79

## Converse Relation Theorems

$$\overline{\alpha}^{-1} = \overline{\alpha^{-1}}.$$

$$\begin{aligned} & (b, a) \in \overline{\alpha}^{-1} \\ \Leftrightarrow & (a, b) \in \overline{\alpha} \\ \Leftrightarrow & (a, b) \notin \alpha \\ \Leftrightarrow & (b, a) \notin \alpha^{-1} \\ \Leftrightarrow & (b, a) \in \overline{\alpha^{-1}} \end{aligned}$$

□

14 / 79

## Converse Relation Theorems

$$(\alpha \cap \beta)^{-1} = \alpha^{-1} \cap \beta^{-1}.$$

$$\begin{aligned} & (b, a) \in (\alpha \cap \beta)^{-1} \\ \Leftrightarrow & (a, b) \in (\alpha \cap \beta) \\ \Leftrightarrow & (a, b) \in \alpha \wedge (a, b) \in \beta \\ \Leftrightarrow & (b, a) \in \alpha^{-1} \wedge (b, a) \in \beta^{-1} \\ \Leftrightarrow & (b, a) \in \alpha^{-1} \cap \beta^{-1} \end{aligned}$$

□

15 / 79

## Converse Relation Theorems

$$(\alpha - \beta)^{-1} = \alpha^{-1} - \beta^{-1}.$$

$$\begin{aligned} (\alpha - \beta)^{-1} &= (\alpha \cap \overline{\beta})^{-1} \\ &= \alpha^{-1} \cap \overline{\beta}^{-1} \\ &= \alpha^{-1} \cap \overline{\beta^{-1}} \\ &= \alpha^{-1} - \beta^{-1} \end{aligned}$$

□

16 / 79

## Relation Composition Converse

**Theorem**  
 $(\alpha\beta)^{-1} = \beta^{-1}\alpha^{-1}$

**Proof.**

$$\begin{aligned} & (c, a) \in (\alpha\beta)^{-1} \\ \Leftrightarrow & (a, c) \in \alpha\beta \\ \Leftrightarrow & \exists b [(a, b) \in \alpha \wedge (b, c) \in \beta] \\ \Leftrightarrow & \exists b [(b, a) \in \alpha^{-1} \wedge (c, b) \in \beta^{-1}] \\ \Leftrightarrow & (c, a) \in \beta^{-1}\alpha^{-1} \end{aligned}$$

□

17 / 79

## Relation Properties

- ▶  $\alpha \subseteq A \times A$ 
  - ▶ *binary relation on A*
- ▶ let  $\alpha^n$  mean  $\alpha\alpha \cdots \alpha$
- ▶ **identity relation:**  $E = \{(x, x) \mid x \in A\}$

18 / 79

## Reflexivity

reflexive

$$\alpha \subseteq A \times A$$

$$\forall a [a\alpha a]$$

- ▶  $E \subseteq \alpha$
- ▶ nonreflexive:  
 $\exists a [\neg(a\alpha a)]$
- ▶ irreflexive:  
 $\forall a [\neg(a\alpha a)]$

19 / 79

## Reflexivity Examples

Example

$$\mathcal{R}_1 \subseteq \{1, 2\} \times \{1, 2\}$$

$$\mathcal{R}_1 = \{(1, 1), (1, 2), (2, 2)\}$$

- ▶  $\mathcal{R}_1$  is reflexive

Example

$$\mathcal{R}_2 \subseteq \{1, 2, 3\} \times \{1, 2, 3\}$$

$$\mathcal{R}_2 = \{(1, 1), (1, 2), (2, 2)\}$$

- ▶  $\mathcal{R}_2$  is nonreflexive

20 / 79

## Reflexivity Examples

Example

$$\mathcal{R} \subseteq \{1, 2, 3\} \times \{1, 2, 3\}$$

$$\mathcal{R} = \{(1, 2), (2, 1), (2, 3)\}$$

- ▶  $\mathcal{R}$  is irreflexive

21 / 79

## Reflexivity Examples

Example

$$\mathcal{R} \subseteq \mathbb{Z} \times \mathbb{Z}$$

$$\mathcal{R} = \{(a, b) \mid ab \geq 0\}$$

- ▶  $\mathcal{R}$  is reflexive

22 / 79

## Symmetry

symmetric

$$\alpha \subseteq A \times A$$

$$\forall a, b [(a = b) \vee (a\alpha b \wedge b\alpha a) \vee (\neg(a\alpha b) \wedge \neg(b\alpha a))]$$

$$\forall a, b [(a = b) \vee (a\alpha b \leftrightarrow b\alpha a)]$$

- ▶  $\alpha^{-1} = \alpha$
- ▶ asymmetric:  
 $\exists a, b [(a \neq b) \wedge (a\alpha b \wedge \neg(b\alpha a)) \vee (\neg(a\alpha b) \wedge b\alpha a)]$
- ▶ antisymmetric:  
 $\forall a, b [(a = b) \vee (a\alpha b \rightarrow \neg(b\alpha a))]$   
 $\Leftrightarrow \forall a, b [(a = b) \vee \neg(a\alpha b) \vee \neg(b\alpha a)]$   
 $\Leftrightarrow \forall a, b [\neg(a\alpha b \wedge b\alpha a) \vee (a = b)]$   
 $\Leftrightarrow \forall a, b [(a\alpha b \wedge b\alpha a) \rightarrow (a = b)]$

23 / 79

## Symmetry Examples

Example

$$\mathcal{R} \subseteq \{1, 2, 3\} \times \{1, 2, 3\}$$

$$\mathcal{R} = \{(1, 2), (2, 1), (2, 3)\}$$

- ▶  $\mathcal{R}$  is asymmetric

24 / 79

## Symmetry Examples

### Example

$$\mathcal{R} \subseteq \mathbb{Z} \times \mathbb{Z}$$

$$\mathcal{R} = \{(a, b) \mid ab \geq 0\}$$

- $\mathcal{R}$  is symmetric

25 / 79

## Symmetry Examples

### Example

$$\mathcal{R} \subseteq \{1, 2, 3\} \times \{1, 2, 3\}$$

$$\mathcal{R} = \{(1, 1), (2, 2)\}$$

- $\mathcal{R}$  is symmetric and antisymmetric

26 / 79

## Transitivity

### transitive

$$\alpha \subseteq A \times A$$

$$\forall a, b, c [(a\alpha b \wedge b\alpha c) \rightarrow (a\alpha c)]$$

- $\alpha^2 \subseteq \alpha$
- nontransitive:  
 $\exists a, b, c [(a\alpha b \wedge b\alpha c) \wedge \neg(a\alpha c)]$
- antitransitive:  
 $\forall a, b, c [(a\alpha b \wedge b\alpha c) \rightarrow \neg(a\alpha c)]$

27 / 79

## Transitivity Examples

### Example

$$\mathcal{R} \subseteq \{1, 2, 3\} \times \{1, 2, 3\}$$

$$\mathcal{R} = \{(1, 2), (2, 1), (2, 3)\}$$

- $\mathcal{R}$  is antitransitive

28 / 79

## Transitivity Examples

### Example

$$\mathcal{R} \subseteq \mathbb{Z} \times \mathbb{Z}$$

$$\mathcal{R} = \{(a, b) \mid ab \geq 0\}$$

- $\mathcal{R}$  is nontransitive

29 / 79

## Converse Relation Properties

### Theorem

*The reflexivity, symmetry and transitivity properties are preserved in the converse relation.*

30 / 79

## Closures

- ▶ reflexive closure:  
 $r_\alpha = \alpha \cup E$
- ▶ symmetric closure:  
 $s_\alpha = \alpha \cup \alpha^{-1}$
- ▶ transitive closure:  
 $t_\alpha = \bigcup_{i=1,2,3,\dots} \alpha^i = \alpha \cup \alpha^2 \cup \alpha^3 \cup \dots$

31 / 79

## Special Relations

### predecessor - successor

$$\mathcal{R} \subseteq \mathbb{Z} \times \mathbb{Z}$$
$$\mathcal{R} = \{(a, b) \mid a - b = 1\}$$

- ▶ irreflexive
- ▶ antisymmetric
- ▶ antitransitive

32 / 79

## Special Relations

### adjacency

$$\mathcal{R} \subseteq \mathbb{Z} \times \mathbb{Z}$$
$$\mathcal{R} = \{(a, b) \mid |a - b| = 1\}$$

- ▶ irreflexive
- ▶ symmetric
- ▶ antitransitive

33 / 79

## Special Relations

### strict order

$$\mathcal{R} \subseteq \mathbb{Z} \times \mathbb{Z}$$
$$\mathcal{R} = \{(a, b) \mid a < b\}$$

- ▶ irreflexive
- ▶ antisymmetric
- ▶ transitive

34 / 79

## Special Relations

### partial order

$$\mathcal{R} \subseteq \mathbb{Z} \times \mathbb{Z}$$
$$\mathcal{R} = \{(a, b) \mid a \leq b\}$$

- ▶ reflexive
- ▶ antisymmetric
- ▶ transitive

35 / 79

## Special Relations

### preorder

$$\mathcal{R} \subseteq \mathbb{Z} \times \mathbb{Z}$$
$$\mathcal{R} = \{(a, b) \mid |a| \leq |b|\}$$

- ▶ reflexive
- ▶ asymmetric
- ▶ transitive

36 / 79

## Special Relations

### limited difference

$$\mathcal{R} \subseteq \mathbb{Z} \times \mathbb{Z}, m \in \mathbb{Z}^+$$

$$\mathcal{R} = \{(a, b) \mid |a - b| \leq m\}$$

- ▶ reflexive
- ▶ symmetric
- ▶ nontransitive

37 / 79

## Special Relations

### comparability

$$\mathcal{R} \subseteq \mathbb{U} \times \mathbb{U}$$

$$\mathcal{R} = \{(a, b) \mid (a \subseteq b) \vee (b \subseteq a)\}$$

- ▶ reflexive
- ▶ symmetric
- ▶ nontransitive

38 / 79

## Special Relations

### sibling

- ▶ irreflexive
- ▶ symmetric
- ▶ transitive
- ▶ how can a relation be symmetric, transitive and nonreflexive?

39 / 79

## Compatibility Relations

### Definition

compatibility relation:  $\gamma$

- ▶ reflexive
- ▶ symmetric
- ▶ when drawing, lines instead of arrows
- ▶ matrix representation as a triangle matrix
- ▶  $\alpha\alpha^{-1}$  is a compatibility relation


40 / 79

## Compatibility Relation Example

### Example

$$A = \{a_1, a_2, a_3, a_4\}$$

$$\mathcal{R} = \{(a_1, a_1), (a_2, a_2), (a_3, a_3), (a_4, a_4), (a_1, a_2), (a_2, a_1), (a_2, a_4), (a_4, a_2), (a_3, a_4), (a_4, a_3)\}$$



$$\begin{vmatrix} 1 & 1 & 0 & 0 \\ 1 & 1 & 0 & 1 \\ 0 & 0 & 1 & 1 \\ 0 & 1 & 1 & 1 \end{vmatrix}$$

$$\begin{vmatrix} 1 & & & \\ 0 & 0 & & \\ 0 & 1 & 1 & \\ 0 & 1 & 1 & \end{vmatrix}$$

41 / 79

## Compatibility Relation Example

### Example ( $\alpha\alpha^{-1}$ )

$P$ : persons,  $L$ : languages

$$P = \{p_1, p_2, p_3, p_4, p_5, p_6\}$$

$$L = \{l_1, l_2, l_3, l_4, l_5\}$$

$$\alpha \subseteq P \times L$$

$$M_\alpha = \begin{vmatrix} 1 & 1 & 0 & 0 & 0 \\ 1 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 1 \\ 1 & 0 & 1 & 1 & 0 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 1 & 1 & 0 & 0 \end{vmatrix}$$

$$M_{\alpha^{-1}} = \begin{vmatrix} 1 & 1 & 0 & 1 & 0 & 0 \\ 1 & 1 & 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 1 & 0 & 1 \\ 0 & 0 & 0 & 1 & 1 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 \end{vmatrix}$$


42 / 79

## Compatibility Relation Example

Example ( $\alpha\alpha^{-1}$ )

$$\alpha\alpha^{-1} \subseteq P \times P$$

$$M_{\alpha\alpha^{-1}} = \begin{vmatrix} 1 & 1 & 0 & 1 & 0 & 1 \\ 1 & 1 & 0 & 1 & 0 & 1 \\ 0 & 0 & 1 & 1 & 0 & 1 \\ 1 & 1 & 1 & 1 & 1 & 1 \\ 0 & 0 & 0 & 1 & 1 & 0 \\ 1 & 1 & 1 & 1 & 0 & 1 \end{vmatrix}$$



43 / 79

## Compatibility Block

Definition

compatibility block:  $C \subseteq A$


$$\forall a, b [a \in C \wedge b \in C \rightarrow a\gamma b]$$

- ▶ maximal compatibility block:
  - not a subset of another compatibility block
- ▶ an element can be a member of more than one MCB
- ▶ complete cover:  $C_\gamma$ 
  - set of all MCBs

44 / 79

## Compatibility Block Example

Example ( $\alpha\alpha^{-1}$ )



- ▶  $C_1 = \{a_4, a_6\}$
- ▶  $C_2 = \{a_2, a_4, a_6\}$
- ▶  $C_3 = \{a_1, a_2, a_4, a_6\}$  (MCB)

$$C_\gamma(A) = \{ \{a_1, a_2, a_4, a_6\}, \{a_3, a_4, a_6\}, \{a_4, a_5\} \}$$

45 / 79

## Equivalence Relations

Definition

equivalence relation:  $\epsilon$

- ▶ reflexive
- ▶ symmetric
- ▶ transitive
- ▶ equivalence classes (partitions)
- ▶ every element is a member of exactly one equivalence class
- ▶ complete cover:  $C_\epsilon$

46 / 79

## Equivalence Relation Example

Example

$$\mathcal{R} \subseteq \mathbb{Z} \times \mathbb{Z}$$

$$\mathcal{R} = \{(a, b) \mid \exists m \in \mathbb{Z} [a - b = 5m]\}$$

- ▶  $\mathcal{R}$  partitions  $\mathbb{Z}$  into 5 equivalence classes

47 / 79

## References

Required Reading: Grimaldi

- ▶ Chapter 5: Relations and Functions
  - ▶ 5.1. Cartesian Products and Relations
- ▶ Chapter 7: Relations: The Second Time Around
  - ▶ 7.1. Relations Revisited: Properties of Relations
  - ▶ 7.4. Equivalence Relations and Partitions

Supplementary Reading: O'Donnell, Hall, Page

- ▶ Chapter 10: Relations

48 / 79



## Functions

### Definition

**function:**  $f : X \rightarrow Y$

$$\forall x \in X \forall y_1, y_2 \in Y (x, y_1), (x, y_2) \in f \Rightarrow y_1 = y_2$$

- ▶  $X$ : **domain**,  $Y$ : **codomain** (or *range*)
- ▶  $y = f(x)$  is the same as  $(x, y) \in f$
- ▶  $y$  is the *image* of  $x$  under  $f$
- ▶ let  $f : X \rightarrow Y$ , and  $X_1 \subseteq X$   
subset image:  $f(X_1) = \{f(x) \mid x \in X_1\}$

49 / 79

## Subset Image Examples

### Example

$$f : \mathbb{R} \rightarrow \mathbb{R}$$

$$f(x) = x^2$$

$$f(\mathbb{Z}) = \{0, 1, 4, 9, 16, \dots\}$$

$$f(\{-2, 1\}) = \{1, 4\}$$

50 / 79

## Function Properties

### Definition

$f : X \rightarrow Y$  is **one-to-one** (or **injective**):

$$\forall x_1, x_2 \in X f(x_1) = f(x_2) \Rightarrow x_1 = x_2$$

### Definition

$f : X \rightarrow Y$  is **onto** (or **surjective**):

$$\forall y \in Y \exists x \in X f(x) = y$$

- ▶  $f(X) = Y$

### Definition

$f : X \rightarrow Y$  is **bijective**:

$f$  is one-to-one and onto

51 / 79

## One-to-one Function Examples

### Example

$$f : \mathbb{R} \rightarrow \mathbb{R}$$

$$f(x) = 3x + 7$$

$$\begin{aligned} f(x_1) &= f(x_2) \\ \Rightarrow 3x_1 + 7 &= 3x_2 + 7 \\ \Rightarrow 3x_1 &= 3x_2 \\ \Rightarrow x_1 &= x_2 \end{aligned}$$

### Counterexample

$$g : \mathbb{Z} \rightarrow \mathbb{Z}$$

$$g(x) = x^4 - x$$

$$\begin{aligned} g(0) &= 0^4 - 0 = 0 \\ g(1) &= 1^4 - 1 = 0 \end{aligned}$$

52 / 79

## Onto Function Examples

### Example

$$f : \mathbb{R} \rightarrow \mathbb{R}$$

$$f(x) = x^3$$

### Counterexample

$$f : \mathbb{Z} \rightarrow \mathbb{Z}$$

$$f(x) = 3x + 1$$

53 / 79

## Function Composition

### Definition

let  $f : X \rightarrow Y, g : Y \rightarrow Z$

$$g \circ f : X \rightarrow Z$$

$$(g \circ f)(x) = g(f(x))$$

- ▶ function composition is not commutative
- ▶ function composition is associative:  
 $f \circ (g \circ h) = (f \circ g) \circ h$

54 / 79

## Function Composition Examples

### Example (commutativity)

$$f : \mathbb{R} \rightarrow \mathbb{R}$$

$$f(x) = x^2$$

$$g : \mathbb{R} \rightarrow \mathbb{R}$$

$$g(x) = x + 5$$

$$g \circ f : \mathbb{R} \rightarrow \mathbb{R}$$

$$(g \circ f)(x) = x^2 + 5$$

$$f \circ g : \mathbb{R} \rightarrow \mathbb{R}$$

$$(f \circ g)(x) = (x + 5)^2$$

55 / 79

## Composite Function Theorems

### Theorem

let  $f : X \rightarrow Y, g : Y \rightarrow Z$

$f$  is one-to-one  $\wedge$   $g$  is one-to-one  $\Rightarrow g \circ f$  is one-to-one

Proof.

$$\begin{aligned} (g \circ f)(a_1) &= (g \circ f)(a_2) \\ \Rightarrow g(f(a_1)) &= g(f(a_2)) \\ \Rightarrow f(a_1) &= f(a_2) \\ \Rightarrow a_1 &= a_2 \end{aligned}$$

□

56 / 79

## Composite Function Theorems

### Theorem

let  $f : X \rightarrow Y, g : Y \rightarrow Z$

$f$  is onto  $\wedge$   $g$  is onto  $\Rightarrow g \circ f$  is onto

Proof.

$$\forall z \in Z \exists y \in Y g(y) = z$$

$$\forall y \in Y \exists x \in X f(x) = y$$

$$\Rightarrow \forall z \in Z \exists x \in X g(f(x)) = z$$

□

57 / 79

## Identity Function

### Definition

identity function:  $1_X$

$$1_X : X \rightarrow X$$

$$1_X(x) = x$$

58 / 79

## Inverse Function

### Definition

$f : X \rightarrow Y$  is invertible:

$$\exists f^{-1} : Y \rightarrow X [f^{-1} \circ f = 1_X \wedge f \circ f^{-1} = 1_Y]$$

►  $f^{-1}$ : inverse of function  $f$

59 / 79

## Inverse Function Examples

### Example

$$f : \mathbb{R} \rightarrow \mathbb{R}$$

$$f(x) = 2x + 5$$

$$f^{-1} : \mathbb{R} \rightarrow \mathbb{R}$$

$$f^{-1}(x) = \frac{x-5}{2}$$

$$(f^{-1} \circ f)(x) = f^{-1}(f(x)) = f^{-1}(2x + 5) = \frac{(2x+5)-5}{2} = \frac{2x}{2} = x$$

$$(f \circ f^{-1})(x) = f(f^{-1}(x)) = f\left(\frac{x-5}{2}\right) = 2\frac{x-5}{2} + 5 = (x-5) + 5 = x$$

60 / 79

## Inverse Function

### Theorem

If a function is invertible, its inverse is unique.

### Proof.

let  $f : X \rightarrow Y$

let  $g, h : Y \rightarrow X$  such that:

$$g \circ f = 1_X \wedge f \circ g = 1_Y$$

$$h \circ f = 1_X \wedge f \circ h = 1_Y$$

$$h = h \circ 1_Y = h \circ (f \circ g) = (h \circ f) \circ g = 1_X \circ g = g \quad \square$$

61 / 79

## Invertible Function

### Theorem

A function is invertible if and only if it is one-to-one and onto.

62 / 79

## Invertible Function

If invertible then one-to-one.

$f : A \rightarrow B$

$$f(a_1) = f(a_2)$$

$$\Rightarrow f^{-1}(f(a_1)) = f^{-1}(f(a_2))$$

$$\Rightarrow (f^{-1} \circ f)(a_1) = (f^{-1} \circ f)(a_2)$$

$$\Rightarrow 1_A(a_1) = 1_A(a_2)$$

$$\Rightarrow a_1 = a_2 \quad \square$$

If invertible then onto.

$f : A \rightarrow B$

$$b$$

$$= 1_B(b)$$

$$= (f \circ f^{-1})(b)$$

$$= f(f^{-1}(b)) \quad \square$$

63 / 79

## Invertible Function

If bijective then invertible.

$f : A \rightarrow B$

►  $f$  is onto  $\Rightarrow \forall b \in B \exists a \in A f(a) = b$

► let  $g : B \rightarrow A$  be defined by  $a = g(b)$

► is it possible that  $g(b) = a_1 \neq a_2 = g(b)$  ?

► this would mean:  $f(a_1) = b = f(a_2)$

► but  $f$  is one-to-one □

64 / 79

## Pigeonhole Principle

### Definition

**Pigeonhole Principle** (Dirichlet drawers):

If  $m$  pigeons go into  $n$  holes and  $m > n$ , then at least one hole contains more than one pigeon.

- let  $f : X \rightarrow Y$   
if  $|X| > |Y|$  then  $f$  cannot be one-to-one
- $\exists x_1, x_2 \in X [x_1 \neq x_2 \wedge f(x_1) = f(x_2)]$

65 / 79

## Pigeonhole Principle Examples

### Example

- Among 367 people, at least two have the same birthday.
- In an exam where the grades integers between 0 and 100, how many students have to take the exam to make sure that at least two students will have the same grade?

66 / 79

## Generalized Pigeonhole Principle

### Definition

#### Generalized Pigeonhole Principle:

If  $m$  objects are distributed to  $n$  drawers, then at least one of the drawers contains  $\lceil m/n \rceil$  objects.

### Example

Among 100 people, at least 9 ( $\lceil 100/12 \rceil$ ) were born in the same month.

67 / 79

## Pigeonhole Principle Example

### Theorem

In any subset of cardinality 6 of the set  $S = \{1, 2, 3, \dots, 9\}$ , there are two elements which total 10.

68 / 79

## Pigeonhole Principle Example

### Theorem

Let  $S$  be a set of positive integers smaller than or equal to 14, with cardinality 6. The sums of the elements in all nonempty subsets of  $S$  cannot be all different.

### Proof Trial

$A \subseteq S$

$s_A$  : sum of the elements of  $A$

- ▶ holes:  
 $1 \leq s_A \leq 9 + \dots + 14 = 69$
- ▶ pigeons:  $2^6 - 1 = 63$

### Proof.

look at the subsets for which  $|A| \leq 5$

- ▶ holes:  
 $1 \leq s_A \leq 10 + \dots + 14 = 60$
- ▶ pigeons:  $2^6 - 2 = 62$

□

69 / 79

## Pigeonhole Principle Example

### Theorem

There is at least one pair of elements among 101 elements chosen from set  $S = \{1, 2, 3, \dots, 200\}$ , so that one of the elements of the pair divides the other.

### Proof Method

- ▶ we first show that  
 $\forall n \exists! p [n = 2^r p \wedge r \in \mathbb{N} \wedge \exists t \in \mathbb{Z} [p = 2t + 1]]$
- ▶ then, by using this theorem we prove the main theorem

70 / 79

## Pigeonhole Principle Example

### Theorem

$\forall n \exists! p [n = 2^r p \wedge r \in \mathbb{N} \wedge \exists t \in \mathbb{Z} [p = 2t + 1]]$

### Proof of existence.

$n = 1$ :  $r = 0, p = 1$

$n \leq k$ : assume  $n = 2^r p$

$n = k + 1$ :

$n = 2$ :  $r = 1, p = 1$   
 $n$  prime ( $n > 2$ ):  $r = 0, p = n$   
 $\neg(n \text{ prime})$ :  $n = n_1 n_2$   
 $n = 2^{r_1} p_1 \cdot 2^{r_2} p_2$   
 $n = 2^{r_1 + r_2} \cdot p_1 p_2$

□

### Proof of uniqueness.

if not unique:

$$\begin{aligned} n &= 2^{r_1} p_1 = 2^{r_2} p_2 \\ \Rightarrow 2^{r_1 - r_2} p_1 &= p_2 \\ \Rightarrow 2 | p_2 \end{aligned}$$

□

71 / 79

## Pigeonhole Principle Example

### Theorem

There is at least one pair of elements among 101 elements chosen from set  $S = \{1, 2, 3, \dots, 200\}$  so that one of the elements of the pair divides the other.

### Proof.

- ▶  $T = \{t \mid t \in S, \exists i \in \mathbb{Z} [t = 2i + 1]\}, |T| = 100$
- ▶  $f : S \rightarrow T, r \in \mathbb{N}$  olsun  
 $s = 2^r t \rightarrow f(s) = t$ 
  - ▶ if 101 elements are chosen from  $S$ , at least two of them will have the same image in  $T$ :  $f(s_1) = f(s_2) \Rightarrow 2^{m_1} t = 2^{m_2} t$

$$\frac{s_1}{s_2} = \frac{2^{m_1} t}{2^{m_2} t} = 2^{m_1 - m_2}$$

□

72 / 79

## Recursive Functions

### Definition

**recursive function:** a function defined in terms of itself

$$f(n) = h(f(m))$$

- ▶ *inductively defined function:* a recursive function where the size is reduced at every step

$$f(n) = \begin{cases} k & n = 0 \\ h(f(n-1)) & n > 0 \end{cases}$$

73 / 79

## Recursion Examples

### Example

$$f_{91}(n) = \begin{cases} n - 10 & n > 100 \\ f_{91}(f_{91}(n + 11)) & n \leq 100 \end{cases}$$

### Example (factorial)

$$f(n) = \begin{cases} 1 & n = 0 \\ n \cdot f(n-1) & n > 0 \end{cases}$$

74 / 79

## Euclid Algorithm

### Example (greatest common divisor)

$$\gcd(a, b) = \begin{cases} b & b|a \\ \gcd(b, a \bmod b) & b \nmid a \end{cases}$$

$$\begin{aligned} \gcd(333, 84) &= \gcd(84, 333 \bmod 84) \\ &= \gcd(84, 81) \\ &= \gcd(81, 84 \bmod 81) \\ &= \gcd(81, 3) \\ &= 3 \end{aligned}$$

75 / 79

## Fibonacci Series

### Fibonacci series

$$F_n = \text{fib}(n) = \begin{cases} 1 & n = 1 \\ 1 & n = 2 \\ \text{fib}(n-1) + \text{fib}(n-2) & n > 2 \end{cases}$$

$$\begin{array}{ccccccccccc} F_1 & F_2 & F_3 & F_4 & F_5 & F_6 & F_7 & F_8 & \dots \\ 1 & 1 & 2 & 3 & 5 & 8 & 13 & 21 & \dots \end{array}$$

76 / 79

## Fibonacci Series

### Theorem

$$\sum_{i=1}^n F_i^2 = F_n \cdot F_{n+1}$$

### Proof.

$$n = 2: \quad \sum_{i=1}^2 F_i^2 = F_1^2 + F_2^2 = 1 + 1 = 1 \cdot 2 = F_2 \cdot F_3$$

$$n = k: \quad \sum_{i=1}^k F_i^2 = F_k \cdot F_{k+1}$$

$$\begin{aligned} n = k+1: \quad \sum_{i=1}^{k+1} F_i^2 &= \sum_{i=1}^k F_i^2 + F_{k+1}^2 \\ &= F_k \cdot F_{k+1} + F_{k+1}^2 \\ &= F_{k+1} \cdot (F_k + F_{k+1}) \\ &= F_{k+1} \cdot F_{k+2} \end{aligned}$$

□

77 / 79

## Ackermann Function

### Ackermann function

$$\text{ack}(x, y) = \begin{cases} y + 1 & x = 0 \\ \text{ack}(x-1, 1) & y = 0 \\ \text{ack}(x-1, \text{ack}(x, y-1)) & x > 0 \wedge y > 0 \end{cases}$$

78 / 79

## References

### Required Reading: Grimaldi

- ▶ Chapter 5: Relations and Functions
  - ▶ 5.2. Functions: Plain and One-to-One
  - ▶ 5.3. Onto Functions: Stirling Numbers of the Second Kind
  - ▶ 5.5. The Pigeonhole Principle
  - ▶ 5.6. Function Composition and Inverse Functions

### Supplementary Reading: O'Donnell, Hall, Page

- ▶ Chapter 11: Functions