

Discrete Mathematics

Predicates and Sets

H. Turgut Uyar Ayşegül Gençata Yayımlı Emre Harmancı

2001-2013

1 / 43

License



©2001-2013 T. Uyar, A. Yayımlı, E. Harmancı

You are free:

- ▶ to Share – to copy, distribute and transmit the work
- ▶ to Remix – to adapt the work

Under the following conditions:

- ▶ Attribution – You must attribute the work in the manner specified by the author or licensor (but not in any way that suggests that they endorse you or your use of the work).
- ▶ Noncommercial – You may not use this work for commercial purposes.
- ▶ Share Alike – If you alter, transform, or build upon this work, you may distribute the resulting work only under the same or similar license to this one.

Legal code (the full license):

<http://creativecommons.org/licenses/by-nc-sa/3.0/>

2 / 43

Topics

Predicates

Introduction
Quantifiers
Multiple Quantifiers

Sets

Introduction
Subset
Set Operations
Inclusion-Exclusion

3 / 43

Predicate

Definition

predicate (or **open statement**): a declarative sentence which

- ▶ contains one or more variables, and
- ▶ is not a proposition, but
- ▶ becomes a proposition when the variables in it are replaced by certain allowable choices

4 / 43

Universe of Discourse

Definition

universe of discourse: \mathcal{U}
set of allowable choices

- ▶ examples:
 - ▶ \mathbb{Z} : integers
 - ▶ \mathbb{N} : natural numbers
 - ▶ \mathbb{Z}^+ : positive integers
 - ▶ \mathbb{Q} : rational numbers
 - ▶ \mathbb{R} : real numbers
 - ▶ \mathbb{C} : complex numbers

5 / 43

Predicate Examples

Example

$\mathcal{U} = \mathbb{N}$

$p(x)$: $x + 2$ is an even integer

$p(5)$: F

$p(8)$: T

$\neg p(x)$: $x + 2$ is not an even integer

Example

$\mathcal{U} = \mathbb{N}$

$q(x, y)$: $x + y$ and $x - 2y$ are even integers

$q(11, 3)$: F , $q(14, 4)$: T

6 / 43

Quantifiers

Definition

existential quantifier:

predicate is true for some values

- ▶ symbol: \exists
- ▶ read: *there exists*
- ▶ symbol: $\exists!$
- ▶ read: *there exists only one*

Definition

universal quantifier:

predicate is true for all values

- ▶ symbol: \forall
- ▶ read: *for all*

7 / 43

Quantifiers

existential quantifier

$$\mathcal{U} = \{x_1, x_2, \dots, x_n\}$$

$$\exists x \, p(x) \equiv p(x_1) \vee p(x_2) \vee \dots \vee p(x_n)$$

- ▶ $p(x)$ is true for some x

universal quantifier

$$\mathcal{U} = \{x_1, x_2, \dots, x_n\}$$

$$\forall x \, p(x) \equiv p(x_1) \wedge p(x_2) \wedge \dots \wedge p(x_n)$$

- ▶ $p(x)$ is true for all x

8 / 43

Quantifier Examples

Example

$$\mathcal{U} = \mathbb{R}$$

- ▶ $p(x) : x \geq 0$
- ▶ $q(x) : x^2 \geq 0$
- ▶ $r(x) : (x-4)(x+1) = 0$
- ▶ $s(x) : x^2 - 3 > 0$
- ▶ $\exists x [p(x) \wedge r(x)]$
- ▶ $\forall x [p(x) \rightarrow q(x)]$
- ▶ $\forall x [q(x) \rightarrow s(x)]$
- ▶ $\forall x [r(x) \vee s(x)]$
- ▶ $\forall x [r(x) \rightarrow p(x)]$

are the following expressions true?

9 / 43

Negating Quantifiers

- ▶ replace \forall with \exists , and \exists with \forall
- ▶ negate the predicate

$$\neg \exists x \, p(x) \Leftrightarrow \forall x \, \neg p(x)$$

$$\neg \exists x \, \neg p(x) \Leftrightarrow \forall x \, p(x)$$

$$\neg \forall x \, p(x) \Leftrightarrow \exists x \, \neg p(x)$$

$$\neg \forall x \, \neg p(x) \Leftrightarrow \exists x \, p(x)$$

10 / 43

Negating Quantifiers

Theorem

$$\neg \exists x \, p(x) \Leftrightarrow \forall x \, \neg p(x)$$

Proof.

$$\begin{aligned} \neg \exists x \, p(x) &\equiv \neg[p(x_1) \vee p(x_2) \vee \dots \vee p(x_n)] \\ &\Leftrightarrow \neg p(x_1) \wedge \neg p(x_2) \wedge \dots \wedge \neg p(x_n) \\ &\equiv \forall x \, \neg p(x) \end{aligned}$$

□

11 / 43

Predicate Equivalences

Theorem

$$\exists x [p(x) \vee q(x)] \Leftrightarrow \exists x \, p(x) \vee \exists x \, q(x)$$

Theorem

$$\forall x [p(x) \wedge q(x)] \Leftrightarrow \forall x \, p(x) \wedge \forall x \, q(x)$$

12 / 43

Predicate Implications

Theorem

$$\forall x \, p(x) \Rightarrow \exists x \, p(x)$$

Theorem

$$\exists x \, [p(x) \wedge q(x)] \Rightarrow \exists x \, p(x) \wedge \exists x \, q(x)$$

Theorem

$$\forall x \, p(x) \vee \forall x \, q(x) \Rightarrow \forall x \, [p(x) \vee q(x)]$$

13 / 43

Multiple Quantifiers

- ▶ $\exists x \exists y \, p(x, y)$
- ▶ $\forall x \exists y \, p(x, y)$
- ▶ $\exists x \forall y \, p(x, y)$
- ▶ $\forall x \forall y \, p(x, y)$

14 / 43

Multiple Quantifier Examples

Example

$$\mathcal{U} = \mathbb{Z}$$

$$p(x, y) : x + y = 17$$

- ▶ $\forall x \exists y \, p(x, y)$:
for every x there exists a y such that $x + y = 17$
- ▶ $\exists y \forall x \, p(x, y)$:
there exists a y so that for all x , $x + y = 17$
- ▶ what if $\mathcal{U} = \mathbb{N}$?

15 / 43

Multiple Quantifiers

Example

$$\mathcal{U}_x = \{1, 2\} \wedge \mathcal{U}_y = \{A, B\}$$

$$\begin{aligned}\exists x \exists y \, p(x, y) &\equiv [p(1, A) \vee p(1, B)] \vee [p(2, A) \vee p(2, B)] \\ \exists x \forall y \, p(x, y) &\equiv [p(1, A) \wedge p(1, B)] \vee [p(2, A) \wedge p(2, B)] \\ \forall x \exists y \, p(x, y) &\equiv [p(1, A) \vee p(1, B)] \wedge [p(2, A) \vee p(2, B)] \\ \forall x \forall y \, p(x, y) &\equiv [p(1, A) \wedge p(1, B)] \wedge [p(2, A) \wedge p(2, B)]\end{aligned}$$

16 / 43

References

Required Reading: Grimaldi

- ▶ Chapter 2: Fundamentals of Logic
 - ▶ 2.4. The Use of Quantifiers

Supplementary Reading: O'Donnell, Hall, Page

- ▶ Chapter 7: Predicate Logic

17 / 43

Set

Definition

set: a collection of elements that are

- ▶ distinct
- ▶ unordered
- ▶ non-repeating

18 / 43

Set Representation

- ▶ *explicit representation*
elements are listed within braces: $\{a_1, a_2, \dots, a_n\}$
- ▶ *implicit representation*
elements that validate a predicate: $\{x | x \in G, p(x)\}$
- ▶ \emptyset : empty set
- ▶ let S be a set, and a be an element
 - ▶ $a \in S$: a is an element of set S
 - ▶ $a \notin S$: a is not an element of set S
- ▶ $|S|$: number of elements (**cardinality**)

19 / 43

Explicit Representation Example

Example

$\{3, 8, 2, 11, 5\}$
 $11 \in \{3, 8, 2, 11, 5\}$
 $|\{3, 8, 2, 11, 5\}| = 5$

20 / 43

Implicit Representation Examples

Example

$\{x | x \in \mathbb{Z}^+, 20 < x^3 < 100\} \equiv \{3, 4\}$
 $\{2x - 1 | x \in \mathbb{Z}^+, 20 < x^3 < 100\} \equiv \{5, 7\}$

Example

$A = \{x | x \in \mathbb{R}, 1 \leq x \leq 5\}$

Example

$E = \{n | n \in \mathbb{N}, \exists k \in \mathbb{N} [n = 2k]\}$
 $A = \{x | x \in E, 1 \leq x \leq 5\}$

21 / 43

Set Dilemma

- ▶ There is a barber who lives in a small town.
He shaves all those men who don't shave themselves.
He doesn't shave those men who shave themselves.
Does the barber shave himself?
- ▶ yes \rightarrow but he doesn't shave men who shave themselves
 \rightarrow no
- ▶ no \rightarrow but he shaves all men who don't shave themselves
 \rightarrow yes

22 / 43

Set Dilemma

- ▶ let S be the set of sets that are not an element of themselves
 $S = \{A | A \notin A\}$
Is S an element of itself?
- ▶ yes \rightarrow but the predicate is not valid \rightarrow no
- ▶ no \rightarrow but the predicate is valid \rightarrow yes

23 / 43

Subset

Definition

$A \subseteq B \Leftrightarrow \forall x [x \in A \rightarrow x \in B]$

- ▶ **set equality**:
 $A = B \Leftrightarrow (A \subseteq B) \wedge (B \subseteq A)$
- ▶ **proper subset**:
 $A \subset B \Leftrightarrow (A \subseteq B) \wedge (A \neq B)$
- ▶ $\forall S [\emptyset \subseteq S]$

24 / 43

Subset

not a subset

$$\begin{aligned} A \not\subseteq B &\Leftrightarrow \neg \forall x [x \in A \rightarrow x \in B] \\ &\Leftrightarrow \exists x \neg [x \in A \rightarrow x \in B] \\ &\Leftrightarrow \exists x \neg [\neg(x \in A) \vee (x \in B)] \\ &\Leftrightarrow \exists x [(x \in A) \wedge \neg(x \in B)] \\ &\Leftrightarrow \exists x [(x \in A) \wedge (x \notin B)] \end{aligned}$$

25 / 43

Power Set

Definition

power set: $\mathcal{P}(S)$

the set of all subsets of a set, including the empty set and the set itself

- ▶ if a set has n elements, its power set has 2^n elements

26 / 43

Example of Power Set

Example

$$\mathcal{P}(\{1, 2, 3\}) = \{ \emptyset, \{1\}, \{2\}, \{3\}, \{1, 2\}, \{1, 3\}, \{2, 3\}, \{1, 2, 3\} \}$$

27 / 43

Set Operations

complement

$$\overline{A} = \{x | x \notin A\}$$

intersection

$$A \cap B = \{x | (x \in A) \wedge (x \in B)\}$$

- ▶ if $A \cap B = \emptyset$ then A and B are **disjoint**

union

$$A \cup B = \{x | (x \in A) \vee (x \in B)\}$$

28 / 43

Set Operations

difference

$$A - B = \{x | (x \in A) \wedge (x \notin B)\}$$

- ▶ $A - B = A \cap \overline{B}$
- ▶ **symmetric difference:**
 $A \triangle B = \{x | (x \in A \cup B) \wedge (x \notin A \cap B)\}$

29 / 43

Cartesian Product

Definition

Cartesian product:

$$A \times B = \{(a, b) | a \in A, b \in B\}$$

$$A \times B \times C \times \cdots \times N = \{(a, b, \dots, n) | a \in A, b \in B, \dots, n \in N\}$$

- ▶ $|A \times B \times C \times \cdots \times N| = |A| \cdot |B| \cdot |C| \cdots |N|$

30 / 43

Cartesian Product Example

Example

$$A = \{a_1, a_2, a_3, a_4\}$$

$$B = \{b_1, b_2, b_3\}$$

$$A \times B = \{ \\ (a_1, b_1), (a_1, b_2), (a_1, b_3), \\ (a_2, b_1), (a_2, b_2), (a_2, b_3), \\ (a_3, b_1), (a_3, b_2), (a_3, b_3), \\ (a_4, b_1), (a_4, b_2), (a_4, b_3) \\ \}$$

31 / 43

Equivalences

Double Complement

$$\overline{\overline{A}} = A$$

Commutativity

$$A \cap B = B \cap A$$

$$A \cup B = B \cup A$$

Associativity

$$(A \cap B) \cap C = A \cap (B \cap C) \quad (A \cup B) \cup C = A \cup (B \cup C)$$

Idempotence

$$A \cap A = A$$

$$A \cup A = A$$

Inverse

$$A \cap \overline{A} = \emptyset$$

$$A \cup \overline{A} = \mathcal{U}$$

32 / 43

Equivalences

Identity

$$A \cap \mathcal{U} = A$$

$$A \cup \emptyset = A$$

Domination

$$A \cap \emptyset = \emptyset$$

$$A \cup \mathcal{U} = \mathcal{U}$$

Distributivity

$$A \cap (B \cup C) = (A \cap B) \cup (A \cap C) \quad A \cup (B \cap C) = (A \cup B) \cap (A \cup C)$$

Absorption

$$A \cap (A \cup B) = A$$

$$A \cup (A \cap B) = A$$

DeMorgan's Laws

$$\overline{A \cap B} = \overline{A} \cup \overline{B}$$

$$\overline{A \cup B} = \overline{A} \cap \overline{B}$$

33 / 43

DeMorgan's Laws

Proof.

$$\begin{aligned} \overline{A \cap B} &= \{x | x \notin (A \cap B)\} \\ &= \{x | \neg(x \in (A \cap B))\} \\ &= \{x | \neg((x \in A) \wedge (x \in B))\} \\ &= \{x | \neg(x \in A) \vee \neg(x \in B)\} \\ &= \{x | (x \notin A) \vee (x \notin B)\} \\ &= \{x | (x \in \overline{A}) \vee (x \in \overline{B})\} \\ &= \{x | x \in \overline{A} \cup \overline{B}\} \\ &= \overline{A} \cup \overline{B} \end{aligned}$$

□

34 / 43

Example of Equivalence

Theorem

$$A \cap (B - C) = (A \cap B) - (A \cap C)$$

35 / 43

Equivalence Example

Proof.

$$\begin{aligned} (A \cap B) - (A \cap C) &= (A \cap B) \cap \overline{(A \cap C)} \\ &= (A \cap B) \cap (\overline{A} \cup \overline{C}) \\ &= ((A \cap B) \cap \overline{A}) \cup ((A \cap B) \cap \overline{C}) \\ &= \emptyset \cup ((A \cap B) \cap \overline{C}) \\ &= (A \cap B) \cap \overline{C} \\ &= A \cap (B \cap \overline{C}) \\ &= A \cap (B - C) \end{aligned}$$

□

36 / 43

Principle of Inclusion-Exclusion

- ▶ $|A \cup B| = |A| + |B| - |A \cap B|$
- ▶ $|A \cup B \cup C| = |A| + |B| + |C| - (|A \cap B| + |A \cap C| + |B \cap C|) + |A \cap B \cap C|$

Theorem

$$|A_1 \cup A_2 \cup \dots \cup A_n| = \sum_i |A_i| - \sum_{i,j} |A_i \cap A_j| + \sum_{i,j,k} |A_i \cap A_j \cap A_k| - \dots + (-1)^{n-1} |A_1 \cap A_2 \cap \dots \cap A_n|$$

37 / 43

Inclusion-Exclusion Example

Example (sieve of Eratosthenes)

- ▶ a method to identify prime numbers

2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17
18	19	20	21	22	23	24	25	26	27	28	29	30			
2	3		5		7		9		11		13		15		17
	19		21		23		25		27		29				
2	3		5		7				11		13				17
	19				23		25				29				
2	3		5		7				11		13				17
	19				23						29				

38 / 43

Inclusion-Exclusion Example

Example (sieve of Eratosthenes)

- ▶ number of primes between 1 and 100
- ▶ numbers that are not divisible by 2, 3, 5 and 7
 - ▶ A_2 : set of numbers divisible by 2
 - ▶ A_3 : set of numbers divisible by 3
 - ▶ A_5 : set of numbers divisible by 5
 - ▶ A_7 : set of numbers divisible by 7
- ▶ $|A_2 \cup A_3 \cup A_5 \cup A_7|$

39 / 43

Inclusion-Exclusion Example

Example (sieve of Eratosthenes)

- ▶ $|A_2| = \lfloor 100/2 \rfloor = 50$
- ▶ $|A_3| = \lfloor 100/3 \rfloor = 33$
- ▶ $|A_5| = \lfloor 100/5 \rfloor = 20$
- ▶ $|A_7| = \lfloor 100/7 \rfloor = 14$
- ▶ $|A_2 \cap A_3| = \lfloor 100/6 \rfloor = 16$
- ▶ $|A_2 \cap A_5| = \lfloor 100/10 \rfloor = 10$
- ▶ $|A_2 \cap A_7| = \lfloor 100/14 \rfloor = 7$
- ▶ $|A_3 \cap A_5| = \lfloor 100/15 \rfloor = 6$
- ▶ $|A_3 \cap A_7| = \lfloor 100/21 \rfloor = 4$
- ▶ $|A_5 \cap A_7| = \lfloor 100/35 \rfloor = 2$

40 / 43

Inclusion-Exclusion Example

Example (sieve of Eratosthenes)

- ▶ $|A_2 \cap A_3 \cap A_5| = \lfloor 100/30 \rfloor = 3$
- ▶ $|A_2 \cap A_3 \cap A_7| = \lfloor 100/42 \rfloor = 2$
- ▶ $|A_2 \cap A_5 \cap A_7| = \lfloor 100/70 \rfloor = 1$
- ▶ $|A_3 \cap A_5 \cap A_7| = \lfloor 100/105 \rfloor = 0$
- ▶ $|A_2 \cap A_3 \cap A_5 \cap A_7| = \lfloor 100/210 \rfloor = 0$

41 / 43

Inclusion-Exclusion Example

Example (sieve of Eratosthenes)

$$\begin{aligned}
 |A_2 \cup A_3 \cup A_5 \cup A_7| &= (50 + 33 + 20 + 14) \\
 &\quad - (16 + 10 + 7 + 6 + 4 + 2) \\
 &\quad + (3 + 2 + 1 + 0) \\
 &\quad - (0) \\
 &= 78
 \end{aligned}$$

- ▶ number of primes: $(100 - 78) + 4 - 1 = 25$

42 / 43

References

Required Reading: Grimaldi

- ▶ Chapter 3: Set Theory
 - ▶ 3.1. [Sets and Subsets](#)
 - ▶ 3.2. [Set Operations and the Laws of Set Theory](#)
- ▶ Chapter 8: The Principle of Inclusion and Exclusion
 - ▶ 8.1. [The Principle of Inclusion and Exclusion](#)

Supplementary Reading: O'Donnell, Hall, Page

- ▶ Chapter 8: Set Theory