

TA: Ondřej Čertík
web: <http://hpfem.math.unr.edu/~ondrej/>
class: MATH 181
date: January 29, 2009

1 Introduction

We did examples from the 2.4 section and Quiz 4.

2 Problem 1

Excercise 3, section 2.4. Solution is on the page A89.

3 Problem 2

Excercise 5:

Sketch the graph of a function that is continuous everywhere except at $x = 3$ and is continuous from the left at 3.

Solution: one such function is given in the solution at the page A89.

4 Problem 3

Excercise 15. The graph is sketched on the page A90. The function needs to be rewritten by factoring out the nominator:

$$\frac{x^2 - x - 12}{x + 3} = \frac{(x - 4)(x + 3)}{x + 3} = x - 4$$

See the solution to Quiz 4 for more information.

5 Problem 4

Excercise 21. Given the function $G(t) = \ln(t^4 - 1)$, find the domain, why the function is continuous at every number in its domain?

Solution: The logarithm $\ln x$ is defined only for $x > 0$. We want the logarithm to be real, e.g. $\ln(-1)$ is not defined. Otherwise, if you remember complex numbers, logarithm can be defined for any (complex) number $z = \rho e^{i\theta}$ as $\ln \rho e^{i\theta} = \ln \rho + i\theta$ and it depends on the branch cut (you can add any number of 2π to θ).

6 Quizzes

We did Quiz 4.