

A geometric account of the solution to the Dixmier unitarizability problem

Vadim Alekseev

ABSTRACT. This note presents a more geometric account of OpenAI’s solution to the Dixmier unitarizability problem. Some parts of the original proof are reinterpreted using the existence of a positive divergence flow on a non-amenable group, and the randomized construction for operator norm control of the Markov operator with matrix coefficients is replaced by one coming from free groups.

Introduction

A representation $\pi : G \rightarrow \mathrm{GL}(H)$ on a complex Hilbert space is *uniformly bounded* if $|\pi| := \sup_{g \in G} \|\pi(g)\| < \infty$, and *unitarizable* if a bounded invertible operator V makes $V\pi(g)V^{-1}$ unitary for every $g \in G$. A group is unitarizable if all its uniformly bounded Hilbert space representations are unitarizable. Sz.-Nagy’s theorem on operators with uniformly bounded integer powers established the case $G = \mathbb{Z}$ [SzN47]. Day and Dixmier extended the invariant-mean argument to amenable groups [Day50, Dix50]; see also Nakamura–Takeda [NT51]. Dixmier immediately asked about the converse [Dix50, § 5].

OpenAI’s preprint [OAI26] proves this converse for discrete groups. The purpose of this note is to give a geometric account of its operator obstruction and a construction of the required positive matrix coefficients using free products of finite groups. We work throughout with countable discrete groups (since the reduction to the general case is straightforward) and prove the following statement. Together with the Day–Dixmier theorem [Day50, Dix50], this characterizes amenability among discrete groups.

MAIN THEOREM. *Every countable nonamenable discrete group is nonunitarizable.*

The history of the problem involves several different ways of detecting nonunitarizability. Ehrenpreis and Mautner produced the first examples for $\mathrm{SL}_2(\mathbb{R})$ in 1955 [EM55], with a detailed development by Kunze and Stein [KS60]. Explicit constructions for nonabelian free groups were obtained subsequently, including the analytic families of Pytlik and Szwarc [PS86]. Since unitarizability passes to subgroups, these results imply nonunitarizability for every group containing a nonabelian free subgroup. The remaining difficulty was to obtain an obstruction for general nonamenable groups, including those without such subgroups.

Later work developed quantitative criteria for unitarizability [Pis98, Pis05]. Random-forest methods and Osin’s constructions produced nonunitarizable groups without free subgroups [EM09, Osi09], while measurable-group techniques yielded results for wreath products, later extended to finite lamp groups by Alpeev [Mon06, GL09, MO10, Alp23]. Further work connected unitarizability with fixed price and approximation properties, including for groups acting on finite-dimensional $\mathrm{CAT}(0)$ cube complexes [Tho16, Ver23].

The analytic approaches closest to the present proof are Littlewood-function methods for amenability and unitarizability [Wys88, BF91, EM09], later related to asymptotic Cheeger constants by Gerasimova, Gruber, Monod, and Thom [GGMT20].

The general idea of the proof given here can be explained as follows. Since the group is nonamenable, it has a flow of positive divergence which splits the random-walk operator on the Cayley graph into contributions from walks with positive and non-positive total flow. Thus we can decompose the Markov operators (acting from the right) on bigger and bigger powers of the generating set into the “non-positive flow” and “positive flow” parts, called T resp. Q (we omit the index counting the powers of the generating set here). Positive divergence makes the relevant non-positive part sparse, which allows to uniformly control the operator norm displacement of T under the G -action. At the same time, the resulting tracial 2-norms of the Markov operators, if one allows suitable matrix amplifications, tend to infinity, which together with sparsity control on T shows via a tracial distance estimate that T move further and further from the right group von Neumann algebra. Therefore the cocycle given at each stage by $T - U(g)TU(g)^{-1}$ remains bounded in the direct sum, but becomes non-inner there, which yields a nonunitarizable uniformly bounded representation.

AI assistance. AI tools contributed to several stages of the work leading to this paper, starting with OpenAI’s solution to the Dixmier unitarizability problem.

The present exposition was developed through interaction of GPT-6 Astra with subsequent editing by the author. The author supplied ideas reinterpreting the original proof in a geometric way as well as placing the relevant quantities in the proper operator-algebraic context, which were then made precise by Astra; the usage of group-theoretic sources instead of random matrix methods was suggested by the author and worked out by Astra in detail with subsequent manual editing by the author to improve presentation. The author had written the relevant explanations of the proof ideas and steps and takes responsibility for the mathematical claims, references, and final presentation.

1. Preliminaries and notation

Operator algebras. Let G be a countable discrete group, with identity e . Inner products are linear in the second variable. On $\ell^2(G)$ let

$$(1.1) \quad (\lambda(g)\xi)(x) = \xi(g^{-1}x), \quad (\rho(g)\xi)(x) = \xi(xg), \quad LG = \lambda(G)'', \quad RG = \rho(G)'' = LG'.$$

The canonical traces on LG and RG , the normalized matrix traces, and their tensor products are denoted by τ :

$$(1.2) \quad \tau(X) = \langle \delta_e, X\delta_e \rangle, \quad \tau(B) = \frac{1}{k} \text{Tr}(B), \quad \|A\|_2 = \tau(A^*A)^{1/2}.$$

Every element of $LG \otimes M_k(\mathbb{C})$, respectively $RG \otimes M_k(\mathbb{C})$, has a unique Fourier expansion converging in L^2 :

$$(1.3) \quad A = \sum_{g \in G} \lambda(g) \otimes A_g \quad \text{or} \quad A = \sum_{g \in G} \rho(g) \otimes A_g, \quad \|A\|_2^2 = \sum_g \|A_g\|_2^2.$$

Put $H_k = \ell^2(G) \otimes \mathbb{C}^k$. Viewing δ_x as a map $\mathbb{C} \rightarrow \ell^2(G)$, define for $X \in \mathcal{B}(H_k)$

$$(1.4) \quad \text{row}_x(X) = (\delta_x^* \otimes I_k)X, \quad \text{col}_x(X) = X(\delta_x \otimes I_k).$$

For $A \in RG \otimes M_k(\mathbb{C})$, the trace identities give

$$(1.5) \quad \|A\|_2^2 = \tau(\text{row}_e(A) \text{row}_e(A)^*) = \tau(\text{col}_e(A)^* \text{col}_e(A)).$$

Hence $\|A\|_2 \leq \|\text{row}_e(A)\|$ and $\|A\|_2 \leq \|\text{col}_e(A)\|$.

Littlewood functions. Recall that $f \in T^1(G)$ if $f(x^{-1}y) = u(x, y) + v(x, y)$ with

$$(1.6) \quad \|f\|_{T^1} = \inf_{f(x^{-1}y)=u+v} (\alpha(u) + \gamma(v)) < \infty, \quad \alpha(u) = \sup_x \sum_y |u(x, y)|, \quad \gamma(v) = \sup_y \sum_x |v(x, y)|.$$

See [GGMT20, Definition 2.3 and Propositions 2.5–2.6]. For a finitely supported f and a decomposition $\rho(f) = T + Q$ with kernels u, v , Schur’s test gives

$$(1.7) \quad \sup_g \|T - \lambda(g)T\lambda(g)^*\| \leq 2\sqrt{\alpha(u)\gamma(v)}.$$

Indeed the difference has row bound $2\alpha(u)$ and, since it also equals $\lambda(g)Q\lambda(g)^* - Q$, column bound $2\gamma(v)$. Lemma 4.1 gives the corresponding estimate for positive matrix coefficients.

For unitarizable G , the standard Littlewood estimate is $\|f\|_{\ell^2(G)} \leq C_G \|f\|_{T^1(G)}$; see Bożejko–Fendler [BF91] and Epstein–Monod [EM09, Proposition 2.4]. Here $\|f\|_{\ell^2(G)} = \|\rho(f)\|_2$. The construction below uses the same row–column estimates and tracial norm, with matrix coefficients.

2. Positive and non-positive walks

Let G be a countable nonamenable discrete group. By the Følner criterion, there is a finite symmetric set $S \subset G \setminus \{e\}$ such that, with $d = |S|$,

$$(2.1) \quad h := \inf_{\emptyset \neq F \subset G \text{ finite}} \frac{|\partial_E F|}{d|F|} > 0, \quad \partial_E F = \{(x, xs) : x \in F, s \in S, xs \notin F\}.$$

Fix $0 < \eta < \min(h, 1)$ and set $\beta = \sqrt{1 - \eta^2}$.

On the right Cayley graph, an antisymmetric flow gives a signed value to each oriented edge. Its integral along a walk changes sign under reversal. Positive divergence means that a random walk has positive expected flow at every step. Consequently, non-positive walks starting at a fixed vertex are rare; by reversal, positive walks ending at a fixed vertex are also rare. These are the two estimates that will control the two parts of the Markov operator.

LEMMA 2.1 (see [BLPS99, Proposition 2.1]). *There is a real function b on oriented Cayley edges satisfying*

$$(2.2) \quad b(x, y) = -b(y, x), \quad |b(x, y)| \leq 1, \quad \frac{1}{d} \sum_{s \in S} b(x, xs) = \eta \quad (x \in G).$$

PROOF. For a finite set $K \subset G$, form a network with a source, the vertices of K , and a sink. Give each source edge capacity $d\eta$, each orientation of an internal Cayley edge capacity one, and each boundary edge to the sink capacity one. A cut with source-side vertices $F \subseteq K$ has capacity

$$d\eta(|K| - |F|) + |\partial_E F| \geq d\eta|K|.$$

The max-flow/min-cut theorem gives a flow saturating all source edges. Subtract opposite internal flows, extend antisymmetrically across the boundary, and set the remaining values to zero. The resulting function has modulus at most one and divergence $d\eta$ on K . A diagonal limit along a finite exhaustion of G proves the assertion. $\square$

Let $\mathcal{W}_\ell = S^{\times \ell}$, $n_\ell = d^\ell$, and $s_w = s_1 \cdots s_\ell$ for $w = (s_1, \dots, s_\ell)$. Write Ω_ℓ for the set of length- ℓ walks. A walk $\omega = (x_0, \dots, x_\ell)$ has initial vertex $o(\omega) = x_0$, terminal vertex $t(\omega) = x_\ell$, and word $w(\omega) \in \mathcal{W}_\ell$, with $x_j = x_0 s_1 \cdots s_j$. Define

$$(2.3) \quad \Phi(\omega) = \sum_{j=1}^{\ell} b(x_{j-1}, x_j), \quad \Omega_\ell^- = \{\Phi \leq 0\}, \quad \Omega_\ell^+ = \{\Phi > 0\}.$$

Thus $\Omega_\ell = \Omega_\ell^- \sqcup \Omega_\ell^+$. Figure 1 shows the two kinds of walks inside a Cayley graph. For $E \subseteq \Omega_\ell$, use the endpoint fibres

$$E_x^o = \{\omega \in E : o(\omega) = x\}, \quad E_y^t = \{\omega \in E : t(\omega) = y\}.$$

We now show that the proportion of walks with non-positive total flow is negligible.

PROPOSITION 2.2. *For every $\ell \geq 1$ and $x, y \in G$,*

$$(2.4) \quad |(\Omega_\ell^-)_x^o| \leq n_\ell \beta^\ell, \quad |(\Omega_\ell^+)_y^t| \leq n_\ell \beta^\ell.$$

In particular, both quantities are at most $r_\ell := \lceil n_\ell \beta^\ell \rceil$, and $r_\ell/n_\ell \rightarrow 0$.

PROOF. For the uniform random walk starting at x , put $Z_j = b(X_{j-1}, X_j)$. Then $|Z_j| \leq 1$ and $\mathbb{E}[Z_j \mid \mathcal{F}_{j-1}] = \eta$. With $u = \operatorname{arctanh} \eta$, convexity gives

$$e^{-uz} \leq \cosh u - z \sinh u \quad (-1 \leq z \leq 1), \quad \mathbb{E}[e^{-uZ_j} \mid \mathcal{F}_{j-1}] \leq \beta.$$




FIGURE 1. A fragment of the 4-regular Cayley tree of F_2 . Orient edges away from a fixed end and put $b(x, y) = c$ in that direction, where $0 < c < 1$ is a constant; then the normalized divergence is $c/2$. The blue walk goes against the flow and belongs to Ω_2^- ; the orange walk follows the flow and belongs to Ω_2^+ . Dashed edges continue the tree.

Iterating and applying Markov's inequality yields

$$(2.5) \quad \mathbb{P}_x\{\Phi \leq 0\} \leq \mathbb{E}_x e^{-u\Phi} \leq \beta^\ell.$$

There are n_ℓ equally likely walks starting at x , proving the first estimate. Reversal maps positive walks ending at y bijectively to strictly negative walks starting at y , proving the second. Finally, $r_\ell/n_\ell \leq \beta^\ell + d^{-\ell} \rightarrow 0$. $\square$

For $\mu = d^{-1}\mathbf{1}_S$, the decomposition $\Omega_\ell = \Omega_\ell^- \sqcup \Omega_\ell^+$ gives

$$\mu^{*\ell}(x^{-1}y) = m_\ell^-(x, y) + m_\ell^+(x, y), \quad m_\ell^\pm(x, y) = \frac{|(\Omega_\ell^\pm)_x^o \cap (\Omega_\ell^\pm)_y^t|}{n_\ell}.$$

Proposition 2.2 says precisely that

$$(2.6) \quad \alpha(m_\ell^-) \leq \beta^\ell, \quad \gamma(m_\ell^+) \leq \beta^\ell, \quad \|\mu^{*\ell}\|_{T^1} \leq 2\beta^\ell.$$

3. Controlled matrix coefficients from free products

We construct positive matrix coefficients whose small partial sums have uniformly bounded operator norm, while their total squared tracial 2-norm can grow. The construction starts with averaging projections for the cyclic factors of a free product, passes to a finite quotient preserving short words, and compresses away a small spectral subspace. Fix the absolute constant $C_0 = 8\sqrt{2}$.

LEMMA 3.1. *Let $n \geq 2$ and $r \geq 1$ be integers. There are an integer $k \geq 1$ and positive matrices $B_1, \dots, B_n \in M_k(\mathbb{C})$ such that*

$$(3.1) \quad \left\| \sum_{i \in J} B_i \right\| \leq C_0 \quad (J \subseteq \{1, \dots, n\}, |J| \leq 2r), \quad \tau(B_i^2) \geq \frac{1}{2r} \quad (1 \leq i \leq n).$$

We recall two standard facts about free products used in the proof. For $n, q \geq 2$, let

$$(3.2) \quad \Gamma = H_1 * \dots * H_n, \quad H_i \cong C_q, \quad p_i = \frac{1}{q} \sum_{h \in H_i} \lambda_\Gamma(h) \in L\Gamma.$$

Here C_q is the cyclic group of order q , and τ_Γ is the canonical trace on $L\Gamma$. Each p_i is the orthogonal projection onto the H_i -invariant vectors, and

$$(3.3) \quad p_i^2 = p_i = p_i^*, \quad \tau_\Gamma(p_i^2) = \tau_\Gamma(p_i) = \frac{1}{q}.$$

LEMMA 3.2. For every nonempty $J \subseteq \{1, \dots, n\}$ with $s = |J| \leq q$,

$$(3.4) \quad \left\| \sum_{i \in J} p_i \right\| \leq \frac{s}{q} + \frac{2\sqrt{s(q-1)}}{q} + \frac{q-1}{q} < 4.$$

PROOF. Put $a_i = p_i - q^{-1}I$. The length-one free-product estimate of Ricard–Xu [RX06, Lemmas 2.1–2.3 and Corollary 2.4] gives, since $a_i = a_i^*$ and $\tau_\Gamma(a_i) = 0$,

$$\left\| \sum_{i \in J} a_i \right\| \leq \max_{i \in J} \|a_i\| + 2 \left(\sum_{i \in J} \tau_\Gamma(a_i^2) \right)^{1/2}.$$

Here $\|a_i\| = (q-1)/q$ and $\tau_\Gamma(a_i^2) = (q-1)/q^2$. Adding the scalar term s/q proves (3.4). $\square$

PROOF OF LEMMA 3.1. Set $q = 2r$, $p = \min(q, n)$ and $M = \binom{n}{p}$. Choose

$$(3.5) \quad t = \lceil \log_2(4qM) \rceil, \quad M2^{-t} \leq \frac{1}{4q}.$$

The group $\Gamma = H_1 * \dots * H_n$ is residually finite [Gru57, Theorem 4.1, Corollary (ii)], so there is a finite quotient F such that the quotient map is injective on all words in the syllable length at most t . On $\ell^2(F)$ define

$$(3.6) \quad P_i = \frac{1}{q} \sum_{h \in H_i} \lambda_F(\varphi(h)), \quad S_J = \sum_{i \in J} P_i \quad (|J| = p).$$

Thus P_i is a projection and $\tau(P_i) = 1/q$, where τ is the normalized trace in the regular representation of F .

All summands of $(\sum_{i \in J} p_i)^t$ map injectively to the quotient. Since the normalized trace in the regular representation of a finite group is the coefficient of the identity, Lemma 3.2 gives

$$(3.7) \quad \tau(S_J^t) = \tau_\Gamma \left(\left(\sum_{i \in J} p_i \right)^t \right) \leq 4^t.$$

For each p -element set J , let $Q_J = \mathbf{1}_{(8, \infty)}(S_J)$ be the spectral projection for eigenvalues greater than 8. Positivity gives $\tau(Q_J) \leq 8^{-t} \tau(S_J^t) \leq 2^{-t}$. Letting $E = \bigvee_{|J|=p} Q_J$ be the orthogonal projection onto the sum of their ranges, and putting $R = 1 - E$, we get $\delta := \tau(E) \leq M2^{-t} \leq \frac{1}{4q}$. The range of R lies in the spectral subspace of every S_J for eigenvalues at most 8. Hence for $|J| \leq p$ we get $RS_JR \leq 8R$. Define $B_i = \sqrt{2}(RP_iR)|_{R\ell^2(F)}$; the estimate on the norm follows.

It remains to estimate the tracial mass after compression. For $P = P_i$, we have

$$(3.8) \quad \tau((RPR)^2) = \tau(P) - 2\tau(PE) + \tau(PEPE) \geq \frac{1}{q} - 2\delta.$$

Indeed $0 \leq \tau(PE) \leq \tau(E)$ and $\tau(PEPE) = \tau((EPE)^2) \geq 0$. The normalized trace on the compressed space is $\tau' = \tau/\tau(R)$. Therefore

$$(3.9) \quad \tau'(B_i^2) = \frac{2\tau((RP_iR)^2)}{1 - \delta} \geq \frac{2(1/q - 2\delta)}{1 - \delta} \geq \frac{1}{q} = \frac{1}{2r}.$$

$\square$

4. Decomposing the matrix-weighted Markov operator

Fix ℓ and write $n = n_\ell$, $r = r_\ell$. Apply Lemma 3.1 with the words $w \in \mathcal{W}_\ell$ as labels, obtaining $B_w \in M_k(\mathbb{C})$. The unnormalized right Markov operator with coefficients (B_w) is defined as

$$(4.1) \quad \mathbf{M}_\ell = \sum_{w \in \mathcal{W}_\ell} \rho(s_w) \otimes B_w \in RG \overline{\otimes} M_k(\mathbb{C}).$$

For $E \subseteq \Omega_\ell$, let

$$(4.2) \quad \mathbf{M}_\ell(E) = \sum_{\omega \in E} (\delta_{o(\omega)} \delta_{t(\omega)}^*) \otimes B_{w(\omega)}.$$

These sums converge strongly: each vertex is the initial or terminal vertex of n walks, and all partial sums have norm at most $\sum_w \|B_w\|$. In particular $M_\ell(\Omega_\ell) = M_\ell$. The decomposition into non-positive and positive walks gives

$$(4.3) \quad T_\ell = M_\ell(\Omega_\ell^-), \quad Q_\ell = M_\ell(\Omega_\ell^+), \quad M_\ell = T_\ell + Q_\ell.$$

The sums retain the individual walks, including distinct walks with the same endpoints.

The next lemma converts sparsity at both endpoints into an operator norm bound. The factorization through the walk space applies Lemma 3.1 separately at initial and terminal vertices.

LEMMA 4.1. *Let $E \subseteq \Omega_\ell$ satisfy $|E_x^o|, |E_y^t| \leq 2r$ for all x, y . For any function $\varepsilon : E \rightarrow \mathbb{C}$ with $|\varepsilon| \leq 1$,*

$$(4.4) \quad \left\| \sum_{\omega \in E} \varepsilon(\omega) (\delta_{o(\omega)} \delta_{t(\omega)}^*) \otimes B_{w(\omega)} \right\| \leq C_0.$$

PROOF. On the walk space $\ell^2(E) \otimes \mathbb{C}^k$, define

$$\begin{aligned} (F\xi)(\omega) &= B_{w(\omega)}^{1/2} \xi(t(\omega)), \\ (D_\varepsilon \zeta)(\omega) &= \varepsilon(\omega) \zeta(\omega), \\ (S\zeta)(x) &= \sum_{\omega \in E_x^o} B_{w(\omega)}^{1/2} \zeta(\omega). \end{aligned}$$

At either endpoint, distinct walks have distinct words. Thus Lemma 3.1 gives

$$\|F\xi\|^2 = \sum_y \left\langle \xi(y), \left(\sum_{\omega \in E_y^t} B_{w(\omega)} \right) \xi(y) \right\rangle \leq C_0 \|\xi\|^2, \quad \|S\| \leq \sqrt{C_0}.$$

The operator in (4.4) equals $SD_\varepsilon F$, so its norm is at most C_0 . $\square$

We can now use sparsity to control the norms of rows and columns of T_ℓ as well as the its displacement under the conjugation action.

PROPOSITION 4.2. *With $U_k(g) = \lambda(g) \otimes I_k$, the operators (4.3) satisfy*

$$(4.5) \quad \sup_x \|\text{row}_x(T_\ell)\| \leq C_0, \quad \sup_y \|\text{col}_y(Q_\ell)\| \leq C_0, \quad \sup_g \|T_\ell - U_k(g)T_\ell U_k(g)^*\| \leq C_0.$$

PROOF. The operators $(\delta_x \delta_x^* \otimes I_k)T_\ell = M_\ell((\Omega_\ell^-)_x^o)$ and $Q_\ell(\delta_y \delta_y^* \otimes I_k) = M_\ell((\Omega_\ell^+)_y^t)$ involve at most r walks each. Lemma 4.1 gives the first two bounds.

Left translation acts on walks by $g\omega = (gx_0, \dots, gx_\ell)$ and preserves their words. Hence

$$U_k(g)T_\ell U_k(g)^* = M_\ell(g\Omega_\ell^-).$$

The difference is supported on $E = \Omega_\ell^- \triangle g\Omega_\ell^-$, with coefficients ± 1 . At an initial vertex, E is contained in $\Omega_\ell^- \cup g\Omega_\ell^-$; at a terminal vertex, use $E = \Omega_\ell^+ \triangle g\Omega_\ell^+ \subseteq \Omega_\ell^+ \cup g\Omega_\ell^+$. Proposition 2.2 therefore bounds both endpoint fibres by $2r$. Apply Lemma 4.1. $\square$

We now record the growing 2-norm estimate on our Markov operators.

PROPOSITION 4.3. *For the same operators,*

$$(4.6) \quad \|M_\ell\|_2^2 \geq \frac{n_\ell}{2r_\ell} \rightarrow \infty.$$

PROOF. By the L^2 Fourier decomposition (1.3),

$$(4.7) \quad \|M_\ell\|_2^2 = \sum_{s_w=s_u} \tau(B_w B_u) \geq \sum_w \tau(B_w^2) \geq \frac{n_\ell}{2r_\ell}.$$

The cross terms are nonnegative because $\tau(B_w B_u) = \tau(B_w^{1/2} B_u B_w^{1/2}) \geq 0$. Now apply Proposition 2.2. $\square$

5. Separation from the commutant and the bounded cocycle

The trace converts the local row and column bounds into a distance estimate from the commutant.

LEMMA 5.1. *Suppose $A \in RG \overline{\otimes} M_k(\mathbb{C})$ and $A = T + Q$, with $T, Q \in \mathcal{B}(H_k)$. For every $C \in RG \overline{\otimes} M_k(\mathbb{C})$,*

$$(5.1) \quad \|A\|_2 \leq \|\text{row}_e(T)\| + \|\text{col}_e(Q)\| + 2\|T - C\|.$$

Consequently,

$$(5.2) \quad \text{dist}(T, RG \overline{\otimes} M_k(\mathbb{C})) \geq \frac{1}{2} (\|A\|_2 - \|\text{row}_e(T)\| - \|\text{col}_e(Q)\|).$$

PROOF. Set $B = T - C$. Both C and $A - C$ lie in $RG \overline{\otimes} M_k(\mathbb{C})$. Equation (1.5) gives

$$\|C\|_2 \leq \|\text{row}_e(C)\| \leq \|\text{row}_e(T)\| + \|B\|, \quad \|A - C\|_2 \leq \|\text{col}_e(A - C)\| \leq \|\text{col}_e(Q)\| + \|B\|.$$

Apply the triangle inequality to $A = C + (A - C)$ in $L^2(RG \overline{\otimes} M_k(\mathbb{C}), \tau)$, and take the infimum over C . $\square$

For a unitary representation $U : G \rightarrow \mathcal{U}(H)$, an operator cocycle is a map $D : G \rightarrow \mathcal{B}(H)$ satisfying $D(gh) = D(g) + U(g)D(h)U(g)^*$. It is bounded if $\sup_g \|D(g)\| < \infty$, and inner if $D(g) = B - U(g)BU(g)^*$ for some $B \in \mathcal{B}(H)$. We now show that T_ℓ give rise to a non-inner bounded cocycle.

THEOREM 5.2. *For each ℓ , let k_ℓ and T_ℓ be obtained above, and put*

$$(5.3) \quad H = \bigoplus_{\ell \geq 1} (\ell^2(G) \otimes \mathbb{C}^{k_\ell}), \quad U(g) = \bigoplus_{\ell \geq 1} (\lambda(g) \otimes I_{k_\ell}).$$

Then

$$(5.4) \quad D(g) = \bigoplus_{\ell \geq 1} (T_\ell - (\lambda(g) \otimes I_{k_\ell})T_\ell(\lambda(g)^* \otimes I_{k_\ell}))$$

is a bounded non-inner operator cocycle for U , with $\sup_g \|D(g)\| \leq C_0$.

PROOF. Write $U_\ell(g) = \lambda(g) \otimes I_{k_\ell}$ and $D_\ell(g) = T_\ell - U_\ell(g)T_\ell U_\ell(g)^*$. Proposition 4.2 gives a uniformly bounded direct sum in (5.4). Direct expansion yields

$$D_\ell(gh) = D_\ell(g) + U_\ell(g)D_\ell(h)U_\ell(g)^*,$$

and hence the cocycle identity for D .

Suppose D were implemented by $B \in \mathcal{B}(H)$. Let $j_\ell : H_{k_\ell} \rightarrow H$ be the inclusion and set $B_\ell = j_\ell^* B j_\ell$. Since the summand projections commute with $U(g)$, compression gives

$$D_\ell(g) = B_\ell - U_\ell(g)B_\ell U_\ell(g)^*, \quad \|B_\ell\| \leq \|B\|.$$

It follows that $C_\ell = T_\ell - B_\ell \in RG \overline{\otimes} M_{k_\ell}(\mathbb{C})$. Applying Lemma 5.1 to $M_\ell = T_\ell + Q_\ell$ gives

$$\|M_\ell\|_2 \leq 2C_0 + 2\|B_\ell\| \leq 2C_0 + 2\|B\| \quad (\ell \geq 1).$$

This contradicts Proposition 4.3. Thus D is non-inner. $\square$

PROOF OF THE MAIN THEOREM. Let U and D be supplied by Theorem 5.2. Use the standard triangular construction; see [MO10, § 2]:

$$(5.5) \quad \pi(g) = \begin{pmatrix} U(g) & D(g)U(g) \\ 0 & U(g) \end{pmatrix} \quad \text{on } H \oplus H.$$

The cocycle identity gives $\pi(gh) = \pi(g)\pi(h)$, and $\sup_g \|\pi(g)\| \leq 1 + \sup_g \|D(g)\| \leq 1 + C_0 < 13$. If π were unitarizable, the invariant subspace $H \oplus 0$ would have an invariant complement, obtained by taking its orthogonal complement in an equivalent invariant Hilbert space inner product. The corresponding bounded projection would have the form

$$P = \begin{pmatrix} I & -B \\ 0 & 0 \end{pmatrix} \quad (B \in \mathcal{B}(H))$$

and commute with every $\pi(g)$. Comparing the upper-right entries gives

$$D(g)U(g) - BU(g) = -U(g)B, \quad D(g) = B - U(g)BU(g)^*.$$

This contradicts the non-innerness of D . Thus π is a uniformly bounded nonunitarizable representation of G . $\square$

References

- [Alp23] A. Alpeev, *Lamplighters over non-amenable groups are not strongly Ulam stable*, preprint, arXiv:2009.11738v3 (2023). arXiv:2009.11738v3.
- [BLPS99] I. Benjamini, R. Lyons, Y. Peres, and O. Schramm, *Group-invariant percolation on graphs*, Geom. Funct. Anal. **9** (1999), 29–66. Author-hosted text.
- [BF91] M. Bożejko and G. Fendler, *Herz-Schur multipliers and uniformly bounded representations of discrete groups*, Arch. Math. **57** (1991), 290–298.
- [Day50] M. M. Day, *Means for the bounded functions and ergodicity of the bounded representations of semi-groups*, Trans. Amer. Math. Soc. **69** (1950), no. 2, 276–291. doi:10.1090/S0002-9947-1950-0044031-5.
- [Dix50] J. Dixmier, *Les moyennes invariantes dans les semi-groupes et leurs applications*, Acta Sci. Math. (Szeged) **12** (1950), 213–227. Journal archive.
- [EM55] L. Ehrenpreis and F. I. Mautner, *Uniformly bounded representations of groups*, Proc. Nat. Acad. Sci. U.S.A. **41** (1955), no. 4, 231–233. doi:10.1073/pnas.41.4.231.
- [EM09] I. Epstein and N. Monod, *Nonunitarizable representations and random forests*, Int. Math. Res. Not. IMRN **2009** (2009), no. 22, 4336–4353. arXiv:0811.3422; doi:10.1093/imrn/rnp090.
- [GL09] D. Gaboriau and R. Lyons, *A measurable-group-theoretic solution to von Neumann’s problem*, Invent. Math. **177** (2009), no. 3, 533–540. arXiv:0711.1643; doi:10.1007/s00222-009-0187-5.
- [GGMT20] M. Gerasimova, D. Gruber, N. Monod, and A. Thom, *Asymptotics of Cheeger constants and unitarisability of groups*, J. Funct. Anal. **278** (2020), no. 11, 108457. arXiv:1801.09600; doi:10.1016/j.jfa.2019.108457.
- [GT21] M. Gerasimova and A. Thom, *On the isometrisability of group representations on p -spaces*, J. Operator Theory **86** (2021), no. 1, 51–60. arXiv:1901.07496; doi:10.7900/jot.2020jan22.2275.
- [Gru57] K. W. Gruenberg, *Residual properties of infinite soluble groups*, Proc. London Math. Soc. (3) **7** (1957), 29–62. doi:10.1112/plms/s3-7.1.29.
- [KS60] R. A. Kunze and E. M. Stein, *Uniformly bounded representations and harmonic analysis of the 2×2 real unimodular group*, Amer. J. Math. **82** (1960), no. 1, 1–62. doi:10.2307/2372876.
- [Mon06] N. Monod, *An invitation to bounded cohomology*, International Congress of Mathematicians, Vol. II, European Mathematical Society, Zürich, 2006, 1183–1211. doi:10.4171/022-2/56.
- [MO10] N. Monod and N. Ozawa, *The Dixmier problem, lamplighters and Burnside groups*, J. Funct. Anal. **258** (2010), no. 1, 255–259. arXiv:0902.4585; doi:10.1016/j.jfa.2009.06.029.
- [NT51] M. Nakamura and Z. Takeda, *Group representation and Banach limit*, Tohoku Math. J. (2) **3** (1951), no. 2, 132–135. doi:10.2748/tmj/1178245513.
- [OAI26] OpenAI, *Unitarizability implies amenability for discrete groups*, OpenAI Math, September 23, 2026. GitHub preprint.
- [Osi09] D. V. Osin, *L^2 -Betti numbers and non-unitarizable groups without free subgroups*, Int. Math. Res. Not. IMRN **2009** (2009), no. 22, 4220–4231. arXiv:0812.2093; doi:10.1093/imrn/rnp085.
- [Pis98] G. Pisier, *The similarity degree of an operator algebra*, Algebra i Analiz **10** (1998), no. 1, 132–186; English translation, St. Petersburg Math. J. **10** (1999), no. 1, 103–146. arXiv:math/9706211.
- [Pis05] G. Pisier, *Are unitarizable groups amenable?*, Infinite Groups: Geometric, Combinatorial and Dynamical Aspects, Progr. Math., vol. 248, Birkhäuser, Basel, 2005, 323–362. arXiv:math/0405282; doi:10.1007/3-7643-7447-0_8.
- [PS86] T. Pytlik and R. Szwarc, *An analytic family of uniformly bounded representations of free groups*, Acta Math. **157** (1986), no. 3–4, 287–309. doi:10.1007/BF02392596.
- [RX06] É. Ricard and Q. Xu, *Khintchine type inequalities for reduced free products and applications*, J. Reine Angew. Math. **599** (2006), 27–59. arXiv:math/0505302; doi:10.1515/CRELLE.2006.077.
- [SzN47] B. Sz.-Nagy, *On uniformly bounded linear transformations in Hilbert space*, Acta Sci. Math. (Szeged) **11** (1947), 152–157. Journal archive.
- [Tho15] A. Thom, *A remark about the spectral radius*, Int. Math. Res. IMRN **2015** (2015), no. 10, 2856–2864. arXiv:1306.1767; doi:10.1093/imrn/rnu018.
- [Tho16] A. Thom, *The expected degree of minimal spanning forests*, Combinatorica **36** (2016), no. 5, 591–600. arXiv:1306.0303; doi:10.1007/s00493-014-3160-x.
- [Ver23] I. Vergara, *The M_a -approximation property and unitarisability*, Proc. Amer. Math. Soc. **151** (2023), no. 3, 1209–1220. arXiv:2202.12198; doi:10.1090/proc/16204.
- [Wys88] J. Wysocański, *Characterization of amenable groups and the Littlewood functions on free groups*, Colloq. Math. **55** (1988), no. 2, 261–265. doi:10.4064/cm-55-2-261-265.