

MARN3002-Physics: 01 Introduction

Cesar B Rocha

UConn Avery Point, 2020

1 What is Physical Oceanography?

Oceanography is the scientific study of the ocean, including its physics, chemistry, biology and geology. Oceanographers aim to produce a mechanistic understanding of the ocean, with its myriad of multidisciplinary processes, as part of the Earth System. The process-oriented focus and its interdisciplinary nature are distinguishing characteristics of Oceanography.

The branch of Oceanography that focusses on ocean physics is known as *Physical Oceanography*. It studies why and how the ocean flows, what sets the distribution of ocean properties such as temperature and salinity, and how ocean motions influence (and sometimes control) ocean biogeochemistry and weather and climate.

Initially, Physical Oceanography emerged as a branch of Geography. Its principal products were maps of the distribution of ocean properties from observations collected in multi-year transoceanic expeditions, beginning in the 1800s. This oceanic branch of Geography developed into the subdiscipline *Descriptive Physical Oceanography*, which aims to quantitatively *describe* ocean motions. Historically, this description was developed with sparse ship-based observations, and more recently, with a combination of multi-platform observations and realistic computer simulations.

In the early 1900s, Physical Oceanography also began to be viewed as part of Geophysics, the branch of Physics concerned with Earth phenomena. The field of *Dynamical Physical Oceanography* emerged, approaching the understanding of ocean flows as a fundamental problem in Fluid Mechanics, and studying it with theoretical and computer models. A similar historical development occurred in Atmospheric Sciences, with the advent of Dynamical Meteorology in the late 1800s. In fact, Dynamical Meteorology and Dynamical Physical Oceanography are considered sister sciences, and most of their topics fall under the umbrella of Geophysical Fluid Dy-




Figure 1: Surface salinity map based on data collected aboard the German ship R/V Meteor in 1925-1927. Dots represent oceanographic stations, where in-situ measurements and water samples were collected. Maps like this were the main product of Ocean Geography, which later evolved into Descriptive Physical Oceanography. Note the high-salinity tongue in the Northeastern Atlantic, a footprint of the salty Mediterranean water that overflows into the Atlantic through the Strait of Gibraltar. Source: Adapted from Wüst (1978).

namics—the study of fluid flows influenced by rotation and stratification on Earth as well as on other planets and stars.




Figure 2: Sea-surface temperature in the Tropical Pacific from a state-of-the-art realistic ocean simulation. A realistic ocean numerical model is a computer code that solves the equations of the Dynamical Physical Oceanography, given initial and boundary conditions provided by observations. Note the texture of sea-surface temperature, associated with a number of ocean phenomena. Source: NOAA/GFDL.

While today we still talk about observational (descriptive) and theoretical (dynamical) oceanography, this division has been significantly eroded in the last couple of decades. The goal of a modern observational study is rarely to only survey a particular region; it generally aims to understand the mechanisms behind the observed phenomena. And dynamical process studies, both theoretical and numerical, are strongly influenced by ocean observations. Modern Physical Oceanography focusses on understanding the *physics of ocean processes*, their causes and consequences, using a combination of observations, theoretical fluid mechanics, and computer models.

Ocean physical processes can be local or regional or global, small or large, fast or slow. Figure 1 shows a space-time diagram of ocean phenomena, including bubbles, surface-gravity waves, tides, tsunamis, and global overturning (thermohaline) circulation. The range of scales of ocean variability is extraordinary, varying from from milliseconds and milimeters to thousands of years and tens of thousands of kilometers. Understanding these myriad phenomena and modeling their interactions is the main goal




Figure 3: Diagram of space and time scales of a number of ocean phenomena. In general, larger spatial scales are associated with longer time scales. Source: Talley et al., *Descriptive Physical Oceanography*.

of Physical Oceanography.

In this module of MARN3002, *we will study the basic physics of some of these phenomena* from observations and theory. I will emphasize *major concepts and ideas* of Physical Oceanography. Your expected learning outcome is a newfound *physical intuition* for how ocean motions work. Such an intuition is critical to learning the details of Physical Oceanography and its relationship to other areas of Oceanography and Climate. While Calculus is a pre-requisite for this class, I will strive to keep Math at a moderate level, focussing on what the relevant equations mean (what physics they contain) rather than deriving and solving them.

2 The physical/geographical setting

We begin by describing the container where ocean motions occur—planet Earth and its oceanic basins.




Figure 4: A spheroid representing Earth, with coordinates latitude (θ), longitude (λ) and radius r . The radius of the sphere (the distance from the center to any point on the sphere's surface) is the radius R . Source: Adapted from geokov.com.

2.1 Spherical Earth

Earth is an oblate spheroid, a sphere slightly flattened at the poles. This flattening is due to Earth's rotation around its vertical axis, and it implies that the polar radius is a bit (0.3 %) smaller than the equatorial radius. The rate of Earth's rotation is

$$\begin{aligned} |\vec{\Omega}| &= \frac{2\pi}{1 \text{ day}} + \frac{2\pi}{365 \text{ days}} \\ &= \frac{2\pi}{86400 \text{ sec}} + \frac{2\pi}{365 \times 86400 \text{ sec}} \approx 7.29 \times 10^{-5} \text{ radians/sec.} \end{aligned} \quad (1)$$

The direction of the rotation vector $\vec{\Omega}$ is perpendicular to the equator, pointing north (see figure 4).

It is natural to use a spherical coordinate system to describe the position on Earth. This consists of the distance from the center of the sphere (r) plus two angles θ and λ . Earth Scientists adopt the convention of measuring θ relative to the equator (the circle in the direction perpendicular to the rotation axis that splits Earth in half). The *latitude* θ varies from -90° (South

Pole) to $+90^\circ$ (North Pole). For mathematical calculations, we almost *always* measure angles in radians (rad). Recall that $360^\circ = 2\pi$ rad, so

$$\text{angle in radians} = \frac{\pi}{180} \times \text{angle in degrees}. \quad (2)$$

Thus, in radians, θ ranges from $-\pi/2$ to $\pi/2$. Longitude λ goes around the globe, varying from 0° (0) to 360° (2π). The reference longitude goes through the Greenwich, UK. An equivalent longitude range is from -180° ($-\pi$) to $+180^\circ$ ($+\pi$), with negative angles west of Greenwich. The curves along constant longitudes are called *meridians*. We refer to longitudinal direction as the *meridional direction*. The latitudinal circles are called parallels, and we refer to direction along parallels as the *zonal direction*.

The average radius of Earth is $R = 6371$ km. Hence one degree of latitude is equivalent to:

$$1^\circ \text{ latitude} = \frac{2\pi R}{360} = 111.20 \text{ km}. \quad (3)$$

Things are slightly different for longitude. The circumference around Earth at the constant latitude θ has a radius $R_\theta = R \cos \theta$. Thus

$$1^\circ \text{ longitude} = \frac{2\pi R}{360} \cos \theta. \quad (4)$$

Note that 1° longitude ≤ 111.20 km, with the equality holding at the equator, because $\cos 0 = 1$. The meridians converge poleward, with the distance between them diminishing with increasing latitude. For instance, at the latitude of Avery Point $\theta = 41.3179^\circ\text{N}$, 1° longitude = 83.52 km. Further poleward, at $\theta = 85^\circ$, 1° longitude = 9.67 km.

Distances in the ocean are also reported in *nautical miles* (Nm). One Nm is equivalent to one minute (i.e., one-sixtieth) of latitude:

$$1 \text{ Nm} = \frac{1^\circ}{60} = \frac{111.20}{60} \text{ km} = 1.853 \text{ km}. \quad (5)$$

2.2 Ocean topography

Figure 5 shows a map of the sea-floor elevation (the ocean bottom topography or bathymetry) on the global ocean. The values are negative, representing the distance from the sea surface to the ocean bottom. These data are from the ETOPO5 dataset, compiled by NOAA's National Center for Environmental Information; the resolution is 5 minutes of latitude (about 9 km).




Figure 5: World map of ocean topography and the location of a transatlantic bathymetric section along 41.3179° N, the latitude of Avery Point. Plotted with ETOPO5 dataset.

The ocean topography varies from shallow regions near the continents, ... through . The continental margin is an extension of the continent, wherein the seawater depth increases seaward. It has three main components: shelf, slope and elevation. These differentiated primarily by their inclination, the topographic gradient. The continental shelf starts at the coast and is the flatter regions of the continental margin. On the shelf, the bottom depth increases by about 1 m every 500 m. In other words, the topographic gradient is 1:500. Continental shelves vary in extension, from 10 km to 200 km, width an average with of 70 km. At the seaward edge of the shelf is the shelf break, where the topographic gradient increases significantly to 1:20. The average depth at the shelf break is 130 m. The slope is a narrow and steep region of the continental margin, where the bottom depth rapidly increases seaward. At about 3000-4000 m, the topographic gradient becomes gentler, given way to the continental elevation or rise, separating the continental margin from the abyss. By and large, the ocean abyss has rough topography, with oceanic mountain chains (such as mid-ocean ridges), isolated seamounts and trenches. The average depth of the ocean is 3734 m. The deepest documented region of the world ocean is the Mariana trench, in the Northwest Pacific, where depths reach more than 11000 m (11 km).

Figure 6 shows the cumulative probability distribution (CTD) the ocean topography in figure 5. In essence, a CDF is a fractional histogram, which




Figure 6: Cumulative probability function (CDF) of ocean topography, based on the ETOPO5 dataset.

measures the cumulative probability of the data. In figure 6, the cumulative probability starts at the deepest levels, moving to the right. The maximum probability is 1 (100 %), stating the obvious fact that 100 % of the ocean has depths below 0 m. The topography CDF shows that continental shelves (roughly regions shallower than 200 m) occupy less than 15 % of the world ocean, and the continental slope accounts for about 5 % the total area. Most of the ocean is deeper than 2000 %, although it is rarely deeper than 8000 m (0.05 %).

2.3 The aspect ratio

While most of the ocean is deep for our human scale, oceanic basins are extremely shallow on a planetary scale. Even the deepest point in the Mariana Trench is tiny in comparison with Earth's radius or circumference. The ocean is a *thin shell* in outer 0.08 % of Earth. In fact, when we plot long vertical sections of ocean topography, as in Figure 7, the vertical coordinate needs to be grossly exaggerated in order to render a detailed view of topographic features. From figure 7, the North Atlantic basin spans 60° of longitude at 41.3179° N. Thus, the North Atlantic width is

$$60^\circ \text{ longitude} \times \frac{111.20 \text{ km} \times \cos 41.3179^\circ}{1^\circ \text{ longitude}} \approx 5011 \text{ km}. \quad (6)$$




Figure 7: A transatlantic bathymetric section along 41.3179° N, the latitude of Avery Point, indicated in figure 5. Plotted with ETOPO5 dataset.

The average depth of the North Atlantic at this latitude is about 3000 m.

To calculate the relative size of the North Atlantic depth to its width, we calculate their ratio using the same *units*:

$$\frac{\text{depth}}{\text{width}} = \frac{3000 \text{ m}}{501110^3 \text{ m}} = 0.0006. \quad (7)$$

In other words, the North Atlantic thickness is about 0.06 % of its width. The relative size of thickness to width is called **aspect ratio**. The aspect ratio is a *non-dimensional number*. For most practical purposes, we are more interested in the order of magnitude of the terms. We thus define the aspect ratio as

$$\alpha \stackrel{\text{def}}{=} \frac{H}{L}, \quad (8)$$

where H and L is the order of magnitude (or scale) of the thickness and width. For the North Atlantic, $H \sim 10^3$ m and $L \sim 10^6$ m, and so $\alpha = 10^{-3}$.

As we will see, this tiny aspect ratio of the ocean is a powerful geometrical constraint on the ocean circulation, because it implies that many circulation features are much more wide than deep. For example, the Gulf Stream, the warm current that flows poleward off the US East coast, is about 200 km and 2000 m deep. The aspect ratio of the current is 10^{-2} . As we will see, small aspect ratio flows have tiny vertical velocities compared to horizontal velocities. The Gulf Stream flows laterally.