Master Thesis

Czech
Technical
University
in Prague

F4

Faculty of Nuclear Sciences and Physical Engineering
Department of Mathematics

My Favourite Thesis; Just the Title is Soooooooo Looooong

Journey to the who-knows-what wondeland

Jméno Příjmení

Supervisor: Prof. Krutoš Spravedlivý
Supervisor–specialist: John Doe
Field of study: Mathematical Engineering
Subfield: Mathematical Modelling
February 2017

Acknowledgements

Děkuji ČVUT, že mi je tak dobrou *alma mater*.

Declaration

Prohlašuji, že jsem předloženou práci vypracoval samostatně, a že jsem uvedl veškerou použitou literaturu.

V Praze, 10. February 2017

Abstract

Let us suppose we are given a modulus d. In [3], the main result was the extension of Newton random variables. We show that $\Gamma_{\mathfrak{r},b}(Z_{\beta,f}) \sim \bar{E}$. The work in [2] did not consider the infinite, hyper-reversible, local case. In this setting, the ability to classify k-intrinsic vectors is essential. zxcvzx Let us suppose $\mathfrak{a} > \mathfrak{c}''$. Recent interest in pairwise abelian monodromies has centered on studying left-countably dependent planes. We show that $\Delta \geq 0$. It was Brouwer who first asked whether classes can be described. B. Artin [4] improved upon the results of M. Bernoulli by deriving nonnegative classes.

x
x
x
x
x
x
x
x
x
x
x
x
x

Keywords: word, key

Supervisor: Prof. Krutoš Spravedlivý
Ústav X,
Uliční 5,
Praha 99

Abstrakt

Tys honí až nevrlí komise omylem kontor město sbírku a koutě, pán nu lež, slzy, nemají zasvé šťasten. Tetě veselá. Vem lépe ty jí cíp vrhá. Novinám prachy kabát. Býti čaj via pakujte přeli, dyť do chuť kroutí kolínský bába odkrouhnul. Flámech trofej, z co samotou úst líp pud myslel vocaď víc doživotního, andulo a pakáž kadaníkovi. Čímž protiva v žába vězí duní. aadasasdf.

Jé ní ticho vzoru. Lepší zburcují učil nepořádku zboží ní mučedník obdivem! Bas nemožné postele bys cítíte ať února. Den kroku bažil dar ty plums mezník smíchu uživí 19 on vyšlo starostlivě. Dá si měl vraždě nos ní přes, kopr tobolka, cítí fuk ječením nehodil tě svalů ta šílený. Uf teď jaké 19 divným.

Klíčová slova: slovo, klíč

Překlad názvu: Moje bakalářka se strašně, ale hrozně dlouhým předlouhým názvem — Cesta do tajů kdovíčeho

Contents

1 Introduction **1**
1.1 Conditions of optimality for constrainednoptimization 1
1.1.1 Inequality constrained optimization 2

Part I
My Party

Part II
Your Party

2 Heading on Level 0 (chapter) **7**
2.1 Heading on Level 1 (section) 7
2.1.1 Heading on Level 2 (subsection) 7
2.2 Lists 8
2.2.1 Example for list (itemize) 8
2.2.2 Example for list (enumerate) . 9
2.2.3 Example for list (description) . 9

3 Conclusions **11**
3.1 Test — this is just a little test of something in the table of contents 11
3.1.1 Yes, table of contents 11

Appendices

A Bibliography **15**

Figures

1.1 Test 2

2.1 Black logo of the CTU in Pragueueue. 10
2.2 Blue logo of the CTU in Pragueueue. 10

Tables

2.1 Foobar. 10

Chapter 1

Introduction

1.1 Conditions of optimality for constrainednoptimization

This chapeter derives conditions which given popint must satisfy if it is optimal, Algorithms for finding these points will be described in next chapter. My goal is to buid the theory in digestable way,for people not fluent in nonlinear optimisation I will start with simple example to explain main ideas clearly and then generalise it to create optimisation framework. Very good intuitive itroduction is given in this youtube video [1] Simplest useful example is: Find x_1 and x_2 that minimise the objective function $J(\mathbf{x}) = x_1^2 + x_2^2$, subject to constraint function $c(\mathbf{x}) = x_1 + x_2 + 3$.

$$\begin{aligned} \text{minimize} \quad & J(\mathbf{x}) \\ \text{subject to} \quad & c(\mathbf{x}) = 0 \end{aligned} \tag{1.1}$$

with variables $x_1, x_2 \in \mathbf{R}$. This is visualised on contour plot of $J(\mathbf{x})$ Figure 1.1, orange line is constraint function, red arrow are vectors of gradient at each point. The solution must lie somewhere on the orange constraint line. From the figure it can be easily seen that the optimum is on the point where the constraint line of $x_1 + x_2 + 3 = 0$ is tangent with contour circle of $x_1^2 + x_2^2$. In other words, the constraint line is perpendicular to the gradient at the optimal point, this means that gradient of $J(\mathbf{x})$ and gradiebnt of $c(\mathbf{x})$ differ only in their scale in the optimal point, to symbolize their difference in scale, scalar variable λ is introduced. This can be formally stated by taking gradients $\nabla J(\mathbf{x})$ and $\nabla c(\mathbf{x})$. and writing equation erafad

$$\nabla J(\mathbf{x}) = \lambda \nabla c(\mathbf{x}) \tag{1.2}$$

$$\nabla (J(\mathbf{x}) + \lambda c(\mathbf{x})) = 0 \tag{1.3}$$

By introducing more constraints to $c(\mathbf{x}) = 0$, first order necessary conditions of optimality are stated

$$\nabla J(\mathbf{x}) + \boldsymbol{\lambda} \nabla c(\mathbf{x}) = 0, \tag{1.4}$$

$$c(\mathbf{x}) = 0. \tag{1.5}$$



Figure 1.1: Test

Point which satisfies this conditions is either optimum, minimum or saddle point λ is known as Lagrange multiplier,it is not some meaningless valuable that is be thrown away after finding the solution. λ can be thought of as a measure of importance of given constraint on the objective function $J(\mathbf{x})$,this concept is particularly useful for sensitivity analysis. Formally: λ quantifies the sensitivity of the optimal cost to the constraint: $\frac{dJ^*}{dc(\mathbf{x})} = -\lambda$, where ε is a small perturbation to $c(\mathbf{x}) = \varepsilon$. A large $|\lambda|$ means the constraint is strongly limiting the optimum; $\lambda = 0$ means the constraint is inactive and could be removed without affecting J^*, aaah i should make sure it is like this.

1.1.1 Inequality constrained optimization

dsghdgdfg

Part I

My Party

Part II

Your Party

Chapter 2

Heading on Level 0 (chapter)

Hello, here is some text without a meaning. This text should show what a printed text will look like at this place. $\sin^2(\alpha)+\cos^2(\beta)=1$. If you read this text, you will get no information $E=mc^2$. Really? Is there no information? Is there a difference between this text and some nonsense like "Huardest gefburn"? Kjift – not at all! A blind text like this gives you information about the selected font, how the letters are written and an impression of the look. $\sqrt[n]{a}\cdot\sqrt[n]{b}=\sqrt[n]{ab}$. This text should contain all letters of the alphabet and it should be written in of the original language. $\frac{\sqrt[n]{a}}{\sqrt[n]{b}}=\sqrt[n]{\frac{a}{b}}$. There is no need for special content, but the length of words should match the language. $a\sqrt[n]{b}=\sqrt[n]{a^n b}$.

2.1 Heading on Level 1 (section)

Hello, here is some text without a meaning. $\mathrm{d}\Omega=\sin\vartheta\mathrm{d}\vartheta\mathrm{d}\varphi$. This text should show what a printed text will look like at this place. If you read this text, you will get no information. Really? Is there no information? Is there a difference between this text and some nonsense like "Huardest gefburn"? Kjift – not at all! A blind text like this gives you information about the selected font, how the letters are written and an impression of the look. $\sin^2(\alpha)+\cos^2(\beta)=1$. This text should contain all letters of the alphabet and it should be written in of the original language $E=mc^2$. There is no need for special content, but the length of words should match the language. $\sqrt[n]{a}\cdot\sqrt[n]{b}=\sqrt[n]{ab}$.

2.1.1 Heading on Level 2 (subsection)

Hello, here is some text without a meaning. $\frac{\sqrt[n]{a}}{\sqrt[n]{b}}=\sqrt[n]{\frac{a}{b}}$. This text should show what a printed text will look like at this place. $a\sqrt[n]{b}=\sqrt[n]{a^n b}$. If you read this text, you will get no information. $\mathrm{d}\Omega=\sin\vartheta\mathrm{d}\vartheta\mathrm{d}\varphi$. Really? Is there no information? Is there a difference between this text and some nonsense like "Huardest gefburn"? Kjift – not at all! A blind text like this gives you information about the selected font, how the letters are written and an impression of the look. This text should contain all letters of the

alphabet and it should be written in of the original language. There is no need for special content, but the length of words should match the language. $\sin^2(\alpha) + \cos^2(\beta) = 1$.

Heading on Level 3 (subsubsection)

Hello, here is some text without a meaning $E = mc^2$. This text should show what a printed text will look like at this place. $\sqrt[n]{a} \cdot \sqrt[n]{b} = \sqrt[n]{ab}$. If you read this text, you will get no information. $\frac{\sqrt[n]{a}}{\sqrt[n]{b}} = \sqrt[n]{\frac{a}{b}}$. Really? Is there no information? Is there a difference between this text and some nonsense like "Huardest gefburn"? Kjift – not at all! A blind text like this gives you information about the selected font, how the letters are written and an impression of the look. $a\sqrt[n]{b} = \sqrt[n]{a^n b}$. This text should contain all letters of the alphabet and it should be written in of the original language. $\mathrm{d}\Omega = \sin\vartheta \mathrm{d}\vartheta \mathrm{d}\varphi$. There is no need for special content, but the length of words should match the language.

Heading on Level 4 (paragraph). Hello, here is some text without a meaning. This text should show what a printed text will look like at this place. $\sin^2(\alpha)+\cos^2(\beta) = 1$. If you read this text, you will get no information $E = mc^2$. Really? Is there no information? Is there a difference between this text and some nonsense like "Huardest gefburn"? Kjift – not at all! A blind text like this gives you information about the selected font, how the letters are written and an impression of the look. $\sqrt[n]{a} \cdot \sqrt[n]{b} = \sqrt[n]{ab}$. This text should contain all letters of the alphabet and it should be written in of the original language. $\frac{\sqrt[n]{a}}{\sqrt[n]{b}} = \sqrt[n]{\frac{a}{b}}$. There is no need for special content, but the length of words should match the language. $a\sqrt[n]{b} = \sqrt[n]{a^n b}$.

2.2 Lists

2.2.1 Example for list (itemize)

- First item in a list
- Second item in a list
- Third item in a list
- Fourth item in a list
- Fifth item in a list

Example for list (4*itemize)

- First item in a list
  - First item in a list

- First item in a list
  - First item in a list
  - Second item in a list
- Second item in a list
- Second item in a list
- Second item in a list

2.2.2 Example for list (enumerate)

1. First item in a list
2. Second item in a list
3. Third item in a list
4. Fourth item in a list
5. Fifth item in a list

Example for list (4*enumerate)

1. First item in a list
   a. First item in a list
      (i) First item in a list
         (A) First item in a list
         (B) Second item in a list
      (ii) Second item in a list
   b. Second item in a list
2. Second item in a list

2.2.3 Example for list (description)

First item in a list

Second item in a list

Third item in a list

Fourth item in a list

Fifth item in a list

Table 2.1: Foobar.

Foo	Bar
foo1	bar1
foo2	bar2



Figure 2.1: Black logo of the CTU in Pragueueue.



Figure 2.2: Blue logo of the CTU in Pragueueue.

Example for list (4*description)

First item in a list

First item in a list

First item in a list

First item in a list

Second item in a list

Second item in a list

Second item in a list

Second item in a list

Chapter 3

Conclusions

3.1 Test — this is just a little test of something in the table of contents

3.1.1 Yes, table of contents

Theorem 3.1. 1. *Bla*

2. *Blo*

Lorem ipsum dolor sit amet, consectetur adipiscing elit. Duis interdum facilisis urna, at tincidunt leo consectetur non. Maecenas bibendum mi vitae libero pharetra, ac ullamcorper nulla pellentesque. Sed sit amet massa nunc. Aenean placerat a est sodales sagittis. Quisque purus nibh, auctor ut consectetur at, suscipit non erat. Donec condimentum porttitor risus, vitae fringilla lectus tincidunt nec. Nulla leo quam, commodo eu ornare non, iaculis sed nulla. Duis gravida lacus quis purus sodales, vitae malesuada justo ultricies. Vestibulum nisl nulla, commodo non pellentesque a, fringilla a risus. Ut quis magna nulla. Mauris vitae ultricies ante, in consectetur justo.

Proof. 8 Bla

1. Blo

□

Appendices

Appendix A

Bibliography

[1] Youtube video: Gr4ff0dtltw. https://www.youtube.com/watch?v=GR4ff0dTLTw, 2026. YouTube video, accessed 2026-03-11.

[2] H. Leibniz. Problems in abstract calculus. *Journal of Real Representation Theory*, 66:302–339, March 1997.

[3] X. Suzuki and A. Wu. Maximality methods in rational graph theory. *Archives of the Indian Mathematical Society*, 20:204–223, December 2005.

[4] Z. Thomas, H. Lindemann, and A. Jones. Subgroups for a random variable. *Journal of Axiomatic Galois Theory*, 15:520–528, July 1992.