

$$SEP(p(\sigma x.WN(x)) + p(\sigma x.WIJ(x)))$$

we avoid this problem.

## ANSWERS TO EXERCISES

### CHAPTER ONE

#### Exercise 1

(a) If  $\varphi$  is not true in  $M$ , then for some  $g$ :  $M \not\models \varphi$ . Then (by the lemma) for all  $g$ :  $M \not\models \varphi$ , hence  $\varphi$  is false in  $M$ . Similarly, if  $\varphi$  is not false in  $M$ , by the same argument  $\varphi$  is true in  $M$ . Hence  $\varphi$  cannot be both not true in  $M$  and not false in  $M$ . So  $\varphi$  is either true or false in  $M$ .

(b) We know that for some  $d \in D$ :  $d \in i(P)$  and for some  $d' \in D$ :  $d' \notin i(P)$ . Hence we know that for some  $g$  (a  $g$  where  $g(x) = d$ )  $M \models P(x)[g]$  and that for some  $g'$  (a  $g'$  where  $g'(x) = d'$ )  $M \not\models P(x)[g']$ . The first tells us that  $P(x)$  is not true, the second that  $P(x)$  is not false, so  $P(x)$  is undefined. Of course, the same argument applies to  $\neg P(x)$ .

(c) Take any assignment  $g$ .  $M \models P(x) \vee \neg P(x)[g]$  iff  $M \models P(x)[g]$  or  $M \models \neg P(x)[g]$  iff  $g(x) \in i(P)$  or  $g(x) \notin i(P)$ . The latter is of course true (for any  $g$ ). So for all  $g$ :  $M \models P(x) \vee \neg P(x)[g]$ , so  $P(x) \vee \neg P(x)$  is true in  $M$  (in any  $M$ , for that matter). A similar argument tells you that  $P(x) \wedge \neg P(x)$  is false in  $M$ , because there is no  $g$  such that  $g(x) \in i(P)$  and  $g(x) \notin i(P)$ .

So you see that, although  $P(x)$  and  $\neg P(x)$  themselves are undefined,  $P(x) \vee \neg P(x)$  and  $P(x) \wedge \neg P(x)$  are not.

#### Exercise 2

$\varphi[x]$  and  $\forall x\varphi[x]$  are logically equivalent if for every  $M$ :  $M \models \varphi[x]$  iff  $M \models \forall x\varphi[x]$ .  $M \models \varphi[x]$  iff for every  $g$ :  $M \models \varphi[x][g]$  iff for every  $g$  and every  $d \in D$ :  $M \models \varphi[x][g_x^d]$  iff for every  $g$ :  $M \models \forall x\varphi[x][g]$  iff  $M \models \forall x\varphi[x]$ .

#### Exercise 3

1. Every atomic formula has an equal number of left and right brackets.
2. Assume  $\varphi$  has an equal number of left and right brackets.  $\neg\varphi$

doesn't introduce any brackets, so  $\neg\varphi$  has the same number of left and right brackets. The same holds for  $\exists x\varphi$ .

3. Assume that each of  $\varphi$  and  $\psi$  has an equal number of left and right brackets.  $(\varphi \wedge \psi)$  introduces one left and one right bracket, so  $(\varphi \wedge \psi)$  has an equal number of left and right brackets as well.

#### Exercise 4

(a) Suppose  $\Delta$  is not deductively closed. This means that for some  $\varphi$ :  $\Delta \vdash \varphi$  but  $\varphi \notin \Delta$ . But then  $\Delta \cup \{\varphi\}$  is consistent, and since  $\Delta \subseteq \Delta \cup \{\varphi\}$  and  $\Delta \neq \Delta \cup \{\varphi\}$ ,  $\Delta$  is not maximally consistent.

(b) Suppose that for some  $\varphi$ :  $\Delta \not\vdash \varphi$  and  $\Delta \not\vdash \neg\varphi$ . Since  $\Delta \not\vdash \neg\varphi$ ,  $\Delta \cup \{\varphi\}$  is consistent. But then, because  $\Delta$  is maximally consistent,  $\varphi \in \Delta$ . If  $\varphi \in \Delta$  then  $\Delta \vdash \varphi$ . Contradiction.

#### Exercise 5

##### Soundness

- (1) if  $\Delta \vdash \varphi$  then  $\Delta \models \varphi$
- (2) if  $\Delta$  has a model then  $\Delta$  is consistent

(2)  $\rightarrow$  (1)

Suppose  $\Delta \not\models \varphi$ . If  $\Delta \not\models \varphi$ , Then for some model  $M$ :  $M \models \Delta$  and  $M \not\models \varphi$ , hence  $\Delta \cup \{\neg\varphi\}$  has a model. Then (by 2)  $\Delta \cup \{\neg\varphi\}$  is consistent, but then  $\Delta \not\models \varphi$ .

(1)  $\rightarrow$  (2)

Suppose  $\Delta$  is inconsistent. Then  $\Delta \vdash \perp$ . Hence (by 1)  $\Delta \models \perp$ . But that means that  $\Delta$  does not have a model.

##### Completeness

- (1) if  $\Delta \models \varphi$  then  $\Delta \vdash \varphi$
- (2) if  $\Delta$  is consistent then  $\Delta$  has a model

(2)  $\rightarrow$  (1)

Suppose  $\Delta \not\models \varphi$ . If  $\Delta \not\models \varphi$ , then  $\Delta \cup \{\neg\varphi\}$  is consistent. Then (by 2)  $\Delta \cup \{\neg\varphi\}$  has a model, but then  $\Delta \not\models \varphi$ .

(1)  $\rightarrow$  (2)

Suppose  $\Delta$  doesn't have a model. Then  $\Delta \models \perp$ . Hence (by 1)  $\Delta \vdash \perp$ , so  $\Delta$  is inconsistent.

#### Exercise 6

Suppose every finite subset of  $\Delta$  has a model. Then (by completeness) every finite subset of  $\Delta$  is consistent. Then (by definition of consistency)  $\Delta$  is consistent. Then (by completeness)  $\Delta$  has a model.

#### Exercise 7

*Fact 3:* If  $\varphi$  were to define infinity then  $\neg\varphi$  would define finiteness.

#### Exercise 8

*Fact 4:* Our set of sentences  $\{\varphi_1, \varphi_2, \dots\}$  defines infinity.

#### Exercise 9

$2 + 2 = 4$  means  $SS0 + SS0 = SSSS0$ .

$SS0 + SS0 =$  (axiom 4)  $S(SS0 + S0) =$  (axiom 4)  $SS(SS0 + 0) =$  (axiom 3)  $SSSS0$ .

#### Exercise 10

The statements expressing that  $\subseteq$  is reflective and transitive are just logical tautologies. Antisymmetry follows from logic and extensionality. If  $\forall x[x \in A \rightarrow x \in B]$  and  $\forall x[x \in B \rightarrow x \in A]$  then  $\forall x[x \in A \leftrightarrow x \in B]$ , hence with extensionality  $A = B$ .

#### Exercise 11

Given set  $A$ . Comprehension and extensionality tell you that the unique set  $\{a \in A : a \neq a\}$  exists, this is of course  $\emptyset$ .

## Exercise 12

ZF is a first order theory. If ZF is consistent, then any subset of ZF is obviously also consistent, so  $ZF - \{F\}$  is consistent.

## Exercise 13

Well, nothing because, as I said, we can prove it from the other axioms. But that is not what I meant, suppose we leave out both substitution and separation, what happens? The theory is not going to be inconsistent, because the same argument as above applies. You are just for a lot of sets unable to prove in the resulting theory that they exist.

## CHAPTER TWO

## Exercise 1

(a)  $\langle \mathbf{E}, < \rangle$  and  $\langle \mathbf{0}, < \rangle$  are substructures of  $\langle \mathbf{N}, < \rangle$ .  $\langle \mathbf{E}, <, + \rangle$  is a substructure of  $\langle \mathbf{N}, <, + \rangle$ , But not  $\langle \mathbf{0}, <, + \rangle$ , because  $\mathbf{0}$  is not closed under  $+$ . Also not  $\langle \mathbf{E}, < \rangle$  and  $\langle \mathbf{0}, < \rangle$ , because they are of the wrong type.

(b) Say that the type of  $\mathbf{A}$  is a subtype of the type of  $\mathbf{B}$  if  $\mathbf{B}$  has relations, operations, special elements corresponding to all relations, operations and special elements of  $\mathbf{A}$ , but maybe more. Let  $\mathbf{B}'$  be the result of disregarding in  $\mathbf{B}$  all the relations, operations and special elements that do not correspond to any in  $\mathbf{A}$ . Then  $\mathbf{A}$  is a generalized substructure of  $\mathbf{B}$  iff  $\mathbf{A}$  is a substructure of  $\mathbf{B}'$ .

Besides the ones that are already substructures,  $\langle \mathbf{E}, < \rangle$  is a generalized substructure of  $\langle \mathbf{N}, <, + \rangle$ .

## Exercise 2

We know already that  $\lambda x.g(f(x))$  is a function from  $A$  into  $C$ . Suppose  $R_A(a, a')$ . Then, because  $f$  is a homomorphism,  $R_B(f(a), f(a'))$ , and because  $g$  is a homomorphism,

$$R_C(g(f(a)), g(f(a'))).$$

$$g(f(a *_A a'))$$


$$= g(f(a) *_B f(a')) \text{ (because } f \text{ is a homomorphism)}$$

$$= g(f(a)) *_C g(f(a')) \text{ (because } g \text{ is a homomorphism).}$$

Of course,  $g(f(a)) = a$ , for the special elements.

## Exercise 3

1.  $h$  is an isomorphism between  $A$  and  $h(A)$ .



2.

There is no relation between  $a$  and the other elements in  $A$ , so, trivially, the indicated function preserves the relation. The function is one-one, so indeed it is a bijective homomorphism. But not an isomorphism, because the relation between  $d$  and  $e$  is not preserved in  $A$ .

## Exercise 4

(A) Relational structures

- (a) not a homomorphism, so it is none of the others either.
- (b) only a homomorphism.
- (c) again only a homomorphism
- (d) epimorphism
- (e) embedding
- (f) embedding
- (g) isomorphism
- (h) isomorphism
- (i) automorphism (the trivial automorphism, given by identity)
- (j) a non-trivial automorphism
- (k) isomorphism. Strictly speaking not an automorphism, because the first structure is  $\langle A, \leq \rangle$  and the second  $\langle A, \geq \rangle$ . Of course, up to duality those two structures are the same, so in this sense it is an automorphism.

(l) homomorphism

(B) Algebras.

(a) of course also not a homomorphism.

(b) homomorphism

(c) not a homomorphism:  $b \wedge c = a$ ,  $f(b \wedge c) = f$ , but  $f(b) \wedge f(c) = g \wedge g = g$

(d) not a homomorphism: for the same reason as under (c).

*Remark:* note the difference between (c) and (d): (c) does preserve  $\vee$  (it is called a join-homomorphism), but not  $\wedge$ . (d) preserves neither  $\vee$  nor  $\wedge$ .

(e) embedding


(f) not a homomorphism  $c \vee d = e$ , but  $f(c \vee d) = l$  and  $f(c) \vee f(d) = k$ .

(g) and (h) are isomorphisms; (i) and (j) are automorphisms; (k) is again an isomorphism (and an automorphism up to duality).

(l) homomorphism

### Exercise 5

(a)



(b) Suppose for some  $a$ :  $R(a, a)$ , then of course  $R(a, a)$  and  $R(a, a)$ , and hence for some  $a, b$  ( $a = b$ ):  $R(a, b)$  and  $R(b, a)$ .

(c) Suppose  $R(a, b)$  and  $R(b, a)$ . Then, by transitivity,  $R(a, a)$ , which contradicts irreflexivity.

(d)




### Exercise 6

1.  $\leq$  is reflexive,  $a \geq a$  iff  $a \leq a$ , so  $\geq$  is reflexive.

2. Assume  $a \geq b$  and  $b \geq c$ . Then we know that  $c \leq b$  and  $b \leq a$ .  $\leq$  is transitive, so  $c \leq a$ , hence  $a \geq c$ .

3. Assume  $a \geq b$  and  $b \geq a$ . Then  $b \leq a$  and  $a \leq b$  and hence  $a = b$ .

### Exercise 7



### Exercise 8

1. Suppose  $b \in [a]_{\approx}$ . i.e.  $b \in \{c: c \approx a\}$ , so  $b \approx a$ . Then  $a \approx b$  (because  $\approx$  is symmetric), hence  $a \in [b]_{\approx}$ . Now, if  $c \in [a]_{\approx}$ , then because  $a \in [b]_{\approx}$  and  $\approx$  is transitive,  $c \in [b]_{\approx}$ . The same, the other way round, of course: if  $c \in [b]_{\approx}$ , then  $c \in [a]_{\approx}$ , because  $b \in [a]_{\approx}$  and  $\approx$  is transitive.

2. Since  $\approx$  is reflexive,  $a \approx a$ , so  $a \in [a]_{\approx}$  (for any  $a$ ).

3. Suppose that  $[a]_{\approx} \cap [b]_{\approx} \neq \emptyset$ , say,  $c \in [a]_{\approx}$  and  $c \in [b]_{\approx}$ . Then, by (1)  $[a]_{\approx} = [c]_{\approx}$  and  $[b]_{\approx} = [c]_{\approx}$ , and hence  $[a]_{\approx} = [b]_{\approx}$ .

4. Suppose  $a \in A$ . Then  $a \in [a]_{\approx}$  and hence  $a \in \bigcup \{[b]_{\approx} : b \in A\}$ . Suppose  $a \in \bigcup \{[b]_{\approx} : b \in A\}$ . Then for some  $[b]_{\approx}$  ( $b \in A$ ):  $a \in [b]_{\approx}$ .  $[b]_{\approx} = \{a \in A : a \approx b\}$ , hence by definition of  $[b]_{\approx}$   $a \in A$ .

## Exercise 9

Suppose  $X \leq Y$  and  $Y \leq X$ . Then for some  $a \in X$ ,  $b \in Y$ :  $aRb$  and for some  $b' \in Y$ ,  $a' \in X$ :  $b'Ra'$ . We then know the following:

$aRb$ ;  $bRb'$  (by def. of  $Y$ ),  $b'Ra'$ ,  $a'Ra$  (by def. of  $X$ ), hence, by transitivity,  $bRa$ , but that means (by def. of  $X$ ) that  $b \in X$ . Hence  $X \cap Y \neq \emptyset$ , and thus  $X = Y$  (partition).

## Exercise 10

Proof of the lemma. (a) Suppose  $\exists c: c \leq b$  and  $d(a) = d(c)$ . Obviously  $d(b) = d(c) + |\{d: c < d \leq b\}|$ , so  $d(c) \leq d(b)$  and hence  $d(a) \leq d(b)$ .

(b) Suppose  $d(a) \leq d(b)$ . Clearly the only interesting situation is the one where  $a$  and  $b$  are on different branches, where they are incomparable. So suppose that to be the case and say,  $d(a) = m$  and  $d(b) = n$ , with  $m \leq n$ . So,  $|\{c: c \leq b\}| = n - 1$ . But, of course, we can just count back in  $\{c: c \leq b\}$ , until we find an element  $c \in \{c: c \leq b\}$ , such that  $|\{d: d \leq c\}| = m - 1$ :  $c \leq b$  and  $d(c) = d(a)$ .

*Proof of the theorem.*

*Transitivity.* Suppose  $X \leq Y$  and  $Y \leq Z$ . Then for some  $a \in X$ ,  $b \in Y$ ,  $a \leq b$  and for some  $b' \in Y$ ,  $c \in Z$ :  $b' \leq c$ . We know that  $a \leq b$ , hence  $d(a) \leq d(b)$ ; we know that  $d(b) = d(b')$ , hence we know that  $d(a) \leq d(b')$ . Then, by the lemma, we know that there is an  $a' \in X$ , such that  $a' \leq b'$ . By transitivity,  $a' \leq c$ , hence indeed  $X \leq Z$ .

*Antisymmetry.*

Suppose  $X \leq Y$  and  $Y \leq X$ . That means that for some  $a \in X$   $b \in Y$   $a \leq b$  and for some  $b' \in Y$   $a' \in X$   $b' \leq a'$ . We know (by def.) that  $d(b) = d(b')$  and  $d(a) = d(a')$ . Since  $a \leq b$  and  $b' \leq a'$ , it then follows that  $d(a) \leq d(b)$  and  $d(b) = d(b') \leq d(a') = d(a)$ , so  $d(b) \leq d(a)$ , hence  $d(a) = d(b)$ , and thus, by definition of  $X$ ,  $b \in X$ . Hence,  $X \cap Y \neq \emptyset$  and hence  $X = Y$ .

*Linearity:*  $X \leq Y$  or  $Y \leq X$ .

Suppose  $X \neq Y$ . Then for all  $a \in X$ , all  $b \in Y$ :  $a \not\leq b$ . We know  $X$  and  $Y$  are non-empty, so take  $a \in X$  and  $b \in Y$ .

$a \not\leq b$ . Now suppose that  $d(a) \leq d(b)$ . Then for some  $a' \in X$ :  $a' \leq b$

(by the lemma), contradicting the assumption. So we know that  $d(b) < d(a)$ . But then, again by the lemma, we know that for some  $b' \in Y$ :  $b' < a$ , and hence we know that  $Y \leq X$ .

## Exercise 11

We know that in a semitree for every  $a$  and  $b$ , there is a unique latest point, earlier than both  $a$  and  $b$ , call this  $\bigcirc(a, b)$ , the origin of  $a$  and  $b$ . If  $x \leq y$ , let us define  $d(y, x)$ , the distance from  $y$  to  $x$  as:  $d(y, x) = |\{z: x \leq z \text{ and } z \leq y\}| - 1$ .

Since  $\bigcirc(a, b) \leq a$  and  $\bigcirc(a, b) \leq b$ ,  $d(a, \bigcirc(a, b))$  and  $d(b, \bigcirc(a, b))$  are both well defined. So we can define equidistance as:


$$\approx := \lambda a \lambda b. d(a, \bigcirc(a, b)) = d(b, \bigcirc(a, b))$$

This is obviously going to give us the right result.

## Exercise 12

Assume that for some  $b$ :  $a \leq b$  and  $a' \leq b$ . Look at  $\{c: c \leq b\}$ . This is, by definition a linear order, and since both  $a$  and  $a'$  are in it, it follows that  $a \leq a'$  or  $a' \leq a$ .


## Exercise 13



(b) Take any two minimal elements  $a$  and  $b$ . Either  $a \leq b$  or  $b \leq a$  (by linearity). But  $a < b$  is false, because  $b$  is minimal, and similarly  $b < a$  is false, hence  $a = b$ .

## Exercise 14


(a) The following poset is an example:



So, the poset can be a collection of linear orders.

(b)  $\forall x \forall y \exists z [(z \leq x \wedge z \leq y) \vee (x \leq z \wedge y \leq z)]$ . This excludes the previous example: take any  $a$  and  $b$ : they have a common predecessor  $c$  or a common successor, say a predecessor  $c$ . Then, by no branching to the future  $a \leq b$  or  $b \leq a$ . Similarly, if they have a common successor, you use no branching to the past.

(c)



(d) A treelike structure is a strict partial order, which is not branching to the past and where every two elements have a common predecessor.

#### Exercise 15

(a) Because of the following possibility:



$a$  has a direct successor, but the branch between  $a$  and  $b$  may be dense.

(b)  $\langle T, < \rangle$  is discreet iff

$$\forall a \forall b [a < b \rightarrow \exists c [a < c \leq b \wedge \neg \exists d [a < d < c]]]$$

#### Exercise 16


(a)  $T$  itself is a linear order in  $T$ , so it has a minimal element.

(b) I. Suppose  $T$  is not discreet to the future. Then there is an element  $a$  that has successors, but no direct successor. Look at  $\{b: a < b\}$ , the set of all  $a$ 's successors. This set doesn't have a minimal element (because that would be  $a$ 's direct successor. This contradicts well foundedness.

II. The ordinal numbers ordered by  $<$  are a good example: this structure is linear, continuing to the future and well founded.  $\mathbb{N}$  with a copy of  $\mathbb{N}$  behind it is another example. These structures are well founded all right, but not discreet, because  $\omega$  in the ordinal numbers and the second 0 in  $\mathbb{N} * \mathbb{N}$  don't have a direct predecessor.

#### Exercise 17

Intuitively we want  $\{a, b, d\}$  and  $\{a, c, d\}$  to be intervals in a structure like:



They are chains all right, but not convex sets: not everything between  $a$  and  $d$  is in  $\{a, b, d\}$ .

#### Exercise 18

$T_1$  and  $T_2$  are chains (because  $T$  is already linear). Suppose  $t < t' < t''$  and  $t, t'' \in T_1$ . Since the cut is a bipartition, either  $t' \in T_1$  or  $t' \in T_2$ . Suppose  $t' \in T_2$ : then  $t'' \in T_1$  and  $t' \in T_2$  but not  $t'' < t'$ , contradicting the definition of cut. So  $t' \in T_1$ , and  $T_1$  is convex. The same argument for  $T_2$ .

#### Exercise 19

Suppose  $\langle T_1, T_2 \rangle$  determines a transition. Then either  $T_1$  has a maximum and  $T_2$  no minimum or the other way around. Suppose  $T_1$  has a maximum  $m$ .  $m$  does have successors (because  $T_2$  is non-empty), but doesn't have a direct successor (because  $T_2$  does not have a minimum). So  $T$  is not discreet to the future. A similar argument holds for the other case.

Suppose  $T$  is not discreet; say, not discreet to the future. That means that there is some element  $t$  that has successors, but no direct successor. Look at  $\langle \{t': t' \leq t\}, \{t': t < t'\} \rangle$ . This is a cut that determines a transition ( $t$  is the maximum of the left part, the right part doesn't have a minimum).

The same argument if  $T$  is not discreet to the past.

#### Exercise 20

The simplest way to go is to simply use the definitions of tree and tree-like structure.



A *diabolo* is a structure  $\langle A, \leq, m \rangle$  where

1.  $\langle \{a \in A: m \leq a\}, \leq, m \rangle$  is a tree
2.  $\langle \{a \in A: a \leq m\}, \geq, m \rangle$  is a tree
3.  $\forall a \in A: a \leq m$  or  $m \leq a$ .

A *diabolo-like structure* is a structure  $\langle A, \leq, m \rangle$  where

1.  $\langle \{a \in A: m \leq a\}, \leq, m \rangle$  is a tree-like structure with origin  $m$
2.  $\langle \{a \in A: a \leq m\}, \geq, m \rangle$  is a tree-like structure with origin  $m$
3.  $\forall a \in A: a \leq m$  or  $m \leq a$

### Exercise 21



- (e)  $\llbracket \text{No woman} \rrbracket = \{\emptyset, \{a\}, \{b\}, \{d\}, \{a, b\}, \{a, d\}, \{b, d\}, \{a, b, d\}\}$

*No woman moves* entails *No woman walks* entails *No woman walks fast*

*No woman moves* entails *No woman skates*

Downward entailing:


if  $X \in \llbracket \text{No woman} \rrbracket$  and  $Y \subseteq X$  then  $Y \in \llbracket \text{No woman} \rrbracket$

See the graph under (d).


Other example: *at most three men*

### Exercise 22


The posets with three elements are:




The poset with eight elements is:




1.




2.




3.



4.



5.



## CHAPTER THREE

## Exercise 1

(a)  $PP\varphi \rightarrow P\varphi$  is true on all transitive frames.

Let  $F$  be transitive and assume that in some point  $t$   $PP\varphi$  is true. That means that for some  $t' < t$   $P\varphi$  is true, and hence for some  $t'' < t'$   $\varphi$  is true. Since  $\leq$  is transitive, it follows that  $t'' < t$ , and hence  $P\varphi$  is true in  $t$ .

(b) If  $F$  is not transitive we can give a counterexample to  $PP\varphi \rightarrow P\varphi$ . Let  $F$  be non-transitive, say  $t'' < t'$  and  $t' < t$  but not  $t'' < t$ . Assign 1

to  $p$  in  $t''$ , and 0 to  $p$  in  $t'$  (and to all  $t'$  such that  $t' < t$ ).  $PPp$  is true in  $t$ , but  $Pp$  is not true in  $t$ . So  $PPp \rightarrow Pp$  is false in  $t$ , and hence  $PP\varphi \rightarrow P\varphi$  is not true on the frame.

## Exercise 2

$F\varphi \rightarrow FF\varphi$  expresses density.

(a)  $F\varphi \rightarrow FF\varphi$  is true on all dense frames.

Let  $F$  be a dense frame. Assume  $F\varphi$  is true in  $t$ . That means that for some  $t' > t$   $\varphi$  is true in  $t'$ . Since  $F$  is dense, there is a  $t''$  between  $t$  and  $t'$ . In  $t''$   $F\varphi$  is true, hence in  $t$   $FF\varphi$  is true.

(b) Let  $F$  be a non-dense frame. Say,  $t < t'$  and nothing is in between  $t$  and  $t'$ . Assign 1 to  $p$  in  $t'$ , and 0 in all  $t''$  later than  $t$  besides  $t'$ . Then  $Fp$  is true in  $t$ , but  $FFp$  is false in  $t$ . Hence  $Fp \rightarrow FFp$  is false in  $t$ , hence  $F\varphi \rightarrow FF\varphi$  is not true on the frame.

## Exercise 3

(a) Let  $T$  be a tautology (for instance  $p \vee \neg p$ ).

$$P\varphi := S(\varphi, T)$$

This says that there is a moment in the past where  $\varphi$  was true and at every later moment a tautology was true, which says just that there is a moment in the past where  $\varphi$  was true.

$$F\varphi := U(\varphi, T)$$

$$H\varphi := \neg P \neg \varphi; \quad G\varphi := \neg F \neg \varphi$$

(b)  $U(\varphi \wedge U(\neg \varphi \wedge \neg U(\varphi, T), \varphi), \neg \varphi)$

At some moment  $t$  in the future  $\varphi$  is true and before that it was never true. At some moment  $t'$  later than  $t$   $\neg \varphi$  is true and between  $t$  and  $t'$   $\varphi$  is true at every moment. After  $t'$   $\varphi$  is never true again. In other words: after having been false uninterruptedly (from the present),  $\varphi$  will be uninterruptedly true for a while and then never be true again after.



## Exercise 4

The semantics given for *before* is:

$$pBq(t_0) \text{ iff } \exists t_1 < t_0 [q(t_1) \wedge \exists t_2 < t_1 [p(t_2)]]$$

$pB(q \vee r)$  does not entail  $pBq$ : if there is a moment in the past where  $p$  is true and a later moment where either  $p$  or  $q$  is true, it doesn't follow that there is a moment in the past where  $p$  is true and a later moment where  $q$  is true.

The idea is:  $p$  before  $q$  means: there is a moment in the past where  $p$  is true and that moment is before every moment where  $q$  is true.

Two alternatives:

$$pBq(t_0) \text{ iff } \exists t_1 < t_0 [p(t_1) \wedge \exists t_2: t_1 < t_2 < t_0 [q(t_2) \\ \wedge \forall t [t < t_0 \wedge q(t) \rightarrow t_1 < t]]]$$

or

$$pBq(t_0) \text{ iff } \exists t_1 < t_0 [p(t_1) \wedge \forall t_2 [t_2 < t_0 \wedge q(t_2) \rightarrow t_1 < t_2]]$$

The first extends the given clause, the second changes it. They share the crucial point that the proposition in the *before* context is analyzed as being in the antecedent clause of a universal statement, which is a downward entailing context. The second definition makes the *before* clause indeed downward entailing, the first does not, for the same reason as the original clause was not downward entailing.

Take *John moved before Mary moved* and *John moved after Mary moved*.

Try to cancel the entailments/implicatures:

- (a) John moved before Mary moved. In fact, John never moved.
- (b) John moved before Mary moved. In fact, Mary never moved.
- (c) John moved after Mary moved. In fact, John never moved.
- (d) John moved after Mary moved. In fact, Mary never moved.

Of these sentences (a), (c) and (d) are clearly contradictions. But (b) doesn't seem to be a contradiction. Consequently, *Mary moved* is an implicature in (b), but an entailment in (d), and *John moved* is an entailment in both.

$$\neg(pAq)(t_0) \text{ iff } \forall t_1 < t_0 [p(t_1) \rightarrow \forall t_2 < t_1 [\neg q(t_2)]]$$

$$\neg(pBq)(t_0) \text{ iff } \forall t_1 < t_0 [q(t_1) \rightarrow \exists t [t < t_0 \wedge p(t) \wedge t_1 \not< t]]$$

A model that makes  $pAq$  true but  $qBp$  false is:

$$\begin{array}{ccccc} t & \text{---} & t' & \text{---} & t'' \\ p & & q & & p \end{array}$$

So, First John hit Mary, then she hit him, then he hit her again. If the judge asks me: Did John hit Mary after she hit him, I will certainly say yes. But if the judge asks me: Did Mary hit John before he hit her, I tend to say no, because John started.

## CHAPTER FOUR

## Exercise 1

The illusion of incompatibility arises if in this dense structure you only check the witness\* condition from left to right, i.e. you look at  $z$  and  $z'$  where  $z < z'$ , and for those you can't satisfy witness\*. But  $z'$  and  $z$  are such that  $z'$  doesn't precede  $z$ , and certainly no subperiod of  $z'$  precedes any subperiod of  $z$ , so there is no incompatibility.

## Exercise 2

1.  $<$  is a strict partial order:

Irreflexivity: assume  $p < p$ . Then for every  $t \in p$ :  $t < t$ .

Transitivity: assume  $p < q$  and  $q < r$ . That means that for every  $t \in p$  and every  $t' \in q$ :  $t < t'$ . Similarly for every  $t' \in q$  and for every  $t'' \in r$ ,  $t' < t''$ . Clearly then for every  $t \in p$  and every  $t'' \in r$ :  $t < t''$ , hence  $p < r$ .

2.  $\sqsubseteq$  is a partial order: this is obvious because  $\subseteq$  is a partial order.

3. Monotonicity: Assume  $p < q$  and  $r \sqsubseteq p$ .  $p < q$  means that every point in  $p$  is before every point in  $q$ .  $r \sqsubseteq p$  means that  $r \subseteq p$ , hence clearly every point in  $r$  is before every point in  $q$ . The other one goes similarly.

4. Convexity. Let  $p < q < r$  and  $p \sqsubseteq s$  and  $r \sqsubseteq s$ . This means that  $p$  and  $r$  are subsets of  $s$  and every point in  $q$  is later than every point in  $p$  but earlier than every point in  $r$ .  $s$  is a period, hence it is convex, so every point in  $q$  is in  $s$ , hence  $q \sqsubseteq s$ .

5. Conjunction: this obviously follows from the second clause of period set.

### Exercise 3


1. Assume  $q \in [p]$  and  $q \sqsubseteq r$ . Then  $p \sqsubseteq r$  and hence  $r \in [p]$ .
2. Assume  $q, r \in [p]$ . Then  $p \sqsubseteq q$  and  $p \sqsubseteq r$ , hence  $p \sqsubseteq q \sqcap r$ , hence  $q \sqcap r \in [p]$ .
3.  $p \in [p]$ , hence  $[p]$  is not empty, hence  $[p]$  is consistent.

### Exercise 4

1. Assume  $p \circ_1 q$ . That means that for some  $e$  in  $p$  and some  $e'$  in  $q$ :  $e \circ e'$ . The latter means that for some  $e''$ :  $e'' \sqsubseteq e$  and  $e'' \sqsubseteq e'$ . Take  $r = [e'']_{\sqsubseteq}$ .  $r \sqsubseteq p$  and  $r \sqsubseteq q$ , hence  $p \circ_2 q$ .
2. Assume  $p \circ_2 q$ . Then for some  $r$ :  $r \sqsubseteq p$  and  $r \sqsubseteq q$ . Hence for some  $e \in r$  and some  $e' \in p$ :  $e \sqsubseteq e'$  and for some  $e'' \in r$  and some  $e''' \in q$ :  $e'' \sqsubseteq e'''$ . Since  $e, e'' \in q$ :  $e \approx e''$ . Hence  $e \sqsubseteq e'''$ , so  $e \sqsubseteq e'$  and  $e' \in p$  and  $e \sqsubseteq e'''$  and  $e''' \in q$ , hence  $e' \circ e'''$ , hence  $p \circ_1 q$ .

### Exercise 5

(a)




$$a = \{1, 7\}; b = \{2, 4, 7\}; c = \{3, 4, 8\}; d = \{3, 5, 9\}; e = \{6, 9\}$$

$$a < b < c < d < e$$

$$\begin{array}{ccccccc} & & \{b\} & & \{c, d\} & & \\ \{a\} & & \{b, c\} & & \{d\} & & \{e\} \\ & \{a, b\} & & \{c\} & & \{d, e\} & \end{array}$$

(b)



$$a = \{1\}; b = \{2, 4\}; c = \{3, 4\}; d = \{3, 5\}; e = \{6\}$$

$$a < b < c < d < e$$

Let's try to reconstruct the periods. What we would want is:

$$\begin{array}{ccc} & \{b\} & \{c, d\} \\ \{a\} & & \{e\} \\ & \{b, c\} & \{d\} \end{array}$$

But this is not a period set, because  $\{b, c\} \cap \{c, d\} = \{c\}$ , but there is no period  $\{c\}$ .

## CHAPTER FIVE

### Exercise 1

$q$  overlaps the beginning of  $p$ :  $q \circ p \wedge \neg \exists r[r \sqsubseteq p \wedge r < q]$ . Similarly  $q'$  overlaps the end of  $p$ .

### Exercise 2

Assume that the period structure is dense and assume that  $B\varphi$  holds at  $p$ . That means that  $p$  can be split in  $p'$  where  $\varphi$  is false and  $p''$  where  $\varphi$  is true, with  $p' < p''$ . Since the period structure is dense, we can split  $p'$  into  $q$  and  $q'$ , with  $q < q'$  and  $p''$  into  $r$  and  $r'$  with  $r < r'$ . Consequently,  $\varphi$  is false at  $q'$  and true at  $r$ . But then  $q' \sqcup r$  satisfies condition (1), hence  $B\varphi$  does not hold at  $p$ .

$B\varphi$  can only hold at periods  $p$  that are the union of two atomic periods  $t$  and  $t'$  where at  $t$   $\varphi$  is false and at  $t'$   $\varphi$  is true. Clearly in that case condition 1 is satisfied and there is no subperiod for which that holds (clearly  $B\varphi$  cannot be true in an atomic period).

Suppose that  $p$  is not the union of two atoms but satisfies Condition 1. Partition  $p$  in the part where  $\varphi$  is false and the part where  $\varphi$  is true.

There is either a jump, a transition or a gap between those periods (where a jump means that there is an atom overlapping the end of the part where  $\varphi$  is false and an atom overlapping the beginning of the part where  $\varphi$  is true). In all these cases you can find a subperiod of  $p$  satisfying Condition 1.

## Exercise 3



$\varphi$	$\neg\varphi$	$A\varphi$	$\neg A\varphi$	$\varphi \wedge \neg\varphi$	$\varphi \vee \neg\varphi$	$\varphi \wedge \neg A\varphi$	$\varphi \vee \neg A\varphi$	$A\varphi \wedge \neg A\varphi$	$A\varphi \vee \neg A\varphi$
1	0	1	0	0	1	0	1	0	1
0	1	0	1	0	1	0	1	0	1
*	*	0	1	*	*	*	1	0	1

## CHAPTER SIX

## Exercise 1

(a) The first one is a lattice, the second isn't: there is not a unique smallest element that is the join of the two elements directly above the bottom.

(b)



It is easy to check that all the joins and meets are the same.

We have to show that for all  $X \subseteq A$ :  $\bigvee X \in A$ . Let  $X \neq \emptyset$ . Look at  $\{y \in A: \forall x \in X[x \leq y]\}$ . This is a subset of  $A$ , hence  $\bigwedge\{y \in A: \forall x \in X[x \leq y]\} \in A$ . This is, of course,  $\bigvee X$ . Let  $X = \emptyset$ .

Since there is no  $y \in \emptyset$ , trivially  $\forall y \in A \forall x \in \emptyset[x \leq y]$ . Hence  $\{y \in A: \forall x \in \emptyset[x \leq y]\} = A$ . Hence  $\bigwedge\{y \in A: \forall x \in \emptyset[x \leq y]\} = \bigwedge A$ . So  $\bigvee \emptyset = \bigwedge A$ .

(c) This follows directly from the previous considerations: since  $A$  is complete  $\bigvee A$  and  $\bigwedge A$  are in  $A$ .  $\bigvee A$  is the maximum of  $A$ ,  $\bigwedge A$  is the minimum of  $A$ .

(d) Suppose  $A$  has two distinct maximal elements  $a$  and  $b$ . Since  $A$  is a lattice  $a \vee b$  exists and  $a \leq a \vee b$  and  $b \leq a \vee b$ . But then  $a$  and  $b$  wouldn't be maximal.

(e) If  $A$  is a finite lattice then every subset  $X$  of  $A$  is finite. Then it follows by the theorem given that for every non-empty subset  $X$  of  $A$   $\bigvee X, \bigwedge X \in A$ . This means that  $A$  has a minimum  $\bigwedge A$  and a maximum  $\bigvee A$ . By the argumentation under (a)  $\bigvee \emptyset = \bigwedge A$ , similarly  $\bigwedge \emptyset = \bigvee A$ , hence  $\bigvee \emptyset, \bigwedge \emptyset \in A$ , so  $A$  is complete.

## Exercise 2

Let  $X \subseteq \text{pow } B$ . Then obviously  $\bigcup X, \bigcap X \subseteq B$ , hence  $\bigcup X, \bigcap X \in \text{pow } B$ .

## Exercise 3

The singleton sets.

## Exercise 4

Define in terms of the intersection operation  $\cap$  in the interval lattice the new operation  $\wedge$ :

Let  $i$  and  $i'$  be non-singleton intervals. Define:

$$\begin{aligned} i \wedge i' &:= i \cap i' \text{ if } i \cap i' \text{ is not a singleton} \\ &:= \emptyset \text{ if } i \cap i' \text{ is a singleton} \end{aligned}$$

Then eliminate all singletons from the structure. The structure is a complete atomless lattice. To get a structure that is neither atomless, nor atomic, leave, say, one atom  $\{t\}$  and modify the above definition replacing 'not a singleton' by 'not a singleton except  $\{t\}$ ' twice.

## Exercise 5

A.1.  $\langle A, * \rangle$  is a semilattice, so  $*$  is idempotent, commutative and associative. Let  $\leq_\wedge$  be as given.

- a.  $\leq_\wedge$  is a partial order.
- $\leq_\wedge$  is reflexive:  $a \leq_\wedge a$ . This means  $a * a = a$ , which is idempotency.
- $\leq_\wedge$  is transitive. Assume  $a \leq_\wedge b$  and  $b \leq_\wedge c$ .  $a * b = a$  and  $b * c = b$ . Then  $a * c = a * b * c$  (using the first)  $= a * b$  (using the second)  $= a$  (using the first).
- $\leq_\wedge$  is antisymmetric. Assume  $a \leq_\wedge b$  and  $b \leq_\wedge a$ . This means  $a * b = a$  and  $b * a = b$ , then, by commutativity  $a = b$ .
- b.  $*$  is meet in  $\leq_\wedge$ .
- $a * b \leq_\wedge a$ . This means  $(a * b) * a = a * b$ .  $(a * b) * a = (b * a) * a = b * (a * a) = b * a$ .
- Similarly,  $a * b \leq_\wedge b$ .
- Assume  $c \leq_\wedge a$  and  $c \leq_\wedge b$ . Then  $c * a = c$  and  $c * b = c$ . Then  $c * (a * b) = (c * a) * b = c * b = c$ . Hence  $c \leq_\wedge a * b$ , so indeed  $a * b$  is the meet of  $a$  and  $b$ . So we have proved that  $\langle A, \leq_\wedge \rangle$  is a meet semilattice.

2. The argument that  $\langle A, \leq_\vee \rangle$  is a join semilattice goes in exactly the same way:

- a.  $\leq_\vee$  is a partial order.
- $\leq_\vee$  is reflexive:  $a \leq_\vee a$ . This means  $a * a = a$ , which is idempotency.
- $\leq_\vee$  is transitive. Assume  $a \leq_\vee b$  and  $b \leq_\vee c$ .  $a * b = b$  and  $b * c = c$ . Then  $a * c = a * b * c$  (using the second)  $= b * c$  (using the first)  $= c$  (using the second).
- $\leq_\vee$  is antisymmetric. Assume  $a \leq_\vee b$  and  $b \leq_\vee a$ . This means  $a * b = b$  and  $b * a = a$ , then, by commutativity  $a = b$ .
- b.  $*$  is join in  $\leq_\vee$ .
- $a \leq_\vee a * b$ . This means  $a * (a * b) = a * b$ .  $a * (a * b) = (a * a) * b = a * b$ .
- Similarly,  $b \leq_\vee a * b$ .
- Assume  $a \leq_\vee c$  and  $b \leq_\vee c$ . Then  $a * c = c$  and  $b * c = c$ . Then  $(a * b) * c = a * (b * c) = a * c = c$ , hence  $a * b \leq_\vee c$ , so indeed  $a * b$  is the join of  $a$  and  $b$ . So we have proved that  $\langle A, \leq_\vee \rangle$  is a join semilattice.

B. Let  $\langle A, \leq \rangle$  be a join semilattice and  $\langle A, \vee \rangle$  be the semilattice we get when we take join as an operation. We define in  $\langle A, \vee \rangle$ :  $a \leq' b$  iff  $a \vee b = b$ .

Claim:  $a \leq' b$  iff  $a \leq b$ . This means that we have to prove:  $a \leq b$  iff  $a \vee b = b$ . This just follows from the fact that  $\vee$  is join in  $\langle A, \leq \rangle$ .

Let  $\langle A, * \rangle$  be a semilattice. Define  $a \leq_\vee b := a * b = b$ . Form  $\langle A, \leq_\vee \rangle$ . This is, as proved, a join semilattice. We form  $\langle A, \vee \rangle$ .

Claim:  $a \vee b = a * b$ . This follows obviously from the fact that we defined  $\leq_\vee$  that way.


C. The only thing that does hold is that we can regard a join semilattice as a meet semilattice (by turning it upside down).  $\langle A, \leq_\wedge \rangle$  is a meet semilattice and  $\langle A, \leq_\vee \rangle$  a join semilattice, but they are not the same structure, in particular  $\leq_\wedge \neq \leq_\vee$ . In fact,  $\leq_\wedge = \geq_\vee$ .

## Exercise 6

The definition of atomic says that every non-zero element has an atom below it. Atoms, however are defined as elements that have only 0 below them. The definition of atom, hence, presupposes that the structure has a zero, hence  $\langle A, \leq \rangle$  has a zero. Then  $\forall \emptyset \in A$  and hence  $A$  is complete.


## Exercise 7

(a) No. The diamond is a substructure of this structure:



It can be checked that this is a substructure by checking the joins and meets of  $a, b$  and  $c$  (1 and 0, resp.).


(b) Yes. It may look at first sight as if the structure under (a) is a substructure of this in mirror image, but if you check the distributive laws on  $a, b, c$  in the following structure, you will see that they hold here:





## Exercise 11

(a)



First: obviously none of  $\{a\}$ ,  $\{b\}$ ,  $\{c\}$  generate the whole lattice.  $\{a, b\}$  only generates itself.


$\{b, c\}$  generates  $\{b, c, 0, 2\}$

$\{a, c\}$  generates  $\{a, c, 1, 5\}$

So none of these generate the whole structure.


By the above considerations  $\{a, b, c\}$  generates at least  $\{a, b, c, 0, 1, 2, 5\}$ . 3 is generated as the meet of  $a$  and 2; 4 is generated as the join of  $b$  and 5, hence indeed the whole structure is generated and as argued, minimally.

(b)



1.  $\{\{a\}, \{b\}, \{c\}\}$
2.  $\{\{a, b\}, \{a, c\}, \{b, c\}\}$
3.  $\{\{a\}, \{b\}, \{a, c\}, \{b, c\}\}$
4.  $\{\{a\}, \{c\}, \{a, b\}, \{b, c\}\}$
5.  $\{\{b\}, \{c\}, \{a, b\}, \{a, c\}\}$

(c) Example



For the new sets of minimal generations you have to add  $x$  and  $y$  to the old sets.

(d) It has to be generated under the operations. In a Boolean lattice, the operations are  $\wedge$  and  $\vee$ . You need minimally four elements: the four atoms (the case is similar to (b) above).

As a Boolean algebra the operations are  $\wedge$ ,  $\vee$  and  $\neg$ . To generate it as a Boolean algebra you only need two elements. In the example of Exercise 9c,  $e$  and  $f$  generate the Boolean algebra. They bring in  $\neg e$  and  $\neg f$ . By taking meets this brings in  $0$ ,  $a$ ,  $b$  and  $c$ . Taking complements brings in  $1$ ,  $\neg a$ ,  $\neg b$ ,  $\neg c$ . The join of  $e$  and  $f$  brings in  $\neg d$ , hence complements brings in  $d$ . The join of  $a$  and  $d$  brings in  $g$ , hence complements  $\neg g$ .

## Exercise 12

$\approx$  is clearly an equivalence relation (because  $=$  is). So we have to show: if  $a \approx a'$  and  $b \approx b'$  then:

$$(a \wedge b) \approx (a' \wedge b')$$

$$(a \vee b) \approx (a' \vee b')$$

So assume  $a \approx a'$  and  $b \approx b'$ . That means that  $h(a) = h(a')$  and  $h(b) = h(b')$ . Since  $h$  is a homomorphism  $h(a \wedge b) = h(a) \wedge h(b) = h(a') \wedge h(b') = h(a' \wedge b')$ . Hence  $a \wedge b \approx a' \wedge b'$ . Similarly for  $\vee$ .

## Exercise 13

$\approx$  is the relation  $\lambda a \lambda a'. h(a) = h(b)$ .

1.  $f$  clearly is a surjection.
2.  $f$  is an injection.

Assume  $f([a]_{\approx}) = f([a']_{\approx})$  that means that  $h(a) = h(a')$ , but then  $a \approx a'$  and  $[a]_{\approx} = [a']_{\approx}$ .


3.  $f$  is a homomorphism.

Since  $\approx$  is a congruence relation:  $[a \wedge a']_{\approx} = [a]_{\approx} \wedge [a']_{\approx}$ . Hence

$$\begin{aligned} f([a]_{\approx} \wedge [a']_{\approx}) &= f([a \wedge a']_{\approx}) = h(a \wedge a') = h(a) \wedge h(a') \\ &= f([a]_{\approx}) \wedge f([a']_{\approx}) \end{aligned}$$

Similarly for  $\vee$ .

#### Exercise 14



#### Exercise 15

(a) Let  $F$  be a filter in  $L$ . If  $a, b \in F$  then  $a \wedge b \in F$ , by definition. If  $a, b \in F$  then, since  $a \leq a \vee b$ ,  $a \vee b \in F$ . Hence  $F$  is a sublattice of  $L$ .

Let  $a, b \in F$  and assume  $a \leq x \leq b$ , then obviously  $x \in F$ , since  $F$  is a filter, so  $F$  is convex.

(b) Assume that  $F$  is a filter. Then:

if  $a, b \in F$  then  $a \wedge b \in F$  (filter)  
if  $a \wedge b \in F$  then  $a, b \in F$  because  $a \wedge b \leq a, b$

– Assume that  $F$  is a non-empty subset of  $A$  such that  $a, b \in F$  iff  $a \wedge b \in F$ . Then obviously if  $a, b \in F$   $a \wedge b \in F$ .

Suppose  $a \in F$  and  $a \leq b$ . Then  $a = a \wedge b$ , hence  $a \wedge b \in F$ , hence  $b \in F$ . So  $F$  is a filter.

This proves the equivalence of the first definition.

– If  $F$  is a filter we have proved that it is a sublattice of  $L$ . Let  $a \in F$  and  $b \in L$ .  $a \leq a \vee b$ , hence, because  $F$  is a filter  $a \vee b \in F$ .

– Let  $F$  be a sublattice of  $L$  such that if  $a \in F$  and  $b \in L$  then  $a \vee b \in F$ . If  $a, b \in F$ , then because  $F$  is a sublattice and hence closed under  $\wedge$ ,

$a \wedge b \in F$ . Let  $a \in F$  and  $a \leq b$ . Then  $b = a \vee b$ ,  $a \vee b \in F$ , hence  $b \in F$ . So  $F$  is a filter.

This proves the equivalence of the second definition.

Non-empty subset  $I$  of  $L$  is an ideal in  $L$  iff

$$a \vee b \in I \text{ iff } a, b \in I.$$


$I$  is an ideal in  $L$  iff  $I$  is a sublattice of  $L$  such that:

$$\text{if } a \in I \text{ and } b \leq a \text{ then } a \wedge b \in I.$$

#### Exercise 16


If  $L$  is a finite lattice every subset of  $L$ , hence every filter in  $L$  is finite. Thus  $F = \{a_1, \dots, a_n\}$ . Since  $F$  is closed under conjunction it has a minimal element  $a_1 \wedge \dots \wedge a_n$  and it is the principle filter generated by that minimal element.

#### Exercise 17




Prime filters: none

Ultrafilters:  $b, c, d$



Prime filters:  $b, c, e, f$

Ultrafilters:  $e, f$



$\{1, a, b, d\}$ ,  $\{1, a, e, c\}$ ,  $\{1, b, c, f\}$  are the prime filters and the ultrafilters.

### Exercise 18

Let  $F$  be a prime filter. Hence if  $a \vee b \in F$  then either  $a \in F$  or  $b \in F$ .

Suppose  $a, b \in L - F$ . Then, because  $F$  is prime,  $a \vee b \in L - F$ . Suppose  $a \in L - F$  and  $b \leq a$ . If  $b \in F$ , then  $a \in F$ , because  $F$  is a filter, hence  $b \in L - F$ . So  $L - F$  is an ideal, and since  $F$  is a proper filter  $L - F \neq L$ , so  $L - F$  is a proper ideal.

Assume that  $a \wedge b \in L - F$  and assume that both  $a \in F$  and  $b \in F$ . Then, because  $F$  is a filter  $a \wedge b \in F$ , contradiction. Hence either  $a \in L - F$  or  $b \in L - F$ , hence  $L - F$  is a prime ideal. The other side goes in the same way.

### Exercise 19

Let  $F$  be a proper filter and assume that for every  $a$ : either  $a \in F$  or  $\neg a \in F$ . Assume that  $b \notin F$ , then  $\neg b \in F$ .

Since  $b \wedge \neg b = 0$  this means that  $b$  is incompatible with  $F$ , hence  $F$  is an ultrafilter.

### Exercise 20

Let  $F$  be a prime filter and assume that  $a \notin F$ .  $a \vee \neg a \in F$ , hence, since  $F$  is prime,  $\neg a \in F$ . Hence  $F$  is an ultrafilter.

### Exercise 21

There is no countable powerset Boolean algebra. If  $B$  is finite then  $\text{pow } B$  is finite, if  $B$  is countable, then  $\text{pow } B$  is uncountable.

## REFERENCES

- Anscombe, E., 1964, 'Before and after', in: *Philosophical Review* 73.
- Bach, E., 1979, 'Tenses and aspects as functions of verb-phrases', in: Rohrer (Ed.), *Time, Tense and Quantifiers*, Niemeyer, Tübingen.
- Ballmer, T., 1982, 'The answer is: as faltz as keen(an)', in: *Theoretical Linguistics* 9.
- Bealer, G., 1982, *Quality and Concept*, Clarendon Press, Oxford.
- Bealer, G. and U. Mönnich, 1989, 'Property theories', in: Gabbay and Guentner (Eds.), *Handbook of Philosophical Logic IV*, Kluwer, Dordrecht.
- Bennett, M., 1975, *Some Extensions of a Montague Fragment of English*, Diss. UCLA, IULC, Bloomington.
- van Benthem, J., 1977, 'Tense logic and standard logic', in: *Logique et Analyse* 80.
- van Benthem, J., 1983, *The Logic of Time*, Kluwer, Dordrecht.
- van Benthem, J., 1984, 'Correspondence theory', in: Gabbay and Guentner (Eds.), *Handbook of Philosophical Logic II*, Kluwer, Dordrecht.
- Bunt, H., 1985, *Mass Terms and Model Theoretic Semantics*, Cambridge UP, Cambridge (U.K.).
- Burgess, J., 1984, 'Basic tense logic', in: Gabbay and Guentner (Eds.), *Handbook of Philosophical Logic II*, Kluwer, Dordrecht.
- Chierchia, G., 1984, *Topics in the Syntax and Semantics of Infinitives and Gerunds*, Diss. UMass, Amherst, GLSA, Amherst; also Garland, New York (1989).
- Chierchia, G. and R. Turner, 1988, 'Semantics and property theory', in: *Linguistics and Philosophy* 11.
- Chierchia, G. and S. McConnell-Ginet, 1990, *Meaning and Grammar*, Cambridge UP, Cambridge (MA).
- Cooper, R., 1983, *Quantification and Syntactic Theory*, Kluwer, Dordrecht.
- Cresswell, M., 1977, 'Interval semantics and logical words', in: Rohrer (Ed.), *On the Logical Analysis of Tense and Aspect*, TBL Verlag, Tübingen. Reprinted in Cresswell (1985).
- Cresswell, M., 1985, *Adverbial Modification*, Kluwer, Dordrecht.
- van Dalen, D., 1986, 'Intuitionistic logic', in: Gabbay and Guentner (Eds.), *Handbook of Philosophical Logic III*, Kluwer, Dordrecht.
- Dowty, D., 1979, *Word Meaning and Montague Grammar*, Kluwer, Dordrecht.
- Dowty, D., Wall, R., and S. Peters, 1981, *Introduction to Montague Semantics*, Kluwer, Dordrecht.
- Dummett, M., 1973, 'The philosophical basis of intuitionistic logic', in: Dummett, *Truth and other Enigmas*, Harvard UP, Cambridge (MA).
- Dummett, M., 1977, *Elements of Intuitionism*, Oxford UP, Oxford.
- Enç, M., 1981, *Tense without Scope*, Diss. U. of Wisconsin, Madison.
- Fine, K., 1975, 'Truth, vagueness and logic', in: *Synthese* 30.
- Gallin, D., 1975, *Intensional and Higher-order Modal Logic*, North Holland, Amsterdam.
- Gamut, 1990, *Logic, Language and Meaning* (two volumes), U. of Chicago Press, Chicago.



- Grätzer, G., 1978a, *Universal Algebra*, 2nd edition, Springer, New York.
- Grätzer, G., 1978b, *General Lattice Theory*, Birkhäuser, Basel.
- Groenendijk, J. and M. Stokhof, 1982, 'Semantic analysis of wh-complements', in: *Linguistics and Philosophy* 5.
- Groenendijk, J. and M. Stokhof, 1985, *Studies in the Semantics of Questions and the Pragmatics of Answers*, Diss. U. of Amsterdam.
- Heinäsmäki, O., 1977, *Semantics of English Temporal Connectives*, Diss. U. of Texas, Austin, IULC, Bloomington.
- Hendriks, H., 1987, 'Type change in semantics: the scope of quantification and coordination', in: Klein and van Benthem (Eds.), *Categories, Polymorphisms and Unification*, Edinburgh/Amsterdam.
- Hinrichs, E., 1981, 'Temporale Anaphore im Englischen', Magisterarbeit, U. of Tübingen.
- Hinrichs, E., 1985, *A Compositional Semantics for Aktionsarten and NP reference in English*, Diss. Ohio State U., Columbus.
- Hodges, W., 1983, 'Elementary predicate logic', in: Gabbay and Guenther (Eds.), *Handbook of Philosophical Logic I*, Kluwer, Dordrecht.
- Humberstone, L., 1979, 'Interval semantics for tense logic', in: *Journal of Philosophical Logic* 8.
- Janssen, T., 1983, *Foundations and Applications of Montague Grammar*, Diss. U. of Amsterdam, Mathematical Centre, Amsterdam.
- Kadmon, N. and F. Landman, 1990, 'Polarity sensitive any and free choice any', in: Stokhof and Toorenvliet (Eds.), *Proceedings of the Seventh Amsterdam Colloquium*, ITLI, Amsterdam.
- Kamp, H., 1968, *Tense Logic and the Theory of Linear Order*, Diss. UCLA.
- Kamp, H., 1970, 'Formal properties of now', in: *Theoria* 37.
- Kamp, H., 1975, 'Two theories about adjectives', in: Keenan (Ed.), *Semantics for Natural Language*, Cambridge UP, Cambridge (UK).
- Kamp, H., 1979a, 'Events, instants and temporal reference', in: Bäuerle, Egli and von Stechow (Eds.), *Semantics from Different Points of View*, Springer, Berlin.
- Kamp, H., 1979b, 'Some remarks on the logic of change. Part I', in: Rohrer (Ed.), *Time, Tense and Quantifiers*, Niemeyer, Tübingen.
- Kaplan, D., 1978, 'Dthat', in: Cole (Ed.), *Syntax and Semantics 9: Pragmatics*, Academic Press, New York.
- Keenan, E. and L. Faltz, 1985, *Boolean Semantics for Natural Language*, Kluwer, Dordrecht.
- Krifka, M., 1990, 'Four thousand ships passed through the lock', in: *Linguistics and Philosophy* 13.
- Kripke, S., 1965, 'Semantical analysis of intuitionistic logic I', in: Crossley and Dummett (Eds.), *Formal Systems and Recursive Functions*, North Holland, Amsterdam.
- Ladusaw, W., 1980, *Polarity Sensitivity as Inherent Scope Relations*, Diss. U. of Texas, Garland, New York.
- Landman, F., 1986, *Towards a Theory of Information*, Diss. U. of Amsterdam, GRASS 6, Foris, Dordrecht.
- Landman, F., 1989, 'Groups I' and 'groups II', in: *Linguistics and Philosophy* 12.
- Landman, F., 1990, 'Partial information, modality and intentionality', in: Hanson (Ed.), *Information, Language and Cognition*, U. of British Columbia Press, Vancouver.

- Lewis, D., 1973, *Counterfactuals*, Harvard UP, Cambridge (MA).
- Lewis, D., 1979, 'Scorekeeping in a language game', in: Bäuerle, Egli and von Stechow (Eds.), *Semantics from Different Points of View*, Springer, Berlin.
- Linebarger, M., 1987, 'Negative polarity and grammatical representation', in: *Linguistics and Philosophy* 10.
- Link, G., 1983, 'The logical analysis of plurals and mass terms: a lattice-theoretical approach', in: Bäuerle, Schwarze and von Stechow (Eds.), *Meaning, Use and the Interpretation of Language*, deGruyter, Berlin.
- Link, G., 1984, 'Hydras. On the logic of relative clause constructions with multiple heads', in: Landman and Veltman (Eds.), *Varieties of Formal Semantics*, GRASS 3, Foris, Dordrecht.
- Link, G., forthcoming, 'Plural', in: Wunderlich and von Stechow (Eds.), *Handbook of Semantics*.
- Link, G., 1987, 'Algebraic semantics of event structures', in: Groenendijk, Stokhof and Veltman (Eds.), *Proceedings of the Sixth Amsterdam Colloquium*, ITLI, Amsterdam.
- Montague, R., 1970, 'Universal Grammar', in: *Theoria* 36, reprinted in: Thomason (Ed.), 1974, *Formal Philosophy*, Yale UP, New Haven.
- Montague, R., 1973, 'The proper treatment of quantification in ordinary English', in: Hintikka, Moravcsik and Suppes (Eds.), *Approaches to Natural Language*, Kluwer, Dordrecht. Reprinted in: Thomason (Ed.), 1974, *Formal Philosophy*, Yale UP, New Haven.
- Oversteegen, E., 1989, *Tracking Time*, Diss. U. of Utrecht.
- Parsons, T., 1989, 'The progressive in English: events, states and processes', in: *Linguistics and Philosophy* 12.
- Partee, B., 1973, 'Some structural analogies between tenses and pronouns in English', in: *Journal of Philosophy* 70.
- Partee, B., 1984, 'Nominal and temporal anaphora', in: *Linguistics and Philosophy* 7.
- Partee, B. and M. Rooth, 1983, 'Generalized conjunction and type ambiguity', in: Bäuerle, Schwarze and von Stechow (Eds.), *Meaning, Use and the Interpretation of Language*, de Gruyter, Berlin.
- Partee, B., A. ter Meulen, and R. Wall, 1990, *Mathematical Methods in Linguistics*, Kluwer, Dordrecht.
- Pelletier, J. and L. Schubert, 1989, 'Mass expressions', in: Gabbay and Guenther (Eds.), *Handbook of Philosophical Logic IV*, Kluwer, Dordrecht.
- Pinkal, M., 1984, 'Consistency and context change: the sorites paradox', in: Landman and Veltman (Eds.), *Varieties of Formal Semantics*, GRASS 3, Foris, Dordrecht.
- Prior, A., 1967, *Past, Present and Future*, Oxford UP, Oxford.
- Reichenbach, H., 1947, *Elements of Symbolic Logic*, U. of California Press, Berkeley.
- Scha, R., 1981, 'Distributive, collective and cumulative quantification', in: Groenendijk, Janssen and Stokhof (Eds.), *Formal Methods in the Study of Language*, Mathematical Centre Tracts, Amsterdam. Reprinted in: Groenendijk, Janssen and Stokhof (Eds.), 1983, *Truth, Interpretation, Information*, GRASS 2, Foris, Dordrecht.
- Scott, D., 1979, 'Identity and existence in intuitionistic logic', in: Fourman, Mulvey and Scott (Eds.), *Applications of Sheaves. Proceedings Durham 1977*, Springer, Berlin.
- Stalnaker, R., 1968, 'A theory of conditionals', in: Rescher (Ed.), *Studies in Logical Theory*, Blackwell, London.

- Stalnaker, R., 1978, 'Assertion', in Cole (Ed.), *Syntax and Semantics 9: Pragmatics*, Academic Press, New York.
- Taylor, B., 1977, 'Tense and continuity', in: *Linguistics and Philosophy* 1.
- Thomason, R., 1984, 'Combinations of tense and modality', in: Gabbay and Guenther (Eds.), *Handbook of Philosophical Logic II*, Kluwer, Dordrecht.
- Tichy, P., 1985, 'Do we need interval semantics', in: *Linguistics and Philosophy* 8.
- Van Fraassen, B., 1971, *Formal Semantics and Logic*, Macmillan, New York.
- Veltman, F., 1985, *Logics for Conditionals*, Diss. U. of Amsterdam.
- Vlach, F., 1981, 'The semantics of the progressive', in: Tedeschi and Zaenen (Eds.), *Syntax and Semantics 14: Tense and Aspect*, Academic Press, New York.
- Westerstål, D., 1988, 'Quantifiers in Formal and Natural Languages', in: Gabbay and Guenther (Eds.), *Handbook of Philosophical Logic IV*, Kluwer, Dordrecht.
- Whorf, B., 1956, 'Science and Linguistics', in: Carroll (Ed.), *Language, Thought and Reality*, Wiley, New York.

## INDEX

- Absorption 236–283
- Algebra, Boolean  $x$ , 95, 117, 128, 177, 247–283, 351
- complete atomic 262–267, 284–324
- free complete atomic 287–324
- powerset 267–283, 354
- representation theorem for 280
- representation theorem for complete atomic 280–283
- Kleene 162
- Lindenbaum 95, 283
- relational 72–120
- universal 70–120, 248–267
- Algebraic semantics, functional
- completeness and expressibility 127–135
- Anscombe, E. 140
- Antisymmetric relation 83–105
- Antisymmetry 238, 327, 332, 346
- Approximation, first order 30
- set of 181–186
- Arbitrary consistent theory 18
- Argument, transfer 33
- Arithmetics, Peano 29, 36–69
- Assignment function 4, 5, 26, 28, 128–135, 304, 320
- Associativity 236–283, 346, 348
- Asymmetric relation 83–105
- Atomic lattice 242–283
- Atomicity condition 175
- Atomless lattice 240–283
- Automorphism 77–120, 329, 330
- Axiom, empty set 48
- foundation 53–69
- identity 8
- infinity 58
- of choice 55–69, 115
- of pair 49, 50
- of separation 46
- of sum 49, 50
- propositional 8
- quantifier 8
- schemata 7, 40, 46
- Axiomatic theory 7
- Bach, E. 121
- Background context 160
- Ballmer, T. 294
- Bar 109–120
- Base step 9, 15, 63
- Bealer, G. 32
- Bennett, M. 300
- Benthem, J. van ix, 121, 127, 135, 171, 176, 177, 180, 186, 200
- Beth comb 156
- frame 154–156
- model 150–152, 154–156
- semantics 154–156
- Bijection 51–69, 185, 258–267, 290, 292
- notion of 70–120
- Bijjective homomorphism 76–120, 329, 330
- Binary relation 70–120
- Bivalence principle 211, 222–233
- Blockset, maximal 315–324
- Boolean algebra  $x$ , 95, 117, 128, 177, 247–283, 351
- join homomorphism 301–324
- lattice 247–283
- semantics 284–299
- Bound, lower 234–283
- Bounded distributive lattice 246–283
- lattice 240
- Branch 109–120
- Brouwer intuitionism 146
- Burgess, J. 121
- Cantor mathematical construction 36
- set theory 45

- theorem 65
- Cardinal number 54–69, 258, 259
- Cardinality, finite 21
- infinite 21
- notion of 54–69
- Cartesian product 49, 50, 51, 72
- Chain ix, 109–120
- maximal ix
- Change, intervals events and 197–233
  - Kamp's logic of 221–233
  - moment of 211–218
- Characteristic function 70–120
- Chierchia, G. viii, x, 32, 44
- Choice, axiom of 55–69, 115
- function 55–69
- Closure, deductive 10, 36
- downward 202
- Co-final 103–120
- Co-initial 103–120
- Cohen's forcing technique 60
- Comb, Beth 156
- Common predecessor 334, 335
- successor 334, 335
- Commutativity 236–283
- Compactness theorem 13, 19, 29, 31
- Complemented lattice 247–283
- Completeness 158
- Complete atomic Boolean algebra
  - 262–267, 284–324
  - lattice 269–283
  - bounded 265–267
- Completed change principle 233
- Completeness 326, 327
- functional 128–135
- notion of 256–267
- theorem 13, 18, 19, 24, 29, 31, 36
- Comprehension axiom schema 26
- Comprehension, principle of 45
- Condition, atomicity 175
- conjunction 174
- direction 174
- disjunction 174
- Congruence relation 95–105, 251–254, 352
- Congruence structure 95–105
- Conjunction 188–196, 288–299, 342
- condition 174
- Connected relation 83–105
- Constant function 37, 70–120
- individual 24, 37, 159
- and predicate 33–44
- logical 3, 7, 33–44
- mathematical 33–44, 45
- non-logical 4, 33–44
- non-mathematical 33–44
- non-primitive logical 7
- predicate 37
- Constructions with partial orders
  - 171–196
- Context, background 160
- set 99
- Continuum hypothesis 56–69
- Convex set 110–120
- sublattice 267–283
- Convexity 188–196, 341
- principle 176
- Cooper, R. 30
- Correctness theorem 29
- Countable model 60–69
- Cressell, M. 200, 208
- Dalen, D. van 145, 152
- Deductive closure 10, 36
- Definability, first order 1
- Definition, entailment 7
- truth 5, 26
- Dense linear order 64
- theory of 35–44
- without endpoints 64–69
- Density 339
- principle 176
- Diabolo 335, 336
- Diagram, Hasse 86
- Diamond 245, 265, 347, 348
- Direct interpretation 75
- Direction condition 174
- Discreet change principle 233
- structure 138
- Discreetness 108–120
- Disjunction 288–299
- condition 174
- Distance structure 104–120
- Distributive lattice 244–283
- set theoretic representation of x
- Distributivity 314–324, 348
- Domain function 153

- Downward closure 202
- entailing 118, 142–170, 337, 340
- Dowty, D. viii, 76, 200, 201, 204, 206, 210, 211, 213
- Duality 87–105, 235–283
- Dummett, M. 145
- Element, maximal 158, 181, 236, 305, 345
- Element, minimal 105–120, 139
- Embedding 76–120, 329, 330
- Empty set 51
- axiom 48
- Enç, M. 121
- Endpoints, dense linear order without
  - 64–69
  - theory of 36–44
- Entailing, downward 118, 142–170, 337, 340
- definition of 7
- Epimorphism 76–120, 329
- strong 82–120
- Equivalence relation 84–105
- partial order and 83–105
- partition and ix
- Event model 226
- structure x, 186–196
- Expressibility, algebraic semantics
  - functional completeness and 127–135
- Expression, semantics of x
- Extended 103–120
- Extension 97–105, 292
- negative 158–170
- positive 158–170
- Extensionality 327
- Faltz, L. 258, 284–324
- Filter 352, 353
- ideals and 267–283
- maximally proper 271–283
- notion of 180–186
- prime 271–283, 353, 354
- proper 271–283, 354
- representation ix
- Fine, K. 158, 162–165
- Finitary relation 72–120
- Finite cardinality 21
- intersection property 271–283
- lattice 353
- Finiteness 327
- First order approximation 30
- definability 1
- language 2, 4, 10, 14, 17, 24
- lexicon 2, 3
- logic 1, 2–13, 20, 21, 24, 26
- notion of 1
- predicate logic 2, 7
- quantification 31
- sentence 10, 12, 22, 33
- theory 1, 12, 19, 20, 32–44, 328
- Forcing technique, Cohen's 60
- Formalism, Hilbert's 43
- Formalization in semantics vii
- Formula induction 9
- Foundation axiom 53–69
- Fraassen, B. Van 164
- Frame, Beth 154–156
- Kripke 152–154
- vagueness 158–170
- Free complete atomic Boolean algebra
  - 287–324
- Free join semilattice 304–312
- Free lattice x, 258–267
- Freedom postulate 177
- Frege's reconstruction 42
- Function, assignment 4, 5, 26, 28, 128–135, 304, 320
- characteristic 70–120
- choice 55–69
- constant 37, 70–120
- domain 152
- identity 70–120, 258
- interpretation 28, 122, 156, 304, 319
- inverse 70–120
- n-place truth 128–135
- notion of 70–125
- space 51
- truth 128–135
- Functional completeness 128–135
- type logic 27
- Gallin, D. 28, 132
- Gamut, viii
- Generalized continuum hypothesis
  - 56–69

- theorem 65
- Cardinal number 54–69, 258, 259
- Cardinality, finite 21
  - infinite 21
  - notion of 54–69
- Cartesian product 49, 50, 51, 72
- Chain ix, 109–120
  - maximal ix
- Change, intervals events and 197–233
  - Kamp's logic of 221–233
  - moment of 211–218
- Characteristic function 70–120
- Chierchia, G. viii, x, 32, 44
- Choice, axiom of 55–69, 115
  - function 55–69
- Closure, deductive 10, 36
  - downward 202
- Co-final 103–120
- Co-initial 103–120
- Cohen's forcing technique 60
- Comb, Beth 156
- Common predecessor 334, 335
  - successor 334, 335
- Commutativity 236–283
- Compactness theorem 13, 19, 29, 31
- Complemented lattice 247–283
- Completeness 158
- Complete atomic Boolean algebra
  - 262–267, 284–324
  - lattice 269–283
    - bounded 265–267
- Completed change principle 233
- Completeness 326, 327
  - functional 128–135
  - notion of 256–267
  - theorem 13, 18, 19, 24, 29, 31, 36
- Comprehension axiom schema 26
- Comprehension, principle of 45
- Condition, atomicity 175
  - conjunction 174
  - direction 174
  - disjunction 174
- Congruence relation 95–105, 251–254, 352
- Congruence structure 95–105
- Conjunction 188–196, 288–299, 342
  - condition 174
- Connected relation 83–105
- Constant function 37, 70–120
  - individual 24, 37, 159
    - and predicate 33–44
  - logical 3, 7, 33–44
  - mathematical 33–44, 45
  - non-logical 4, 33–44
  - non-mathematical 33–44
  - non-primitive logical 7
  - predicate 37
- Constructions with partial orders
  - 171–196
- Context, background 160
  - set 99
- Continuum hypothesis 56–69
- Convex set 110–120
  - sublattice 267–283
- Convexity 188–196, 341
  - principle 176
- Cooper, R. 30
- Correctness theorem 29
- Countable model 60–69
- Cressell, M. 200, 208
- Dalen, D. van 145, 152
- Deductive closure 10, 36
- Definability, first order 1
- Definition, entailment 7
  - truth 5, 26
- Dense linear order 64
  - theory of 35–44
  - without endpoints 64–69
- Density 339
  - principle 176
- Diabolo 335, 336
- Diagram, Hasse 86
- Diamond 245, 265, 347, 348
- Direct interpretation 75
- Direction condition 174
- Discreet change principle 233
  - structure 138
- Discreetness 108–120
- Disjunction 288–299
  - condition 174
- Distance structure 104–120
- Distributive lattice 244–283
  - set theoretic representation of x
- Distributivity 314–324, 348
- Domain function 153

- Downward closure 202
  - entailing 118, 142–170, 337, 340
- Dowty, D. viii, 76, 200, 201, 204, 206, 210, 211, 213
- Duality 87–105, 235–283
- Dummett, M. 145
- Element, maximal 158, 181, 236, 305, 345
- Element, minimal 105–120, 139
- Embedding 76–120, 329, 330
- Empty set 51
  - axiom 48
- Eng, M. 121
- Endpoints, dense linear order without
  - 64–69
  - theory of 36–44
- Entailing, downward 118, 142–170, 337, 340
  - definition of 7
- Epimorphism 76–120, 329
  - strong 82–120
- Equivalence relation 84–105
  - partial order and 83–105
  - partition and ix
- Event model 226
  - structure x, 186–196
- Expressibility, algebraic semantics
  - functional completeness and 127–135
- Expression, semantics of x
- Extended 103–120
- Extension 97–105, 292
  - negative 158–170
  - positive 158–170
- Extensionality 327
- Faltz, L. 258, 284–324
- Filter 352, 353
  - ideals and 267–283
  - maximally proper 271–283
  - notion of 180–186
  - prime 271–283, 353, 354
  - proper 271–283, 354
  - representation ix
- Fine, K. 158, 162–165
- Finitary relation 72–120
- Finite cardinality 21
- intersection property 271–283
  - lattice 353
- Finiteness 327
- First order approximation 30
  - definability 1
  - language 2, 4, 10, 14, 17, 24
  - lexicon 2, 3
  - logic 1, 2–13, 20, 21, 24, 26
  - notion of 1
  - predicate logic 2, 7
  - quantification 31
  - sentence 10, 12, 22, 33
  - theory 1, 12, 19, 20, 32–44, 328
- Forcing technique, Cohen's 60
- Formalism, Hilbert's 43
- Formalization in semantics vii
- Formula induction 9
- Foundation axiom 53–69
- Fraassen, B. Van 164
- Frame, Beth 154–156
  - Kripke 152–154
  - vagueness 158–170
- Free complete atomic Boolean algebra
  - 287–324
- Free join semilattice 304–312
- Free lattice x, 258–267
- Freedom postulate 177
- Frege's reconstruction 42
- Function, assignment 4, 5, 26, 28, 128–135, 304, 320
  - characteristic 70–120
  - choice 55–69
  - constant 37, 70–120
  - domain 152
  - identity 70–120, 258
  - interpretation 28, 122, 156, 304, 319
  - inverse 70–120
  - n-place truth 128–135
  - notion of 70–125
  - space 51
  - truth 128–135
- Functional completeness 128–135
  - type logic 27
- Gallin, D. 28, 132
- Gamut, viii
- Generalized continuum hypothesis
  - 56–69

- quantifier theory 256–267
- Generated lattice *x*, 248–267
- Gödel's incompleteness theorem 29, 30, 42, 43, 44
- Grätzer, G. 70, 234, 245, 254, 258, 259
- Grammar, Montague 75, 97
- Graph 83–105
- Groenendijk, J. 97, 98, 101, 135
- Hasse diagram 86
- Heinämäki, O. 140
- Hendriks, H. 294, 295
- Henkin proof 14
  - theory 15–19
- Heyting algebra 153
- Hilbert's formalism 43
- Hinrichs, E. 135, 140, 200
- Hodges, W. 1
- Homomorphic image 259–267
- Homomorphism 74–120, 239, 251–254, 258, 293, 298, 299, 328, 329, 351, 352
  - bijjective 76–120, 329, 330
  - Boolean join 301–32
  - injective 76–120
  - natural 94–120
  - strong 76–120
- Humberstone, L. 201–211
  - deduction system 203
- Hypothesis, continuum 56–69
  - generalized continuum 56–69
  - induction 63–69
- Ideals, filters and 267–283
- Idempotency 236–283, 346, 348
- Identity axiom 8
  - function 70–120, 258
- Image, homomorphic 259–267
- Immediate predecessor 106–120
  - successor 106–120
- Incomparable 86–105
- Incompatibility principle 212–218, 221–233
- Incompleteness theorem 40, 42
- Inconsistency, notion of 19
- Individual and predicate constant 33–44
  - constant 24, 37, 159
  - level term, set of 25
  - variable 24
- Individuals, *n*-place predicate of
  - constant 24
  - predicate constant of *l*-place properties of 24
- Induction, formula 9
  - hypothesis 63–69
  - schema 38
  - step 9, 15, 63
- Inference rule 7, 8, 14, 40
- Infimum notion 234–283
- Infinite cardinality 21
  - structure 33
- Infinity axiom 58
- Information structure *ix*, 145–156
- Initial number 55–69
- Injection 185, 269, 280, 351
  - notion of 70–120
- Injective homomorphism 76–120
- Instant tense logic 121–127, 135–145
  - time structure *ix*
- Intension 97–105
- Interpretation, direct 75
  - function 28, 122, 156, 304, 319
- Intersection 89–105
- Interval semantics 197–201
  - logic of change in 201–211
  - time *ix*
- Intervals, events and change 197–233
- Intransitive relation 83–105
- Intuitionism 43
  - Brouwer 146
- Intuitionistic logic 145–152
- Inverse function 70–120
- Irreflexive relation 83–105
- Irreflexivity 189–196, 330, 341
- Isomorphism 61–69, 76–120, 185, 258–267, 290, 292, 297, 329, 330
  - partial 64–69
- Janssen, T. 76
- Join 234–283, 344
  - semilattice *x*, 242–283, 301–324, 346, 347
- Kadmon, N. *x*, 142

- Kamp, H. 133–135, 158, 162–166, 169–171, 186–196, 211
  - logic of change 221–233
- Kaplan, D. 135
- Keenan, E. 258, 284–324
- Kleene algebra 162
  - logic 161
  - truth table 214
- Krifka, M. 314
- Kripke, S. 145–156
  - frame 152–154
  - model 150–154
  - semantics for predicate logic 153–156
  - semantics for propositional logic 152, 153
- Ladusaw, W. 142
- Landman, F. 138, 142, 160, 168, 301, 303, 309, 322
- Language 303
  - first order 2, 4, 10, 14, 17, 24
  - second order 24–33
  - type logical 27
- Larson, R. 140
- Lattice 234–283
  - atomic 242–283
  - atomless 240–283
  - Boolean 247–283
  - bounded 240
    - distributive 246–283
  - complemented 247–283
  - complete 269–283
    - atomic bounded 265–267
  - distributive 244–283
  - finite 353
  - free *x*, 258–267
  - generated *x*, 248–267
  - Morgan 162
  - powerset 240–283
    - properties of *x*
    - set theoretic 240–283
  - representation theorem for distributive 279
  - semantics with 284–324
- Law, Leibniz' 27
  - Morgan's 247
- Lemma, Lindenbaum's 11, 18
- Zorn's 55, 115–120
- Lewis, D. 138, 167
- Lexicon, first order 2, 3
- Lindenbaum algebra 95, 283
  - lemma 11, 18
- Lindström's theorem 23, 29
- Linear order 84–105, 333
  - properties of *ix*
  - theory of 35–44
- Linearity 224, 332
  - principle 175
- Linebarger, M. 142
- Link, G. 138, 200, 255, 300–313, 317
- Löwenheim-Skolem theorem 13, 19–22, 24, 29, 31–33, 36, 40, 60–69
- Logic *ix*
  - first order 1, 2, 13, 20, 21, 24, 26, 27
  - instant tense 121–127, 135–145
  - intuitionistic 145–152
  - Kleene 161
  - minimal 148
  - minimal tense 122–127
  - Montague's intensional 132, 284–324
  - of change in interval semantics 201–211
  - of plurality and mass nouns 317–324
  - predicate 135
  - Priorian tense *ix*, 121–170, 199
  - propositional tense 134
  - second order 24–33
  - set theory and 1
  - sorted first order 30 31
  - tense *ix*
  - third order 27
  - unsorted first order 30
  - Veltman's data 168
- Logical constant 3, 7, 33–44
- Logicism 42
- Lower bound 234–283
- Mass nouns 312–324
  - logic of plurality and 317–324
- Mathematical and logical structures *viii*
  - techniques *viii*
- Mathematical constant 33–44, 45
  - construction, Cantor 36
- Maximal blockset 315–324
- Maximal chain *ix*
  - principle 116–120, 273–283

- element 158, 181, 236, 305, 345
- filters, representation through 180–186
- Maximally consistent theory 18
  - proper filter 271–283
- McConnell-Ginet, S. viii, x
- Meaning postulate 33–44, 160, 169
- Meet 234–283, 344
  - semilattice 242–283, 346, 347
- Metalogic ix, 13–24
- Meulen, A. ter viii
- Minimal element 105–120, 139
  - logic 148
  - tense logic 122–127
- Mönnich, U. 32
- Model, Beth 150–152, 154–156
  - countable 60–69
  - event 226
  - Kripke 150–154
  - theory 13
  - vagueness 167
- Modus ponens 149
- Modus tollens 149
- Moment of change 211–218
- Monotonicity 188–196, 214, 341
  - principle 176
- Montague, R. 76, 284–324
  - grammar 75, 97
  - intensional logic 132, 284–324
  - semantics 34, 59
- Morgan lattice 162
- Morgan's law 247
- n-place predicate constant 2
  - of individuals 24
  - predicate variable 24
  - quantifier 130–135
  - tense 131–135
  - truth function 128–135
- Natural deduction system 7
  - homomorphism 94–120
  - numbers, Von Neumann's set theoretic definition of 52
- Negation 288–299
- Negative extension 158–170
- Non-contingencies, set of 41
- Non-discreet structure 125
- Non-empty set 53
- Non-logical constant 4, 33–44
  - individual and predicate constant 3
- Non-mathematical constant 33–44
- Non-primitive logical constant 7
- Notion, bijection 70–120
  - cardinality 54–69
  - completeness 256–267
  - filter 180–186
  - first order 1
  - function 70–120
  - inconsistency 19
  - infimum 234–283
  - injection 70–120
  - relation 70–120
  - structure and homomorphism ix
  - supremum 234–283
  - surjection 70–120
  - vague 41
- Nouns, mass 313–324
- Number, cardinal 54–69, 258, 259
  - initial 55–69
  - ordinal 53–69, 334
  - theory 36–44
- Operator, Priorian tense 201–211
- Order, dense linear 64
  - linear 84–105, 333
  - theory of 124
- partial ix, 39, 42, 47, 70–120, 346
- total 84–105
- Ordinal number 53–69, 334
- Oversteegen, E. 138, 140
- Pair, axiom of 49, 50
- Paradox, Russell 42, 45, 59
  - Skolem's 60
- Parsons, T. 200
- Partee, B. viii, 121, 140, 166, 284, 288, 290
- Partial information and vagueness 156–170
  - isomorphism 64–69
  - order ix, 39, 42, 47, 70–120, 346
  - constructions with 171–196
  - equivalence relation and 83–105
  - strict 85–105
  - semantics with 121–170
  - set theoretic representation of ix

- theory of 34–44
- Partition 88–105, 315–324
  - equivalence relation and ix
- Peano arithmetics 29, 36–69
- Pelletier, J. 317
- Pentagon 245, 265
- Period structure 104–120, 171–186
- Peters, S. viii, 76
- Pinkal, M. 170
- Pluralization 306–312
- Plurals 300–312
  - mass nouns and, semantics of x
- Point generated period structure 179–186
  - properties of 182–186
- Positive extension 158–170
- Postulate, freedom 177
  - meaning 33–44, 160, 169
- Power set 50
  - Boolean algebra 267–283, 354
  - lattice 240–283
- Precisification 158
- Precision state 158–170
- Predecessor, common 334, 335
  - immediate 106–120
- Predicate constant 37
  - n-place 2
  - non-logical individual and 3
  - of 1-place properties of individuals 24
  - level term, set of 25
  - logic 135
    - first order 2, 7
    - Kripke semantics for 153–156
- Preorder 84–105
- Prime filter 271–283, 353, 354
- Principle, bivalence 211, 222–233
  - completed change 233
  - convexity 176
  - density 176
  - discreet change 233
  - incompatibility 212–218, 221–233
  - linearity 175
  - maximal chain 116–120, 273–283
  - monotonicity 176
  - of comprehension 45
  - separation 176, 224–233
  - transfer 176
- witness 176, 177
- Prior, A. 121, 127
- Priorian tense logic ix, 121–170, 199
  - tense operator 201–211
- Product, Cartesian 49, 50, 51, 72
- Proper filter 271–283, 354
- Properties of finite intersection 271–283
  - of lattices x
  - of linear order ix
  - of partial orders in temporal setting ix
  - of point 182–186
  - second order 26
- Propositional axiom 8
  - logic, Kripke semantics for 152, 153
  - tense logic 134
- Quantification, first order 31
  - second order 31
- Quantifier axiom 8
  - n-place 130–135
- Realism 43
- Reconstruction, Frege's 42
- Refinement 89–105, 160
- Reflexive relation 83–105
- Reflexivity 238, 331, 332, 346
- Reichenbach, H. 121
- Relation, antisymmetric 83–105
  - asymmetric 83–105
  - binary 70–120
  - congruence 95–105, 251–254, 352
  - connected 83–105
  - equivalence 84–105
  - finitary 72–120
  - intransitive 83–105
  - irreflexive 83–105
  - notion of 70–120
  - reflexive 83–105
  - symmetric 83–105
  - theory of 83–105
  - transitive 83–105
  - unconnected 86–105
- Relational algebra 72–120
  - structure 72–120
- Representation, filter ix
  - theorem for Boolean algebras 280
  - for complete atomic Boolean algebras 280–283

- for distributive lattices 279
- through maximal filters 180–186
- Rooth, M. 284, 288, 290, 294
- Rule, inference 7, 8, 14, 40
- Russell paradox 42, 45, 59
- Scha, R. 300
- Schema, axiom 7, 40, 46
  - comprehension axiom 26
  - induction 38
  - substitution 51
- Schubert, L. 317
- Scott, D. 152
- Second order language 24–33
  - logic 24–33
  - property 26
  - quantification 31
- Semantics 304–312, 319–324
  - Beth 154–156
  - Boolean 284–299
  - formalization in vii
  - interval 197–201
  - Montague 34, 59
  - of expression x
  - of plurals and mass nouns x
  - with lattices 284–324
  - with partial orders 121–170
- Semilattice 346, 347
  - free join 304–312
  - join x, 242–283, 301–324, 346, 347
  - meet 242–283, 346, 347
  - sub-join 305
- Semitree 333
  - structure 103–120
- Sentence 4
  - first order 10, 12, 22, 33
- Separation, axiom of 46
  - principle 176, 224–233
- Set, context 99
  - convex 110–120
  - empty 51
  - non-empty 53
  - of approximation 181–186
  - of individual level term 25
  - of non-contingencies 41
  - of predicate level term 25
  - power 50
  - successor 52
- theoretic lattice 240–283
  - representation of distributive lattices x
  - of partial orders ix
  - universe 52–69
- theory ix, 36–44
  - logic and 1
- Singularization 306–312
- Skolem's paradox 60
- Sorted first order logic 30, 31
- Soundness 326
- Space, function 51
- Stalnaker, R. 99, 135, 167
- State, precision 158–170
- Step, base 9, 15, 63
  - induction 9, 15, 63
- Stokhof, J. 97, 98, 101
- Stokhof, M. 135
- Stone's theorem 275–279
- Strict partial order 85–105
  - theory of 35–44
- Strong epimorphism 82–120
  - homomorphism 76–120
- Structure and homomorphism, notion of ix
  - congruence 95–105
  - discreet 138
  - distance 104–120
  - event x, 186–196
  - infinite 33
  - information ix, 145–156
  - instant time ix
  - mathematical and logical viii
  - non-discreet 125
  - period 104–120, 171–186
  - point generated period 179–186
  - relational 72–120
  - semitree 103–120
  - tree-like 103–120
  - witnessed period 178–186
- Sub-join semilattice 305
- Sublattice, convex 267–283
- Substitution schema 51
- Successor, common 334, 335
  - immediate 106–120
  - set 52
- Sum, axiom of 49, 50
- Supervaluation 164, 218

- structure ix
- Supremum notion 234–283
- Surjection 51–69, 185, 280, 351
  - notion of 70–120
- Symmetric relation 83–105
- Syntax 303, 304, 318–324
- System, Humberstone's deduction 203
  - natural deduction 7
- Table, Kleene truth 214
- Taylor, B. 201
- Techniques, mathematical and logical viii
- Temporal setting, properties of partial orders in ix
- Tense logic ix
  - n-place 131–135
- Theorem, Cantor's 65
  - compactness 19, 29, 31
  - completeness 13, 18, 19, 24, 29, 31, 36
  - correctness 29
  - Gödel's 29, 30, 42–44
  - Lindström's 23
  - Löwenheim-Skolem 13, 19–22, 24, 29, 31–33, 36, 40, 60–69
  - Stone's 275–279
- Theory, arbitrary consistent 18
  - axiomatic 7
  - Cantor set 45
  - first order 1, 12, 19, 20, 32–44, 328
  - generalized quantifier 256–267
  - Henkin 15–19
  - Lindström's 29
  - maximally consistent 18
  - model 13
  - number 36–44
  - of dense linear order 35–44
    - without endpoints 36–44
  - of linear order 35–44, 124
    - without endpoints 35–44
  - of partial order 34–44
  - of relations 83–105
  - of strict partial order 35–44
  - set ix, 36–44
  - type 59
  - Zermelo-Fraenkel set 44–69
- Third order logic 27
- Thomason, R. 101
- Tichy, P. 137
- Time interval ix
- Total order 84–105
- Transfer 224
  - argument 33
  - principle 176
- Transitive relation 83–105
- Transitivity 188–196, 238, 330–332, 338, 341, 346
- Tree-like structure 103–120
- Truth definition 5, 26
  - function 128–135
  - value 122, 128–135, 200, 202, 228–233
- Turner, R. 32
- Type logical language 27
  - theory 59
- Ultrafilter 271–283, 290, 291, 353, 354
- Unconnected relation 86–105
- Unification 90–105
- Universal algebra 70–120, 248–267
- Universe, set theoretic 52–69
- Unsorted first order logic 30
- Vague notion 41
- Vagueness 217, 227–233
  - frame 158–170
  - model 167
  - partial information and 156–170
- Value, truth 122, 128–135, 200, 202, 228–233
- Variable, individual 24
  - n-place predicate 24
- Veltman, F. ix, 168
  - data logic 168
- Vlach, F. 197
- Wall, R. viii, 76
- Well-foundedness 108–120, 126
- Whorf, B. 134
- Witness 17, 18, 314–324, 341
  - principle 176 177
  - period structure 178–186
- Zermelo-Fraenkel set theory 44–69
- Zorn's lemma 55, 115–120