

Name: _____

1. Problem

An experiment has $n_1 = 5$ plants in the treatment group and $n_2 = 3$ plants in the control group. After some time, the plants' heights (in cm) are measured, resulting in the following data:

	value1	value2	value3	value4	value5
sample 1:	9	14.7	8.3	11.4	8.5
sample 2:	18.4	17.9	15.4		

- Determine degrees of freedom.
- Determine t^* for a 96% confidence interval.
- Determine SE .
- Determine a lower bound of the 96% confidence interval of $\mu_2 - \mu_1$.
- Determine an upper bound of the 96% confidence interval of $\mu_2 - \mu_1$.
- Determine $|t_{\text{obs}}|$ under the null hypothesis $\mu_2 - \mu_1 = 0$.
- Determine a lower bound of the two-tail p -value.
- Determine an upper bound of two-tail p -value.
- Do you reject the null hypothesis with a two-tail test using a significance level $\alpha = 0.04$? (yes or no)

1. (a)

					2
--	--	--	--	--	---

 .

0	0	0
---	---	---

(b)

					4
--	--	--	--	--	---

 .

8	5	0
---	---	---

(c)

					1
--	--	--	--	--	---

 .

5	2	7
---	---	---

(d)

				-	0
--	--	--	--	---	---

 .

6	0	6
---	---	---

(e)

				1	4
--	--	--	--	---	---

 .

2	0	6
---	---	---

(f)

					4
--	--	--	--	--	---

 .

4	5	2
---	---	---

(g)

					0
--	--	--	--	--	---

 .

0	4	0
---	---	---

(h)

					0
--	--	--	--	--	---

 .

0	5	0
---	---	---

(i)

no

1. Solution

These data are unpaired. We might as well find the sample means and sample standard deviations (use a calculator's built-in function for standard deviation).

$$\bar{x}_1 = 10.4$$

$$\bar{x}_2 = 17.2$$

$$s_1 = 2.71$$

$$s_2 = 1.61$$

We make a conservative estimate of the degrees of freedom using the appropriate formula.

$$df = \min(n_1, n_2) - 1 = \min(5, 3) - 1 = 2$$

We use the t table to find t^* such that $P(|T| < t^*) = 0.96$

$$t^* = 4.85$$

We use the SE formula for unpaired data.

$$SE = \sqrt{\frac{(s_1)^2}{n_1} + \frac{(s_2)^2}{n_2}} = \sqrt{\frac{(2.71)^2}{5} + \frac{(1.61)^2}{3}} = 1.527$$

We find the bounds of the confidence interval.

$$CI = (\bar{x}_2 - \bar{x}_1) \pm t^* SE$$

$$CI = (-0.606, 14.206)$$

We find t_{obs} .

$$t_{\text{obs}} = \frac{(\bar{x}_2 - \bar{x}_1) - (\mu_2 - \mu_1)_0}{SE} = \frac{(17.2 - 10.4) - 0}{1.527} = 4.45$$

We find $|t_{\text{obs}}|$.

$$|t_{\text{obs}}| = 4.45$$

We use the table to determine bounds on p -value. Remember, $df = 2$ and $p\text{-value} = P(|T| > |t_{\text{obs}}|)$.

$$0.04 < p\text{-value} < 0.05$$

We should consider both comparisons to make our decision.

$$|t_{\text{obs}}| < t^*$$

$$p\text{-value} > \alpha$$

Thus, we retain the null hypothesis. Also notice the confidence interval does contain 0.

- (a) 2
- (b) 4.85
- (c) 1.527
- (d) -0.606
- (e) 14.206
- (f) 4.452
- (g) 0.04
- (h) 0.05
- (i) no