

Name: \_\_\_\_\_

1. Problem

An experiment has  $n_1 = 6$  plants in the treatment group and  $n_2 = 8$  plants in the control group. After some time, the plants' heights (in cm) are measured, resulting in the following data:

	value1	value2	value3	value4	value5	value6	value7	value8
sample 1:	0.85	1.01	1.02	1.29	1.2	0.87		
sample 2:	1.22	1.26	1.01	1.55	1.65	1.02	1.24	1.37

- (a) Determine degrees of freedom.
- (b) Determine  $t^*$  for a 98% confidence interval.
- (c) Determine  $SE$ .
- (d) Determine a lower bound of the 98% confidence interval of  $\mu_2 - \mu_1$ .
- (e) Determine an upper bound of the 98% confidence interval of  $\mu_2 - \mu_1$ .
- (f) Determine  $|t_{obs}|$  under the null hypothesis  $\mu_2 - \mu_1 = 0$ .
- (g) Determine a lower bound of the two-tail  $p$ -value.
- (h) Determine an upper bound of two-tail  $p$ -value.
- (i) Do you reject the null hypothesis with a two-tail test using a significance level  $\alpha = 0.02$ ? (yes or no)

1. (a) 

					5
--	--	--	--	--	---

 . 

0	0	0
---	---	---

(b) 

					3
--	--	--	--	--	---

 . 

3	6	0
---	---	---

(c) 

					0
--	--	--	--	--	---

 . 

1	0	8
---	---	---

(d) 

				-	0
--	--	--	--	---	---

 . 

1	1	3
---	---	---

(e) 

					0
--	--	--	--	--	---

 . 

6	1	3
---	---	---

(f) 

					2
--	--	--	--	--	---

 . 

3	1	5
---	---	---

(g) 

					0
--	--	--	--	--	---

 . 

0	5	0
---	---	---

(h) 

					0
--	--	--	--	--	---

 . 

1	0	0
---	---	---

(i) 

no
----

## 1. Solution

These data are unpaired. We might as well find the sample means and sample standard deviations (use a calculator's built-in function for standard deviation).

$$\bar{x}_1 = 1.04$$

$$\bar{x}_2 = 1.29$$

$$s_1 = 0.176$$

$$s_2 = 0.228$$

We make a conservative estimate of the degrees of freedom using the appropriate formula.

$$df = \min(n_1, n_2) - 1 = \min(6, 8) - 1 = 5$$

We use the  $t$  table to find  $t^*$  such that  $P(|T| < t^*) = 0.98$

$$t^* = 3.36$$

We use the  $SE$  formula for unpaired data.

$$SE = \sqrt{\frac{(s_1)^2}{n_1} + \frac{(s_2)^2}{n_2}} = \sqrt{\frac{(0.176)^2}{6} + \frac{(0.228)^2}{8}} = 0.108$$

We find the bounds of the confidence interval.

$$CI = (\bar{x}_2 - \bar{x}_1) \pm t^* SE$$

$$CI = (-0.113, 0.613)$$

We find  $t_{\text{obs}}$ .

$$t_{\text{obs}} = \frac{(\bar{x}_2 - \bar{x}_1) - (\mu_2 - \mu_1)_0}{SE} = \frac{(1.29 - 1.04) - 0}{0.108} = 2.32$$

We find  $|t_{\text{obs}}|$ .

$$|t_{\text{obs}}| = 2.32$$

We use the table to determine bounds on  $p$ -value. Remember,  $df = 5$  and  $p\text{-value} = P(|T| > |t_{\text{obs}}|)$ .

$$0.05 < p\text{-value} < 0.1$$

We should consider both comparisons to make our decision.

$$|t_{\text{obs}}| < t^*$$

$$p\text{-value} > \alpha$$

Thus, we retain the null hypothesis. Also notice the confidence interval does contain 0.

- (a) 5
- (b) 3.36
- (c) 0.108
- (d) -0.113
- (e) 0.613
- (f) 2.315
- (g) 0.05
- (h) 0.1
- (i) no