

Name: _____

1. Problem

An experiment has $n_1 = 6$ plants in the treatment group and $n_2 = 6$ plants in the control group. After some time, the plants' heights (in cm) are measured, resulting in the following data:

	value1	value2	value3	value4	value5	value6
sample 1:	139	127	120	142	119	142
sample 2:	111	98	94	81	67	125

- (a) Determine degrees of freedom.
- (b) Determine t^* for a 95% confidence interval.
- (c) Determine SE .
- (d) Determine a lower bound of the 95% confidence interval of $\mu_2 - \mu_1$.
- (e) Determine an upper bound of the 95% confidence interval of $\mu_2 - \mu_1$.
- (f) Determine $|t_{\text{obs}}|$ under the null hypothesis $\mu_2 - \mu_1 = 0$.
- (g) Determine a lower bound of the two-tail p -value.
- (h) Determine an upper bound of two-tail p -value.
- (i) Do you reject the null hypothesis with a two-tail test using a significance level $\alpha = 0.05$? (yes or no)

1. (a)

					5
--	--	--	--	--	---

 .

0	0	0
---	---	---

(b)

					2
--	--	--	--	--	---

 .

5	7	0
---	---	---

(c)

					9
--	--	--	--	--	---

 .

5	3	2
---	---	---

(d)

			-	6	0
--	--	--	---	---	---

 .

4	9	7
---	---	---

(e)

			-	1	1
--	--	--	---	---	---

 .

5	0	3
---	---	---

(f)

					3
--	--	--	--	--	---

 .

7	7	7
---	---	---

(g)

					0
--	--	--	--	--	---

 .

0	1	0
---	---	---

(h)

					0
--	--	--	--	--	---

 .

0	2	0
---	---	---

(i)

yes

1. Solution

These data are unpaired. We might as well find the sample means and sample standard deviations (use a calculator's built-in function for standard deviation).

$$\bar{x}_1 = 132$$

$$\bar{x}_2 = 96$$

$$s_1 = 10.8$$

$$s_2 = 20.7$$

We make a conservative estimate of the degrees of freedom using the appropriate formula.

$$df = \min(n_1, n_2) - 1 = \min(6, 6) - 1 = 5$$

We use the t table to find t^* such that $P(|T| < t^*) = 0.95$

$$t^* = 2.57$$

We use the SE formula for unpaired data.

$$SE = \sqrt{\frac{(s_1)^2}{n_1} + \frac{(s_2)^2}{n_2}} = \sqrt{\frac{(10.8)^2}{6} + \frac{(20.7)^2}{6}} = 9.532$$

We find the bounds of the confidence interval.

$$CI = (\bar{x}_2 - \bar{x}_1) \pm t^* SE$$

$$CI = (-60.497, -11.503)$$

We find t_{obs} .

$$t_{\text{obs}} = \frac{(\bar{x}_2 - \bar{x}_1) - (\mu_2 - \mu_1)_0}{SE} = \frac{(96 - 132) - 0}{9.532} = -3.78$$

We find $|t_{\text{obs}}|$.

$$|t_{\text{obs}}| = 3.78$$

We use the table to determine bounds on p -value. Remember, $df = 5$ and $p\text{-value} = P(|T| > |t_{\text{obs}}|)$.

$$0.01 < p\text{-value} < 0.02$$

We should consider both comparisons to make our decision.

$$|t_{\text{obs}}| > t^*$$

$$p\text{-value} < \alpha$$

Thus, we reject the null hypothesis. Also notice the confidence interval does not contain 0.

- (a) 5
- (b) 2.57
- (c) 9.532
- (d) -60.497
- (e) -11.503
- (f) 3.777
- (g) 0.01
- (h) 0.02
- (i) yes