


## Confidence intervals

---

# Confidence intervals

- A plausible range of values for the population parameter is called a *confidence interval*.
- Using only a sample statistic to estimate a parameter is like fishing in a murky lake with a spear, and using a confidence interval is like fishing with a net.



We can throw a spear where we saw a fish but we will probably miss. If we toss a net in that area, we have a good chance of catching the fish.



- If we report a point estimate, we probably won't hit the exact population parameter. If we report a range of plausible values we have a good shot at capturing the parameter.

Photos by Mark Fischer (<http://www.flickr.com/photos/fischerfotos/7439791462>) and Chris Penny

(<http://www.flickr.com/photos/clearlydived/7029109617>) on Flickr.

Remember **standard error**.

Remember **standard error**.

We have a population with a mean  $\mu$  and a standard deviation  $\sigma$ .

Remember **standard error**.

We have a population with a mean  $\mu$  and a standard deviation  $\sigma$ .

We take a sample of size  $n$  and find the sample mean  $\bar{x}$ .

Remember **standard error**.

We have a population with a mean  $\mu$  and a standard deviation  $\sigma$ .

We take a sample of size  $n$  and find the sample mean  $\bar{x}$ .

The standard error is calculated.

$$SE = \frac{\sigma}{\sqrt{n}}$$

Remember **standard error**.

We have a population with a mean  $\mu$  and a standard deviation  $\sigma$ .

We take a sample of size  $n$  and find the sample mean  $\bar{x}$ .

The standard error is calculated.

$$SE = \frac{\sigma}{\sqrt{n}}$$

There is a 68% chance that  $|\bar{x} - \mu| < SE$  and a 95% chance that  $|\bar{x} - \mu| < 2SE$ .

Remember **standard error**.

We have a population with a mean  $\mu$  and a standard deviation  $\sigma$ .

We take a sample of size  $n$  and find the sample mean  $\bar{x}$ .

The standard error is calculated.

$$SE = \frac{\sigma}{\sqrt{n}}$$


There is a 68% chance that  $|\bar{x} - \mu| < SE$  and a 95% chance that  $|\bar{x} - \mu| < 2SE$ .

A sampling distribution (distribution of sample means) is approximately normal!




# A running average


**A single Running Average (6-sided die)**




## Overlay of many Running Averages




68% of data is within  $\mu \pm SE$ . 95% of data is between  $\mu \pm 2SE$ .



### A single running proportion



## Overlay of many running proportions



## Standard error in practice: confidence interval

In practice, we do not know  $\mu$  or  $\sigma$ . We only have a single sample, which is characterized by  $\bar{x}$  and  $s$ .

## Standard error in practice: confidence interval

In practice, we do not know  $\mu$  or  $\sigma$ . We only have a single sample, which is characterized by  $\bar{x}$  and  $s$ .

We estimate that  $\sigma \approx s$ .

## Standard error in practice: confidence interval

In practice, we do not know  $\mu$  or  $\sigma$ . We only have a single sample, which is characterized by  $\bar{x}$  and  $s$ .

We estimate that  $\sigma \approx s$ .

We know that if we did repeat the sampling procedure many times, then 95% of the time  $|\bar{x} - \mu| < 2SE$ .



## Standard error in practice: confidence interval

In practice, we do not know  $\mu$  or  $\sigma$ . We only have a single sample, which is characterized by  $\bar{x}$  and  $s$ .

We estimate that  $\sigma \approx s$ .

We know that if we did repeat the sampling procedure many times, then 95% of the time  $|\bar{x} - \mu| < 2SE$ .

Thus, we say we are 95% confident that  $\mu$  is within 2 SEs of  $\bar{x}$ .

## Standard error in practice: confidence interval

In practice, we do not know  $\mu$  or  $\sigma$ . We only have a single sample, which is characterized by  $\bar{x}$  and  $s$ .

We estimate that  $\sigma \approx s$ .

We know that if we did repeat the sampling procedure many times, then 95% of the time  $|\bar{x} - \mu| < 2SE$ .

Thus, we say we are 95% confident that  $\mu$  is within 2 SEs of  $\bar{x}$ .

To make frequentists happy, we do not say there is a 95% chance that  $\mu$  is within 2 SEs of  $\bar{x}$ .

## Standard error in practice: confidence interval

In practice, we do not know  $\mu$  or  $\sigma$ . We only have a single sample, which is characterized by  $\bar{x}$  and  $s$ .

We estimate that  $\sigma \approx s$ .

We know that if we did repeat the sampling procedure many times, then 95% of the time  $|\bar{x} - \mu| < 2SE$ .

Thus, we say we are 95% confident that  $\mu$  is within 2 SEs of  $\bar{x}$ .

To make frequentists happy, we do not say there is a 95% chance that  $\mu$  is within 2 SEs of  $\bar{x}$ .

Confidence intervals only try to capture the population parameter. A confidence interval says nothing about the confidence of capturing individual observations, a proportion of the observations, or about capturing point estimates. Confidence intervals only attempt to capture population parameters.

## Average number of exclusive relationships

A random sample of 50 college students was asked how many exclusive relationships they have been in so far. This sample yielded a mean of 3.2 and a standard deviation of 1.74. Estimate the true average number of exclusive relationships using this sample.

## Average number of exclusive relationships

A random sample of 50 college students was asked how many exclusive relationships they have been in so far. This sample yielded a mean of 3.2 and a standard deviation of 1.74. Estimate the true average number of exclusive relationships using this sample.

$$\bar{x} = 3.2 \quad s = 1.74$$

## Average number of exclusive relationships

A random sample of 50 college students was asked how many exclusive relationships they have been in so far. This sample yielded a mean of 3.2 and a standard deviation of 1.74. Estimate the true average number of exclusive relationships using this sample.

$$\bar{x} = 3.2 \quad s = 1.74$$

The approximate 95% confidence interval is defined as

$$\text{point estimate} \pm 2 \times SE$$

## Average number of exclusive relationships

A random sample of 50 college students was asked how many exclusive relationships they have been in so far. This sample yielded a mean of 3.2 and a standard deviation of 1.74. Estimate the true average number of exclusive relationships using this sample.

$$\bar{x} = 3.2 \quad s = 1.74$$

The approximate 95% confidence interval is defined as

$$\text{point estimate} \pm 2 \times SE$$

$$SE = \frac{s}{\sqrt{n}} = \frac{1.74}{\sqrt{50}} \approx 0.25$$

## Average number of exclusive relationships

A random sample of 50 college students was asked how many exclusive relationships they have been in so far. This sample yielded a mean of 3.2 and a standard deviation of 1.74. Estimate the true average number of exclusive relationships using this sample.

$$\bar{x} = 3.2 \quad s = 1.74$$

The approximate 95% confidence interval is defined as

$$\text{point estimate} \pm 2 \times SE$$

$$SE = \frac{s}{\sqrt{n}} = \frac{1.74}{\sqrt{50}} \approx 0.25$$

$$\bar{x} \pm 2 \times SE = 3.2 \pm 2 \times 0.25$$



## Average number of exclusive relationships

A random sample of 50 college students was asked how many exclusive relationships they have been in so far. This sample yielded a mean of 3.2 and a standard deviation of 1.74. Estimate the true average number of exclusive relationships using this sample.

$$\bar{x} = 3.2 \quad s = 1.74$$

The approximate 95% confidence interval is defined as

$$\text{point estimate} \pm 2 \times SE$$

$$SE = \frac{s}{\sqrt{n}} = \frac{1.74}{\sqrt{50}} \approx 0.25$$

$$\begin{aligned}\bar{x} \pm 2 \times SE &= 3.2 \pm 2 \times 0.25 \\ &= (3.2 - 0.5, 3.2 + 0.5)\end{aligned}$$

## Average number of exclusive relationships

A random sample of 50 college students was asked how many exclusive relationships they have been in so far. This sample yielded a mean of 3.2 and a standard deviation of 1.74. Estimate the true average number of exclusive relationships using this sample.

$$\bar{x} = 3.2 \quad s = 1.74$$

The approximate 95% confidence interval is defined as

$$\text{point estimate} \pm 2 \times SE$$

$$SE = \frac{s}{\sqrt{n}} = \frac{1.74}{\sqrt{50}} \approx 0.25$$

$$\begin{aligned}\bar{x} \pm 2 \times SE &= 3.2 \pm 2 \times 0.25 \\ &= (3.2 - 0.5, 3.2 + 0.5) \\ &= (2.7, 3.7)\end{aligned}$$

Which of the following is the correct interpretation of this confidence interval?

We are 95% confident that

- (a) the average number of exclusive relationships college students in this sample have been in is between 2.7 and 3.7.
- (b) college students on average have been in between 2.7 and 3.7 exclusive relationships.
- (c) a randomly chosen college student has been in 2.7 to 3.7 exclusive relationships.
- (d) 95% of college students have been in 2.7 to 3.7 exclusive relationships.

Which of the following is the correct interpretation of this confidence interval?

We are 95% confident that

- (a) the average number of exclusive relationships college students in this sample have been in is between 2.7 and 3.7.
- (b) *college students on average have been in between 2.7 and 3.7 exclusive relationships.*
- (c) a randomly chosen college student has been in 2.7 to 3.7 exclusive relationships.
- (d) 95% of college students have been in 2.7 to 3.7 exclusive relationships.

## A more accurate interval

Confidence interval, a general formula

$$\textit{point estimate} \pm z^{\star} \times SE$$

## A more accurate interval

Confidence interval, a general formula

$$\text{point estimate} \pm z^{\star} \times SE$$

Conditions when the point estimate =  $\bar{x}$ :

1. *Independence*: Observations in the sample must be independent
  - random sample/assignment
  - if sampling without replacement,  $n < 10\%$  of population
2. *Sample size / skew*:  $n \geq 30$  and population distribution should not be extremely skewed

## A more accurate interval

Confidence interval, a general formula

$$\text{point estimate} \pm z^{\star} \times SE$$


Conditions when the point estimate =  $\bar{x}$ :

1. *Independence*: Observations in the sample must be independent
  - random sample/assignment
  - if sampling without replacement,  $n < 10\%$  of population
2. *Sample size / skew*:  $n \geq 30$  and population distribution should not be extremely skewed

*Note*: We will discuss working with samples where  $n < 30$  in the next chapter.


## What does 95% confident mean?

- Suppose we took many samples and built a confidence interval from each sample using the equation  $point\ estimate \pm 2 \times SE$ .
- Then about 95% of those intervals would contain the true population mean ( $\mu$ ).
- The figure shows this process with 25 samples, where 24 of the resulting confidence intervals contain the true average number of exclusive relationships, and one does not.





## Same idea shown from running averages view.



About 95% of the time the population mean is within 2 SE of the sample mean.

## Width of an interval

If we want to be more certain that we capture the population parameter, i.e. increase our confidence level, should we use a wider interval or a smaller interval?

## Width of an interval

If we want to be more certain that we capture the population parameter, i.e. increase our confidence level, should we use a wider interval or a smaller interval?


*A wider interval.*

# Width of an interval

If we want to be more certain that we capture the population parameter, i.e. increase our confidence level, should we use a wider interval or a smaller interval?

*A wider interval.*

Can you see any drawbacks to using a wider interval?




# Width of an interval

If we want to be more certain that we capture the population parameter, i.e. increase our confidence level, should we use a wider interval or a smaller interval?

*A wider interval.*

Can you see any drawbacks to using a wider interval?



*If the interval is too wide it may not be very informative.* Image source:

[http://web.as.uky.edu/statistics/users/eao227/misc/garfield\\_weather.gif](http://web.as.uky.edu/statistics/users/eao227/misc/garfield_weather.gif)

## Changing the confidence level

$$\text{point estimate} \pm z^{\star} \times SE$$

- In a confidence interval,  $z^{\star} \times SE$  is called the *margin of error*, and for a given sample, the margin of error changes as the confidence level changes.
- In order to change the confidence level we need to adjust  $z^{\star}$  in the above formula.
- Commonly used confidence levels in practice are 90%, 95%, 98%, and 99%.
- For a 95% confidence interval,  $z^{\star} = 1.96$ .
- However, using the standard normal ( $z$ ) distribution, it is possible to find the appropriate  $z^{\star}$  for any confidence level.

Which of the below Z scores is the appropriate  $z^*$  when calculating a 98% confidence interval?

(a)  $Z = 2.05$

(b)  $Z = 1.96$

(c)  $Z = 2.33$

(d)  $Z = -2.33$

(e)  $Z = -1.65$

Which of the below Z scores is the appropriate  $z^*$  when calculating a 98% confidence interval?

(a)  $Z = 2.05$

(d)  $Z = -2.33$

(b)  $Z = 1.96$

(e)  $Z = -1.65$

(c)  $Z = 2.33$

