
Let *X* be a normal random variable with mean μ = 50 and standard deviation σ = 1.

$$X \sim \mathcal{N}(50, 1)$$

The figure below shows the density of random variable X. Each grid square represents 1% of probability.




- (a) Estimate P(X < 49) by shading and counting.
- (b) Determine P(X < 49) by using the z-table.

2. Problem

Let *X* be a normal random variable with mean μ = 35 and standard deviation σ = 5.

$$X \sim \mathcal{N}(35, 5)$$




- (a) Estimate P(X > 38) by shading and counting.
- (b) Determine P(X > 38) by using the z-table.

Let *X* be a normal random variable with mean μ = 36 and standard deviation σ = 2.

$$X \sim \mathcal{N}(36, 2)$$

The figure below shows the density of random variable X. Each grid square represents 1% of probability.




- (a) Estimate $P(|X \mu| < 2.8)$ by shading and counting.
- (b) Determine $P(|X \mu| < 2.8)$ by using the *z*-table.

4. Problem

Let *X* be a normal random variable with mean μ = 58 and standard deviation σ = 19.

$$X$$
 ∼ \mathcal{N} (58, 19)




- (a) Estimate $P(|X \mu| > 3.8)$ by shading and counting.
- (b) Determine $P(|X \mu| > 3.8)$ by using the z-table.

Let *X* be a normal random variable with mean μ = 60 and standard deviation σ = 18.

X ∼
$$\mathcal{N}$$
(60, 18)

The figure below shows the density of random variable X. Each grid square represents 1% of probability.




- (a) Estimate x such that P(X < x) = 0.88 by shading and counting.
- (b) Determine x such that P(X < x) = 0.88 by using the z-table.

6. Problem

Let *X* be a normal random variable with mean μ = 15 and standard deviation σ = 5.

$$X \sim \mathcal{N}(15, 5)$$




- (a) Estimate x such that P(X > x) = 0.27 by shading and counting.
- (b) Determine z such that P(X > x) = 0.27 by using the z-table.

Let *X* be a normal random variable with mean μ = 50 and standard deviation σ = 17.

X ∼
$$\mathcal{N}$$
(50, 17)

The figure below shows the density of random variable X. Each grid square represents 1% of probability.




- (a) Estimate a such that $P(|X \mu| < a) = 0.31$ by shading and counting.
- (b) Determine a such that $P(|X \mu| < a) = 0.31$ by using the z-table.

8. Problem

Let X be a normal random variable with mean μ = 58 and standard deviation σ = 8.

$$X \sim \mathcal{N}(58, 8)$$




- (a) Estimate a such that $P(|X \mu| > a) = 0.23$ by shading and counting.
- (b) Determine z such that $P(|X \mu| > a) = 0.23$ by using the z-table.

Let *X* be a normal random variable with mean μ = 25 and standard deviation σ = 8.

X ∼
$$N$$
(25, 8)

The figure below shows the density of random variable X. Each grid square represents 1% of probability.




- (a) Estimate x such that P(X > x) = 0.79 by shading and counting.
- (b) Determine z such that P(X > x) = 0.79 by using the z-table.

10. Problem

Let *X* be a normal random variable with mean μ = 57 and standard deviation σ = 13.

X ∼
$$\mathcal{N}$$
(57, 13)




- (a) Estimate $P(|X \mu| < 7.8)$ by shading and counting.
- (b) Determine $P(|X \mu| < 7.8)$ by using the *z*-table.

Let *X* be a normal random variable with mean μ = 16 and standard deviation σ = 3.

X ∼
$$N$$
(16, 3)

The figure below shows the density of random variable X. Each grid square represents 1% of probability.




- (a) Estimate x such that P(X < x) = 0.79 by shading and counting.
- (b) Determine x such that P(X < x) = 0.79 by using the z-table.

12. Problem

Let *X* be a normal random variable with mean $\mu = 17$ and standard deviation $\sigma = 1$.

$$X \sim \mathcal{N}(17, 1)$$




- (a) Estimate P(X > 18.2) by shading and counting.
- (b) Determine P(X > 18.2) by using the z-table.

Let *X* be a normal random variable with mean μ = 29 and standard deviation σ = 10.

X ∼
$$\mathcal{N}$$
(29, 10)

The figure below shows the density of random variable X. Each grid square represents 1% of probability.




- (a) Estimate a such that $P(|X \mu| > a) = 0.07$ by shading and counting.
- (b) Determine z such that $P(|X \mu| > a) = 0.07$ by using the z-table.

14. Problem

Let *X* be a normal random variable with mean μ = 58 and standard deviation σ = 17.

$$X$$
 ∼ \mathcal{N} (58, 17)




- (a) Estimate a such that $P(|X \mu| < a) = 0.45$ by shading and counting.
- (b) Determine a such that $P(|X \mu| < a) = 0.45$ by using the z-table.

Let *X* be a normal random variable with mean μ = 32 and standard deviation σ = 6.

X ∼
$$N$$
(32, 6)

The figure below shows the density of random variable X. Each grid square represents 1% of probability.




- (a) Estimate $P(|X \mu| > 1.2)$ by shading and counting.
- (b) Determine $P(|X \mu| > 1.2)$ by using the *z*-table.

16. Problem

Let *X* be a normal random variable with mean μ = 52 and standard deviation σ = 4.

$$X \sim \mathcal{N}(52, 4)$$




- (a) Estimate P(X < 48.8) by shading and counting.
- (b) Determine P(X < 48.8) by using the z-table.

Let *X* be a normal random variable with mean μ = 11 and standard deviation σ = 1.

$$X \sim \mathcal{N}(11, 1)$$

The figure below shows the density of random variable X. Each grid square represents 1% of probability.




- (a) Estimate x such that P(X > x) = 0.42 by shading and counting.
- (b) Determine z such that P(X > x) = 0.42 by using the z-table.

18. Problem

Let *X* be a normal random variable with mean μ = 48 and standard deviation σ = 9.

X ∼
$$N$$
(48, 9)




- (a) Estimate a such that $P(|X \mu| > a) = 0.07$ by shading and counting.
- (b) Determine z such that $P(|X \mu| > a) = 0.07$ by using the z-table.

Let *X* be a normal random variable with mean μ = 50 and standard deviation σ = 10.

X ∼
$$\mathcal{N}$$
(50, 10)

The figure below shows the density of random variable X. Each grid square represents 1% of probability.




- (a) Estimate x such that P(X < x) = 0.27 by shading and counting.
- (b) Determine x such that P(X < x) = 0.27 by using the z-table.

20. Problem

Let *X* be a normal random variable with mean μ = 44 and standard deviation σ = 7.

$$X \sim \mathcal{N}(44, 7)$$




- (a) Estimate $P(|X \mu| > 9.8)$ by shading and counting.
- (b) Determine $P(|X \mu| > 9.8)$ by using the *z*-table.

Let *X* be a normal random variable with mean μ = 49 and standard deviation σ = 2.

$$X \sim \mathcal{N}(49, 2)$$

The figure below shows the density of random variable X. Each grid square represents 1% of probability.




- (a) Estimate P(X < 47.4) by shading and counting.
- (b) Determine P(X < 47.4) by using the z-table.

22. Problem

Let *X* be a normal random variable with mean μ = 21 and standard deviation σ = 6.

X ∼
$$N$$
(21, 6)




- (a) Estimate a such that $P(|X \mu| < a) = 0.77$ by shading and counting.
- (b) Determine a such that $P(|X \mu| < a) = 0.77$ by using the z-table.

Let *X* be a normal random variable with mean μ = 28 and standard deviation σ = 5.

$$X \sim \mathcal{N}(28, 5)$$

The figure below shows the density of random variable X. Each grid square represents 1% of probability.




- (a) Estimate $P(|X \mu| < 7)$ by shading and counting.
- (b) Determine $P(|X \mu| < 7)$ by using the *z*-table.

24. Problem

Let *X* be a normal random variable with mean μ = 43 and standard deviation σ = 4.

$$X \sim \mathcal{N}(43, 4)$$




- (a) Estimate P(X > 41.4) by shading and counting.
- (b) Determine P(X > 41.4) by using the z-table.

1. (a) You will want a z-score.

$$z = \frac{x - \mu}{\sigma} = -1$$


(b) The shaded region is shown below.



You should count about 16 shaded squares, giving a probability of about 0.16.


(c) The probability is 0.1587.

2. (a) The shaded region is shown below.




You should count about 27 shaded squares, giving a probability of about 0.27.

(b) The probability is 0.2743.




You should count about 84 shaded squares, giving a probability of about 0.84.

- (b) The probability is 0.8385.
- 4. (a) The shaded regions are shown below.




You should count about 84 shaded squares, giving a probability of about 0.84.

- (b) The probability is 0.8415.
- 5. (a) The shaded region is shown below.



When you have shaded 88 squares, starting on the left, you should end around x = 81.6.


(b) $x \approx 81.06$



When you have shaded 27 squares, starting on the right, you should end around x = 18.


(b)
$$x = 11.95$$

7. (a) The shaded region is shown below.



When you have shaded 31 squares, starting in the middle, you should end near x = 56.8, giving a = 6.8.


(b)
$$a = 6.8$$



When you have shaded 23 squares, starting at both tails, you should end near x = 67.6. Really, you want to shade 11.5 squares starting from the left and also 11.5 squares starting from the right. This gives a = 9.6.


(b)
$$a = 9.6$$

9. (a) The shaded region is shown below.




When you have shaded 79 squares, starting on the right, you should end around x = 18.6.

(b)
$$x = 31.48$$




You should count about 45 shaded squares, giving a probability of about 0.45.

- (b) The probability is 0.4515.
- 11. (a) The shaded region is shown below.




When you have shaded 79 squares, starting on the left, you should end around x = 18.4.

- (b) $x \approx 18.43$
- 12. (a) The shaded region is shown below.



You should count about 12 shaded squares, giving a probability of about 0.12.


(b) The probability is 0.1151.



When you have shaded 7 squares, starting at both tails, you should end near x = 47. Really, you want to shade 3.5 squares starting from the left and also 3.5 squares starting from the right. This gives a = 18.


(b)
$$a = 18.1$$

14. (a) The shaded region is shown below.



When you have shaded 45 squares, starting in the middle, you should end near x = 68.2, giving a = 10.2.

(b)
$$a = 10.2$$




You should count about 84 shaded squares, giving a probability of about 0.84.

(b) The probability is 0.8415.

16. (a) You will want a z-score.


$$Z = \frac{x - \mu}{\sigma} = -0.8$$

(b) The shaded region is shown below.



You should count about 21 shaded squares, giving a probability of about 0.21.


(c) The probability is 0.2119.



When you have shaded 42 squares, starting on the right, you should end around x = 11.2.


(b)
$$x = 10.8$$

18. (a) The shaded regions are shown below.



When you have shaded 7 squares, starting at both tails, you should end near x = 64.2. Really, you want to shade 3.5 squares starting from the left and also 3.5 squares starting from the right. This gives a = 16.2.


(b)
$$a = 16.29$$



When you have shaded 27 squares, starting on the left, you should end around x = 44.

(b)
$$x \approx 43.9$$

20. (a) The shaded regions are shown below.




You should count about 16 shaded squares, giving a probability of about 0.16.

(b) The probability is 0.1615.

21. (a) You will want a z-score.

$$Z = \frac{X - \mu}{\sigma} = -0.8$$


(b) The shaded region is shown below.



You should count about 21 shaded squares, giving a probability of about 0.21.


(c) The probability is 0.2119.

22. (a) The shaded region is shown below.



When you have shaded 77 squares, starting in the middle, you should end near x = 28.2, giving a = 7.2.


(b) a = 7.2



You should count about 84 shaded squares, giving a probability of about 0.84.

(b) The probability is 0.8385.

24. (a) The shaded region is shown below.



You should count about 66 shaded squares, giving a probability of about 0.66.

(b) The probability is 0.6554.