

1. Problem:

Your boss wants to know what proportion of a very large population is special. You already know the proportion approximately 0.36. But, your boss wants to guarantee that the margin of error of a 95% confidence interval will be less than 0.004 (which is 0.4 percentage points). How large of a sample is needed?

Solution: We are given the confidence level and the margin of error.

$$\begin{aligned}\gamma &= 0.95 \\ ME &= 0.004\end{aligned}$$

Determine z^* such that $P(|Z| < z^*) = 0.95$. It is helpful to get the percentile of z^* by using $\ell = \frac{1+\gamma}{2} = \frac{1+0.95}{2} = 0.975$. This lets you find z^* such that $P(Z < z^*) = 0.975$.

$$z^* = 1.96$$

Use the appropriate formula.

$$\begin{aligned}n &= \hat{p}(1 - \hat{p}) \left(\frac{z^*}{ME} \right)^2 \\ &= (0.36)(0.64) \left(\frac{1.96}{0.004} \right)^2 \\ &= 55319.04\end{aligned}$$

When determining a necessary sample size, always round up (ceiling).

$$n = 55320$$

2. Problem:

A fair 8-sided die has a discrete uniform distribution with an expected value of $\mu = 4.5$ and a standard deviation $\sigma = 2.29$.

You are told to check if a 8-sided die has an expected value different than 4.5. You are told to roll the die 151 times and do a significance test with a significance level of 0.01.

You then roll the die 151 times and get a mean of 4.011. Should we conclude the die is unfair?

- (a) What kind of hypothesis test is appropriate?
- (b) State the hypotheses
- (c) Determine the p -value.
- (d) Decide whether we reject or retain the null hypothesis.
- (e) Do we conclude the die is unfair?

Solution: We should use a two-tail test of population mean.

State the hypotheses:


$$H_0 \text{ claims } \mu = 4.5$$

$$H_A \text{ claims } \mu \neq 4.5$$

Find the standard error.

$$\sigma_{\bar{x}} = \frac{\sigma}{\sqrt{n}} = \frac{2.29}{\sqrt{151}} = 0.186$$

Make a sketch of the null's sampling distribution.



Find the z score.

$$z = \frac{\bar{x} - \mu_0}{\sigma_{\bar{x}}} = \frac{4.011 - 4.5}{0.186} = -2.62$$

Find the p -value (using formula for left-tail test of mean).

$$\begin{aligned} p\text{-value} &= P(|Z| > 2.62) \\ &= 2 \cdot P(Z < -2.62) \\ &= \boxed{0.0088} \end{aligned}$$

Compare the p -value and the significance level ($\alpha = 0.01$).

$$p\text{-value} < \alpha$$

Yes, we reject the null hypothesis.

We conclude the die is unfair.

- (a) Right-tail single mean test
- (b) Hypotheses: H_0 claims $\mu = 4.5$ and H_A claims $\mu \neq 4.5$.
- (c) p -value = 0.0088
- (d) Yes, we reject the null hypothesis.
- (e) We conclude the die is unfair.

3. Problem:

A traffic engineer wants to determine the average speed of cars on the Longfellow Bridge. She knows the population standard deviation of speeds is 18 mph. To determine a 86% confidence interval with a margin of error of 1 mph, what sample size is needed?

Solution: We are given the population standard deviation, confidence level, and margin of error.

$$\sigma = 18$$

$$\gamma = 0.86$$

$$ME = 1$$

Determine the critical z value, z^* , such that $P(|Z| < z^*) = 0.86$. Remember, $\ell = \frac{\gamma+1}{2} = \frac{0.86+1}{2} = 0.93$

$$z^* = 1.48$$

Use the formula for sample size.

$$n = \left(\frac{z^* \sigma}{ME} \right)^2$$

$$= \left(\frac{(1.48)(18)}{1} \right)^2$$

$$= 709.6896$$

Round up.

$$n = 710$$

4. Problem:

Brahim wants to estimate the average mass of the beans in a large bag. Somehow, Brahim is certain that the standard deviation of the beans in the bag is 58 milligrams. He takes a sample of size 132 and finds the sample mean to be 491 milligrams. What would be the 98% confidence interval?

Solution: We are given the sample size, sample mean, population standard deviation, and confidence level.

$$n = 132$$

$$\bar{x} = 491$$

$$\sigma = 58$$

$$\gamma = 0.98$$

Determine the critical z value, z^* , such that $P(|Z| < z^*) = 0.98$. Remember, $\ell = \frac{\gamma+1}{2} = \frac{0.98+1}{2} = 0.99$

$$z^* = 2.33$$

Use the formula for bounds (mean, σ known).

$$\begin{aligned} LB &= \bar{x} - z^* \frac{\sigma}{\sqrt{n}} & UB &= \bar{x} + z^* \frac{\sigma}{\sqrt{n}} \\ &= 491 - 2.33 \times \frac{58}{\sqrt{132}} & &= 491 + 2.33 \times \frac{58}{\sqrt{132}} \\ &= 479.24 & &= 502.76 \end{aligned}$$

We are 98% confident that the population mean is between 479.24 and 502.76 milligrams.

$$CI = (479.24, 502.76)$$

5. Problem:

A student is taking a multiple choice test with 700 questions. Each question has 5 choices. You want to detect whether the student does better than random guessing, so you decide to run a hypothesis test with a significance level of 0.02.

Then, the student takes the test and gets 159 questions correct.

- (a) What kind of hypothesis test is appropriate?
- (b) State the hypotheses
- (c) Determine the p -value.
- (d) Decide whether we reject or retain the null hypothesis.
- (e) Do we think the student did significantly better than random guessing?

Solution: This is a right-tail (one-tail) proportion test because we only care whether the student does better than random.

Determine the null population proportion.

$$p_0 = \frac{1}{5} = 0.2$$

State the hypotheses.

$$H_0 \text{ claims } p = 0.2$$

$$H_A \text{ claims } p > 0.2$$

Determine the standard error.

$$\sigma_{\hat{p}} = \sqrt{\frac{p_0(1-p_0)}{n}} = \sqrt{\frac{0.2(1-0.2)}{700}} = 0.0151$$


Determine the sample proportion.

$$\hat{p} = \frac{159}{700} = 0.227$$

Determine a z score. For simplicity, we ignore the continuity correction.

$$z = \frac{\hat{p} - p_0}{\sigma_{\hat{p}}} = \frac{0.227 - 0.2}{0.0151} = 1.79$$

Make a sketch of the null's sampling distribution. The p -value is a right area.



.image

To determine that right area, we use the z table.

$$\begin{aligned} p\text{-value} &= P(\hat{p} > 0.227) \\ &= P(Z > 1.79) \\ &= 1 - P(Z < 1.79) \\ &= 0.0367 \end{aligned}$$

Compare p -value to α (which is 0.02).

$$p\text{-value} > \alpha$$

Make the conclusion: we retain the null hypothesis.

We think the student might just be guessing.

- Right tail (one-tail) proportion test
- Hypotheses: H_0 claims $p = 0.2$ and H_A claims $p > 0.2$.
- The p -value is 0.0367
- We retain the null hypothesis.
- We think the student might just be guessing.

6. Problem:

A null hypothesis claims a population has a mean $\mu = 100$ and a standard deviation $\sigma = 22$. You decide to run one-tail test on a sample of size $n = 286$ using a significance level $\alpha = 0.1$ to detect if the actual population mean is more than 100. You then collect the sample and find it has mean $\bar{x} = 101.27$.

- (a) What kind of hypothesis test is appropriate?
- (b) State the hypotheses
- (c) Determine the p -value.
- (d) Decide whether we reject or retain the null hypothesis.

Solution: We should use a right-tail test of population mean.

State the hypotheses:


$$H_0 \text{ claims } \mu = 100$$

$$H_A \text{ claims } \mu > 100$$

Find the standard error.

$$\sigma_{\bar{x}} = \frac{\sigma}{\sqrt{n}} = \frac{22}{\sqrt{286}} = 1.301$$

Make a sketch of the null's sampling distribution.



Find the z score.

$$z = \frac{\bar{x} - \mu_0}{\sigma_{\bar{x}}} = \frac{101.27 - 100}{1.301} = 0.98$$

Find the p -value (using formula for left-tail test of mean).

$$\begin{aligned} p\text{-value} &= P(Z > 0.98) \\ &= 1 - P(Z < 0.98) \\ &= 1 - 0.8365 \\ &= \boxed{0.1635} \end{aligned}$$

Compare the p -value and the significance level ($\alpha = 0.1$).

$$p\text{-value} > \alpha$$

No, we do not reject the null hypothesis.

- (a) Right-tail single mean test
- (b) Hypotheses: H_0 claims $\mu = 100$ and H_A claims $\mu < 100$.
- (c) p -value = 0.1635
- (d) No, we do not reject the null hypothesis.

7. Problem:

A new virus has been devastating corn production. When exposed, 44.7% of common seedlings die within a week. We are trying to develop a resistant strain of corn.

When we expose 700 seedlings of our strain to the virus, 41.1% die within a week. Using a significance level of 0.05, can we conclude that our strain is significantly more resistant?

- (a) What kind of hypothesis test is appropriate?
- (b) State the hypotheses
- (c) Determine the p -value.
- (d) Decide whether we reject or retain the null hypothesis.
- (e) Do we think our strain is significantly more resistant?

Solution: This is a left-tail (one-tail) proportion test because we only care whether a lower percentage of seedlings will die.

State the hypotheses.

$$H_0 \text{ claims } p = 0.447$$

$$H_A \text{ claims } p < 0.447$$


Determine the standard error.

$$\sigma_{\hat{p}} = \sqrt{\frac{p_0(1-p_0)}{n}} = \sqrt{\frac{0.447(1-0.447)}{700}} = 0.0188$$

Determine a z score. For simplicity, we ignore the continuity correction.

$$z = \frac{\hat{p} - p_0}{\sigma_{\hat{p}}} = \frac{0.411 - 0.447}{0.0188} = -1.91$$

Make a sketch of the null's sampling distribution. The p -value is a left area.



To determine that left area, we use the z table.

$$\begin{aligned} p\text{-value} &= P(\hat{p} < 0.411) \\ &= P(Z < -1.91) \\ &= 0.0281 \end{aligned}$$

Compare p -value to α (which is 0.05).

$$p\text{-value} < \alpha$$

Make the conclusion: we reject the null hypothesis.

We think our strain is more resistant than common corn.

- (a) Left-tail (one-tail) proportion test
- (b) Hypotheses: H_0 claims $p = 0.447$ and H_A claims $p < 0.447$.
- (c) The p -value is 0.0281
- (d) We reject the null hypothesis.
- (e) We think our strain is more resistant than common corn.

8. Problem:

Some snails have clockwise shells, and others have counter-clockwise shells. You want to construct a 80% confidence interval of the population proportion with clockwise shells. You hope the margin of error will be about 0.01. How many snails do you need?

Solution: We are given the confidence level and the margin of error.

$$\gamma = 0.8$$

$$ME = 0.01$$

Determine z^* such that $P(|Z| < z^*) = 0.8$. It is helpful to get the percentile of z^* by using $\ell = \frac{1+\gamma}{2} = \frac{1+0.8}{2} = 0.9$. This lets you find z^* such that $P(Z < z^*) = 0.9$.

$$z^* = 1.28$$

Use the appropriate formula.

$$n = \frac{1}{4} \left(\frac{z^*}{ME} \right)^2$$

$$= \frac{1}{4} \left(\frac{1.28}{0.01} \right)^2$$

$$= 4096$$

When determining a necessary sample size, always round up (ceiling).

$$n = 4096$$

9. Problem:

As an ornithologist, you wish to determine the average body mass of *Catharus fuscescens*. You randomly sample 16 adults of *Catharus fuscescens*, resulting in a sample mean of 41.52 grams and a sample standard deviation of 3.32 grams. Determine a 98% confidence interval of the true population mean.

Solution: We are given the sample size, sample mean, sample standard deviation, and confidence level.

$$n = 16$$

$$\bar{x} = 41.52$$

$$s = 3.32$$

$$\gamma = 0.98$$

Find the degrees of freedom.

$$df = n - 1$$

$$= 16 - 1$$

$$= 15$$

Determine the critical t value, t^* , such that $P(|T| < t^*) = 0.98$ and $df = 15$.

$$t^* = 2.6$$

Use the formula for bounds (mean, σ unknown).

$$LB = \bar{x} - t^* \frac{s}{\sqrt{n}}$$

$$= 41.52 - 2.6 \times \frac{3.32}{\sqrt{16}}$$

$$= 39.4$$

$$UB = \bar{x} + t^* \frac{s}{\sqrt{n}}$$

$$= 41.52 + 2.6 \times \frac{3.32}{\sqrt{16}}$$

$$= 43.7$$

We are 98% confident that the population mean is between 39.4 and 43.7.

$$CI = (39.4, 43.7)$$

10. Problem:

A null hypothesis claims a population has a mean $\mu = 120$. You decide to run two-tail test on a sample of size $n = 12$ using a significance level $\alpha = 0.1$.

You then collect the sample:

180.4	152.7	128.5	163.7	72
143.7	101.2	144.5	117.8	142.7
167.9	120.3			

- (a) Determine the p -value.
- (b) Do you reject the null hypothesis?

Solution: State the hypotheses.

$$H_0 \text{ claims } \mu = 120$$

$$H_A \text{ claims } \mu \neq 120$$

Find the mean and standard deviation of the sample.

$$\bar{x} = 136.283$$

$$s = 30.379$$


Determine the degrees of freedom.

$$df = 12 - 1 = 11$$

Find the standard error.

$$\sigma_{\bar{x}} = \frac{s}{\sqrt{n}} = \frac{30.379}{\sqrt{12}} = 8.77$$

Make a sketch of the null's sampling distribution.



Find the t score.

$$t = \frac{\bar{x} - \mu_0}{\sigma_{\bar{x}}} = \frac{136.283 - 120}{8.77} = 1.86$$

Find the p -value.

$$p\text{-value} = P(|T| > 1.86)$$

We can't get an exact value with our table, but we can determine an interval that contains the p -value. (Look at row with $df = 11$.)

$$P(|T| > 2.2) = 0.05$$

$$P(|T| > 1.8) = 0.1$$

Basically, because t is between 2.2 and 1.8, we know the p -value is between 0.05 and 0.1.

$$0.05 < p\text{-value} < 0.1$$

Compare the p -value and the significance level ($\alpha = 0.1$).

$$p\text{-value} < \alpha$$

Yes, we reject the null hypothesis.

(a) $0.05 < p\text{-value} < 0.1$

(b) Yes, we reject the null hypothesis.

11. Problem:

A random sample of size 1900 was found to have a sample proportion of 80% (because there were 1520 successes). Determine a 92% confidence interval of the population proportion.

- (a) Find the lower bound of the confidence interval.
- (b) Find the upper bound of the confidence interval.

Solution: Identify the givens.

$$n = 1900$$

$$\hat{p} = 0.8$$

$$\gamma = 0.92$$

Determine z^* such that $P(|Z| < z^*) = 0.92$. It is helpful to know that $\ell = \frac{\gamma+1}{2} = \frac{0.92+1}{2} = 0.96$. (Use the z-table to find z^* .)

$$z^* = 1.75$$

Use the formula (proportion) for the bounds.

$$\begin{aligned} LB &= \hat{p} - z^* \sqrt{\frac{\hat{p}(1 - \hat{p})}{n}} & UB &= \hat{p} + z^* \sqrt{\frac{\hat{p}(1 - \hat{p})}{n}} \\ &= 0.8 - 1.75 \sqrt{\frac{(0.8)(0.2)}{1900}} & &= 0.8 + 1.75 \sqrt{\frac{(0.8)(0.2)}{1900}} \\ &= 0.784 & &= 0.816 \end{aligned}$$

Determine the interval.

$$CI = (0.784, 0.816)$$

We are 92% confident that the true population proportion is between 78.4% and 81.6%.

- (a) The lower bound = 0.784, which can also be expressed as 78.4%.
- (b) The upper bound = 0.816, which can also be expressed as 81.6%.

12. Problem:

A fair coin should have a 50% chance of landing on either side. Someone is mildly suspicious that a coin is unfair.

You are asked to judge the fairness of the coin by flipping it 6000 times and counting how many heads are flipped. You are told to use a significance level of 0.05.

Then, you actually flip the coin 6000 times and get 2934 heads. Should we conclude this coin is unfair?

- (a) What kind of hypothesis test is appropriate?
- (b) State the hypotheses
- (c) Determine the p -value.
- (d) Decide whether we reject or retain the null hypothesis.
- (e) Do you think the coin is unfair?

Solution: We should use a two-tail proportion test.

State the hypotheses.

$$H_0 \text{ claims } p = 0.5$$

$$H_A \text{ claims } p \neq 0.5$$

Determine the standard error.

$$\sigma_{\hat{p}} = \sqrt{\frac{p_0(1-p_0)}{n}} = \sqrt{\frac{0.5(1-0.5)}{6000}} = 0.00645$$


Determine the sample proportion.

$$\hat{p} = 0.489$$

Determine a z score. For simplicity, we ignore the continuity correction.

$$z = \frac{\hat{p} - p_0}{\sigma_{\hat{p}}} = \frac{0.489 - 0.5}{0.00645} = -1.7$$

Make a sketch of the null's sampling distribution. The p -value is a two-tail area.



To determine that two-tail area, we use the z table.

$$\begin{aligned} p\text{-value} &= P(|Z| > 1.7) \\ &= 2 \cdot P(Z < -1.7) \\ &= 0.0892 \end{aligned}$$

Compare p -value to α (which is 0.05).

$$p\text{-value} > \alpha$$

Make the conclusion: we don't reject the null hypothesis.

We conclude the coin could be fair.

- (a) Two-tail proportion test
- (b) Hypotheses: H_0 claims $p = 0.5$ and H_A claims $p \neq 0.5$.
- (c) The p -value is 0.0892
- (d) We don't reject the null hypothesis.
- (e) We conclude the coin could be fair.