

Non-Relativistic Ten-Dimensional Minimal Supergravity

E. A. Bergshoeff^{1a}, *J. Lahnsteiner*^{1b}, *L. Romano*^{1c},
J. Rosseel^{2d} and *C. Şimşek*^{1e}

¹*Van Swinderen Institute, University of Groningen
Nijenborgh 4, 9747 AG Groningen, The Netherlands*

²*Faculty of Physics, University of Vienna,
Boltzmanngasse 5, A-1090, Vienna, Austria*

Abstract

We construct a non-relativistic limit of ten-dimensional minimal supergravity both from the point of view of the action and the equations of motion. This limit can only be realized provided we impose by hand a set of geometric constraints, invariant under all the symmetries of the non-relativistic theory, that define a so-called ‘self-dual’ String Newton-Cartan geometry. The non-relativistic action exhibits three emerging symmetries: one local scale symmetry and two local conformal supersymmetries. Due to these emerging symmetries the Poisson equation for the Newton potential and two partner fermionic equations do not follow from a variation of the non-relativistic action but, instead, are obtained by a supersymmetry variation of the other equations of motion that do follow from a variation of the non-relativistic action. We show that, after taking into account the geometric constraints that follow from the equations of motion, a natural geometry is given by the same dilatation invariant String Newton-Cartan geometry that we found in the bosonic case. We shortly discuss the inclusion of the Yang-Mills sector that would lead to a non-relativistic heterotic supergravity action.

^aEmail: e.a.bergshoeff@rug.nl

^bEmail: j.m.lahnsteiner@outlook.com

^cEmail: lucaromano2607@gmail.com

^dEmail: jan.rosseel@univie.ac.at

^eEmail: c.simsek@rug.nl

Contents

1	Introduction	2
2	$D = 10$, $\mathcal{N} = 1$ Supergravity	5
3	The NR Limit of the Action	9
4	The NR Limit of the Equations of Motion	13
4.1	Equations of motion from the NR action and missing NR equations of motion . . .	13
4.2	Consistency of all NR equations of motion under supersymmetry and Galilean boosts	19
5	Conclusions	23
A	Notation and Conventions	25
A.1	Bosonic Conventions	25
A.2	Spinor and Clifford Algebra Conventions	25
B	Torsional String Newton-Cartan Geometry	27
C	Bosonic Equations of Motion and Killing Spinor Equations	28
D	Non-Relativistic $D = 10$ Super-Yang-Mills	29
E	Closure of the Non-Relativistic Super-Algebra	31
E.1	Boost- $\delta^{(0)}$	31
E.2	Dilatation- $\delta^{(0)}$	32
E.3	S-symmetry- $\delta^{(0)}$	32
E.4	T-symmetry- $\delta^{(0)}$	32
E.5	$\delta^{(0)}$ - $\delta^{(0)}$	33
E.6	Dilatation-S/T Symmetry	34
E.7	Boost-S/T Symmetry	34
E.8	S/T Symmetry-S/T Symmetry	34

1 Introduction

{sec:intro}

Recently, major progress has been made in understanding the formulation of non-relativistic (NR) string theory in a general curved background thereby generalizing the original proposal for NR string theory in a flat background [1, 2] and its early extensions to special curved backgrounds [3]. These new developments have taken place both in terms of a description via a two-dimensional non-linear sigma model as well as from the point of view of a target space action and equations of motion for the background fields. For the closed bosonic string these results have been obtained either by taking a NR limit [4–7] or by applying a null reduction [7–10]. Moreover, the relation between the sigma model beta-functions and the target space equations of motion has been clarified, both for closed and open strings, proving the one-loop quantum consistency of the NR string theory [11–15]. There is also an intriguing relationship with Double Field Theory [16–21]. For other recent work on NR string theory in a curved background, see [22–26].

At first sight, the natural target space geometry of the NR string theory of [1, 2] generalized to arbitrary backgrounds is given by a Newton-Cartan-like geometry with co-dimension two foliation that is characterized by the following ‘zero torsion constraint’ on the longitudinal Vielbein τ_μ^A :

$$D_{[\mu}(\omega)\tau_{\nu]}^A = 0. \quad (1) \quad \{\text{stringconst}$$

Here, the index $A = 0, 1$ refers to the directions longitudinal to the string and the derivative $D_\mu(\omega)$ is covariant with respect to longitudinal Lorentz transformations, using a spin connection ω_μ^{AB} . The constraints (1) are partly conventional defining the longitudinal spin connection as a composite field and partly geometric. To obtain the geometric constraints one should project the

constraints (1) onto those components where the spin connection cancels out. This leads to the following geometric constraints:

$$e_{A'}{}^\mu \tau_{[A}{}^\nu \partial_{[\mu} \tau_{\nu]}{}_{|B]} = 0, \quad \text{and} \quad e_{A'}{}^\mu e_{B'}{}^\nu \partial_{[\mu} \tau_{\nu]}{}^A = 0. \quad (2) \quad \{\{\text{stringconst}\}\}$$

Here A' refers to the directions transverse to the string, $\tau_A{}^\mu$, $e_{A'}{}^\mu$ are (projective) inverses of the longitudinal and transverse Vielbeine $\tau_\mu{}^A$, $e_\mu{}^{A'}$ and (AB) indicates the symmetric part of AB . The geometry defined by the zero torsion constraint (1) is referred to as String Newton-Cartan (SNC) geometry [27].¹ The NR string then couples to the SNC Vielbeine, as well as to a Kalb-Ramond (KR) and dilaton background field.

In our recent paper [6], we studied the target space action and equations of motion of the NS-NS sector of NR string theory, from the viewpoint of taking a NR limit of the relativistic action and equations of motion. The same NR NS-NS action has also been derived from a Double Field Theory point of view in [16]. We showed in particular that in the NR case a natural geometrical constraint, consistent with the target space equations of motion, is not given by the constraints (2) of SNC geometry, but by the weaker dilatation-invariant geometric constraints²

$$e_{A'}{}^\mu \tau_{\{A}{}^\nu \partial_{[\mu} \tau_{\nu]}{}_{|B\}} = 0, \quad \text{and} \quad e_{A'}{}^\mu e_{B'}{}^\nu \partial_{[\mu} \tau_{\nu]}{}^A = 0, \quad (3) \quad \{\{\text{TTSintro}\}\}$$

where $\{AB\}$ indicates the symmetric traceless part of AB . The constraints (3) characterize what we called a ‘Dilatation invariant String Newton-Cartan’ (DSNC) geometry in [6]. We stress that these constraints are not imposed by hand but follow as natural solutions of the target space equations of motion. The geometry before imposing any constraints is referred to as Torsional String Newton-Cartan (TSNC) geometry [6, 7].

The NR limit of the relativistic equations of motion for the (metric, Kalb-Ramond, dilaton) background fields can be taken such that it preserves the number of independent equations of motion. However, not all of the resulting NR equations follow from a variation of the NR action that is obtained by taking the NR limit of the relativistic one. A distinguishing feature of this NR NS-NS target space action is that it is invariant under an emerging local scale symmetry that is absent in the relativistic case. Due to this emerging symmetry there is one equation, the Poisson equation for the Newton potential, that does not follow from the NR target space action. Instead, this equation constitutes, together with all the other equations that do follow from the NR target space action, a so-called reducible but indecomposable representation of the NR symmetries. For the Poisson equation this means that by varying it under Galilean boosts one can generate the full set of equations but none of these other equations transforms back under Galilean boosts to the Poisson equation. A similar story applies to the beta-functions. The NR string sigma model is also invariant under an emerging dilatation symmetry and consequently the number of beta-functions that one can calculate is one less than the number of relativistic equations of motion. Setting these beta-functions to zero one finds that a certain nonlinear equation that has a dilatation weight opposite to that of the Poisson equation is missing. For an interpretation of this missing nonlinear equation, see the recent paper [31]. Schematically, the situation can be summarized as follows:

$$\text{NR e.o.m.} \quad \rightarrow \quad \text{common equations} + \text{Poisson} + \text{Non-linear}, \quad (4)$$

$$\text{NR NS-NS action} \quad \rightarrow \quad \text{common equations} + \text{Non-linear}, \quad (5)$$

$$\text{NR } \beta\text{-functions} \quad \rightarrow \quad \text{common equations} + \text{Poisson}. \quad (6)$$

We note that if, by taking the NR limit of the equations of motion, one would loose a degree of freedom all the while the number of independent equations of motion, whose linearization is non-trivial, would remain the same, the theory would become over-determined. This is avoided by the fact that one of the equations becomes inherently non-linear. The fact that the NR NS-NS action does not lead to the Poisson equation for the Newton potential is consistent with the fact that no action principle is known for NC gravity based upon the Bargmann algebra.³

¹For earlier work on SNC geometry, see [3, 28, 29].

²These constraints are obtained by first writing the constraints (1) in a dilatation invariant way using a covariant derivative $D_\mu(\omega, b)$ where b_μ is the dilatation gauge field and next projecting to those components where both gauge fields cancel out. The second constraint in (3) is sufficient to define a globally well-defined co-dimension two foliation, called integrable distribution [30].

³For suggestions of such an action based on a larger algebra, see [32]

So far, most calculations have been performed for the bosonic NR string only.⁴ This work is a companion to our previous paper where we enlarge our investigations of the NR NS-NS gravity background to the case of a minimal supergravity background.⁵ To be specific, we will present the NR limit of the ten-dimensional $\mathcal{N} = 1$ supergravity action and equations of motion defining the dynamics of the background fields. This sector is common to all superstring theories. We have a heterotic superstring in mind but we will postpone a discussion of the Yang-Mills sector to the conclusions. To obtain the results of this paper, we will follow the same strategy as used in [6] but there are notable new features in the supersymmetric case. One complication is that, unlike in the bosonic case, there is no direct connection between a two-dimensional sigma model description and the NR target space effective action. A Green-Schwarz sigma model formulation for the NR superstring has been given for a flat background [33] but not for a NR minimal supergravity background. Like in the relativistic case, this will probably require a superspace formulation. Alternatively, starting from a NR sigma model with (1,0) world-sheet supersymmetry⁶, the target space supersymmetry of the background fields is not manifest. In both cases we cannot use the sigma model description to read off the emergent target space fermionic Stueckelberg symmetries that we do expect to team up with the emergent local scale symmetry that we found in the bosonic case.

Another complication, not encountered in the bosonic case, is that taking the naive NR limit of the supersymmetry rules leads to divergent terms in these transformation rules. Concerning the action, we find that, like in the bosonic case, a NR limit of the $\mathcal{N} = 1$ supergravity action can be defined due to a miraculous cancellation of divergent terms when taking this limit. By performing a careful analysis of the NR limit, we will show in this paper that the dangerous divergent terms in the supersymmetry rules are controlled by (1) making use of the fact that the NR action is invariant under two emergent local fermionic Stueckelberg symmetries arising as partners of the emergent local scale symmetry that we already found in the bosonic case and (2) imposing by hand the following constraints on the geometry:

$$e_{A'}{}^\mu \tau_+{}^\nu \partial_{[\mu} \tau_{\nu]}^- = 0, \quad \text{and} \quad e_{A'}{}^\mu e_{B'}{}^\nu \partial_{[\mu} \tau_{\nu]}^- = 0. \quad (7) \quad \{\text{selfdualint}$$

These constraints are needed to show the consistency of the theory. Here \pm refer to (anti-)selfdual projections in the two longitudinal directions. We will call the two emerging fermionic symmetries S and T supersymmetry where the S supersymmetry is of a type that is also encountered in conformal supergravity. We find that the constraints (7) are invariant under all the symmetries of the NR theory, including supersymmetry, and therefore do not lead to additional constraints. They can be substituted into the equations of motion but not into the NR action. In that sense the NR minimal supergravity action is a so-called pseudo action. The constraints (7) constitute half of the constraints (3) defining a DSNC geometry and define what we will call a ‘self-dual’ DSNC geometry. We note that there could be more geometric constraints in general but these follow from the equations of motion for the background fields that we will derive in this paper. The final quantum-consistent geometry can only be determined after the constraints that follow from these equations of motion have been taken into account.

Due to the emergent bosonic and fermionic local symmetries we find that the NR action does not give rise to the full set of equations of motion. There is a Poisson equation for the Newton potential and there are two additional fermionic equations that do not follow from the variation of the NR action. However, unlike in the bosonic case, the NR action knows indirectly about these three missing equations in the sense that they can be obtained by varying the other equations of motion that follow from the variation of the NR action under supersymmetry.

The organization of this paper is as follows. In section 2 we give a brief review of the relativistic $\mathcal{N} = 1$ supergravity theory together with the transformations rules of all fields in a new basis of the fields that contains powers of c for finite c , i.e., before taking the actual NR limit. In the next section we discuss in detail the NR limit of the relativistic supergravity action ending up with a NR action that has emergent dilatations plus an emerging S and T supersymmetry. In section 4 we take the NR limit of the equations of motion and show that we obtain the same

⁴For earlier work on NR strings and supersymmetry, see [3, 33].

⁵The NR supergravity theory is minimal in the sense that, although it contains *two* independent supersymmetries, both are needed to obtain the Hamiltonian as the result of an anti-commutator of two supercharges.

⁶For the recent construction of a NR sigma model with (1,1) worldsheet supersymmetry, see [34].

equations of motion that follow from varying the NR action derived in the previous section plus three more equations: the Poisson equation for the Newton potential together with two fermionic equations. Furthermore, we show how these three missing equations of motion are connected to the ones that do follow from the variation of the NR action via supersymmetry. In the final section we discuss our results. In particular we mention a few subtleties when including the Yang-Mills sector of a heterotic supergravity theory. There are 4 Appendices. Our notations and conventions particular to the supersymmetric case are given in Appendix A. In Appendix B we collect a few useful formulae describing Torsional String Newton-Cartan Geometry. To make this paper more user-friendly for those who wish to investigate compactifications or solutions of the NR superstring we have summarized in Appendix C the bosonic truncation of all equations of motion together with all the bosonic terms in the fermionic supersymmetry rules. Finally, in the last Appendix we show how the NR limit can be defined for the special case of a supersymmetric Yang-Mills theory in a flat background.

2 $D = 10, \mathcal{N} = 1$ Supergravity

In this section, we will briefly review ten-dimensional $\mathcal{N} = 1$ supergravity [35, 36], i.e., the common part of the effective theories for all superstrings. We will summarize its fields and their transformation rules, as well as its action. In order to define the NR limit, one performs an invertible field redefinition that involves a (dimensionless) parameter c , such that the NR limit corresponds to sending $c \rightarrow \infty$. To facilitate taking this limit in the next sections, we will here also give the field redefinition that is involved and apply it to the transformation rules of $\mathcal{N} = 1$ supergravity. The index, spinor and Clifford algebra conventions that we use throughout this paper, are collected in Appendix A.

The bosonic field content of ten-dimensional $\mathcal{N} = 1$ supergravity consists of the Zehnbein $E_\mu^{\hat{A}}$, the Kalb-Ramond (KR) two-form field $B_{\mu\nu}$ and the dilaton Φ . The fermionic fields are given by the gravitino Ψ_μ , and the dilatino λ . Here, Ψ_μ is a left-handed Majorana-Weyl spinor, while λ is a right-handed one. The action of $\mathcal{N} = 1$ supergravity is then given by

$$S = \frac{1}{2\kappa^2} \int d^{10}x E e^{-2\Phi} \left\{ \mathcal{R} + 4 \partial_\mu \Phi \partial^\mu \Phi - \frac{1}{12} \mathcal{H}_{\mu\nu\rho} \mathcal{H}^{\mu\nu\rho} - 2 \bar{\Psi}_\mu \Gamma^{\mu\nu\rho} D_\nu \Psi_\rho - 4 \bar{\lambda} \Gamma^{\mu\nu} D_\mu \Psi_\nu \right. \\ \left. + 2 \bar{\lambda} \not{D} \lambda + \frac{1}{24} \mathcal{H}^{\rho\sigma\tau} \left(2 \bar{\Psi}_\mu \Gamma^{[\mu} \Gamma_{\rho\sigma\tau} \Gamma^{\nu]} \Psi_\nu - 4 \bar{\Psi}_\mu \Gamma^\mu_{\rho\sigma\tau} \lambda - 2 \bar{\lambda} \Gamma_{\rho\sigma\tau} \lambda \right) \right. \\ \left. - 4 \bar{\Psi}_\mu \not{\partial} \Phi \Gamma^\mu \lambda - 4 \bar{\Psi}_\mu \Gamma^\mu \Psi_\nu \partial^\nu \Phi \text{ (+ quartic fermion terms)} \right\}, \quad (8)$$

where κ denotes the gravitational coupling constant and $E = \det(E_\mu^{\hat{A}})$. The Ricci scalar \mathcal{R} is constructed from the Levi-Civita spin connection $\Omega_\mu^{\hat{A}\hat{B}}$ and

$$\mathcal{H}_{\mu\nu\rho} = 3 \partial_{[\mu} B_{\nu\rho]}, \quad (9)$$

is the field strength of the KR field. We moreover defined the (anti-symmetrized when necessary) covariant derivatives of Ψ_μ and λ by

$$D_{[\mu} \Psi_{\nu]} = \partial_{[\mu} \Psi_{\nu]} - \frac{1}{4} \Omega_{[\mu}^{\hat{A}\hat{B}} \Gamma_{\hat{A}\hat{B}} \Psi_{\nu]}, \quad D_\mu \lambda = \partial_\mu \lambda - \frac{1}{4} \Omega_\mu^{\hat{A}\hat{B}} \Gamma_{\hat{A}\hat{B}} \lambda. \quad (10)$$

Note that $\Omega_\mu^{\hat{A}\hat{B}}$, \mathcal{R} and $\mathcal{H}_{\mu\nu\rho}$ in the action (8) do not contain any fermionic contributions (such as supercovariantizations). The first three terms of (8) are thus purely bosonic and only the remaining terms contain fermions. We have not explicitly written the quartic fermion terms that are present in (8). In this paper, we will consistently truncate quartic fermion terms in actions and we will only give the terms in the supersymmetry transformation rules that are consistent with this truncation.

The fields of ten-dimensional $\mathcal{N} = 1$ supergravity transform as follows under local Lorentz transformations with parameter $\Lambda^{\hat{A}\hat{B}}$, a one-form symmetry of the KR field with parameter Θ_μ and supersymmetry with a left-handed Majorana-Weyl spinor parameter ε :

$$\delta E_\mu^{\hat{A}} = \Lambda^{\hat{A}}_{\hat{B}} E_\mu^{\hat{B}} + \bar{\varepsilon} \Gamma^{\hat{A}} \Psi_\mu, \quad \delta B_{\mu\nu} = 2 \partial_{[\mu} \Theta_{\nu]} + 2 \bar{\varepsilon} \Gamma_{[\mu} \Psi_{\nu]}, \quad \delta \Phi = \frac{1}{2} \bar{\varepsilon} \lambda, \quad (11a)$$

$$\delta\Psi_\mu = \frac{1}{4}\Lambda^{\hat{A}\hat{B}}\Gamma_{\hat{A}\hat{B}}\Psi_\mu + D_\mu(\Omega^{(+)})\varepsilon \text{ (+ terms quadratic in } \Psi_\mu \text{ and } \lambda), \quad (11b) \quad \{\text{eq:trafosGr}\}$$

$$\delta\lambda = \frac{1}{4}\Lambda^{\hat{A}\hat{B}}\Gamma_{\hat{A}\hat{B}}\lambda + \Gamma^\mu\varepsilon\partial_\mu\Phi - \frac{1}{12}\Gamma^{\hat{A}\hat{B}\hat{C}}\varepsilon\mathcal{H}_{\hat{A}\hat{B}\hat{C}} \text{ (+ terms quadratic in } \Psi_\mu \text{ and } \lambda), \quad (11c) \quad \{\text{eq:trafosDi}\}$$

where we have defined the following torsionful covariant derivative of ε

$$D_\mu(\Omega^{(+)})\varepsilon = \partial_\mu\varepsilon - \frac{1}{4}\Omega_\mu^{(+)\hat{A}\hat{B}}\Gamma_{\hat{A}\hat{B}}\varepsilon, \quad \text{with} \quad \Omega_\mu^{(+)\hat{A}\hat{B}} = \Omega_\mu^{\hat{A}\hat{B}} + \frac{1}{2}\mathcal{H}_\mu^{\hat{A}\hat{B}}. \quad (12) \quad \{\text{eq:D0megapm}\}$$

In order to take the NR limit in the following sections, we introduce a (dimensionless) parameter c and perform the following field redefinition

$$\begin{aligned} \tau_\mu^A &= c^{-1}E_\mu^A, & e_\mu^{A'} &= E_\mu^{A'}, & b_{\mu\nu} &= B_{\mu\nu} + \epsilon_{AB}E_\mu^AE_\nu^B, & \phi &= \Phi - \log c, \\ \psi_{\mu\pm} &= c^{\mp 1/2}\Pi_\pm\Psi_\mu, & \lambda_\pm &= c^{\mp 1/2}\Pi_\pm\lambda, \end{aligned} \quad (13) \quad \{\text{eq:rescale}\}$$

where we have split the Lorentz index \hat{A} into a longitudinal index $A = 0, 1$ and a transversal index $A' = 2, \dots, 9$. Note that the redefinition of the spinor fields involves the ‘worldsheet chirality’ projection operators Π_\pm , that are defined in (101) [33]. We refer to Appendix A.2 for various properties that are obeyed by worldsheet chirality projected spinors and that are used throughout this paper. For the bosonic fields, the above redefinition coincides with the one used in [6] to derive the NR limit of NS-NS gravity.

As will be seen in the next sections, the fields τ_μ^A , $e_\mu^{A'}$, $b_{\mu\nu}$, ϕ , $\psi_{\mu\pm}$ and λ_\pm correspond, after taking the limit $c \rightarrow \infty$, to the fields of NR ten-dimensional minimal supergravity. As explained in [6], in order to calculate the transformation rules of these fields in the NR theory, it is important that the field redefinition (13) is invertible. This is the case and the inverse of (13) is given by

$$\begin{aligned} E_\mu^A &= c\tau_\mu^A, & E_\mu^{A'} &= e_\mu^{A'}, & B_{\mu\nu} &= -c^2\epsilon_{AB}\tau_\mu^A\tau_\nu^B + b_{\mu\nu}, & \Phi &= \phi + \log c, \\ \Psi_\mu &= c^{1/2}\psi_{\mu+} + c^{-1/2}\psi_{\mu-}, & \lambda &= c^{1/2}\lambda_+ + c^{-1/2}\lambda_-, \end{aligned} \quad (14) \quad \{\text{eq:rescalei}\}$$

where it is understood that $\psi_{\mu\pm}$ and λ_\pm are worldsheet chirality projected spinors (i.e., obey $\psi_{\mu\pm} = \Pi_\pm\psi_{\mu\pm}$ and $\lambda_\pm = \Pi_\pm\lambda_\pm$). It is also useful to introduce objects τ_A^μ and $e_{A'}^\mu$ as the following (invertible) redefinitions of components of the inverse Vielbein $E_{\hat{A}}^\mu$:

$$\tau_A^\mu = cE_A^\mu, \quad e_{A'}^\mu = E_{A'}^\mu. \quad (15) \quad \{\text{eq:invEresc}\}$$

The τ_μ^A , $e_\mu^{A'}$, τ_A^μ , $e_{A'}^\mu$ then satisfy the following ‘projective invertibility’ relations:

$$\begin{aligned} \tau_B^\mu\tau_\mu^A &= \delta_B^A, & e_{B'}^\mu e_\mu^{A'} &= \delta_{B'}^{A'}, & \tau_A^\mu e_\mu^{A'} &= 0, \\ e_{A'}^\mu\tau_\mu^A &= 0, & \tau_\mu^A\tau_A^\nu + e_\mu^{A'}e_{A'}^\nu &= \delta_\mu^\nu. \end{aligned} \quad (16) \quad \{\text{eq:projinv}\}$$

To set the stage for our derivation of the NR limit of the action and equations of motion of ten-dimensional $\mathcal{N} = 1$ supergravity, we will end this section by applying the above field redefinition (13), (14) to the transformation rules (11). To do this, we introduce an analogous invertible redefinition of the parameters $\Lambda^{\hat{A}\hat{B}} = (\Lambda^{AB}, \Lambda^{AA'}, \Lambda^{A'B'})$, Θ_μ and ε of local Lorentz transformations, the KR one-form symmetry and supersymmetry:

$$\begin{aligned} \lambda^{A'B'} &= \Lambda^{A'B'}, & \lambda^{AA'} &= c\Lambda^{AA'}, & \lambda_M\epsilon^{AB} &= \Lambda^{AB}, & \theta_\mu &= \Theta_\mu \\ \epsilon_\pm &= c^{\mp 1/2}\Pi_\pm\varepsilon \Leftrightarrow \varepsilon = c^{1/2}\epsilon_+ + c^{-1/2}\epsilon_- \text{ (with } \Pi_\pm\epsilon_\pm = \epsilon_\pm). \end{aligned} \quad (17) \quad \{\text{eq:parresc}\}$$

After taking the $c \rightarrow \infty$ limit, λ_M will correspond to the parameter of longitudinal Lorentz transformations, $\lambda^{AA'}$ to the parameter of Galilean boosts and $\lambda^{A'B'}$ to the parameter of transversal rotations, while the parameters ϵ_\pm will be those of non-relativistic supersymmetry.

Using the redefinitions (13), (14) and (17), one can easily find how τ_μ^A , $e_\mu^{A'}$, $b_{\mu\nu}$, ϕ , $\psi_{\mu\pm}$ and λ_\pm transform under the symmetries with parameters λ_M , $\lambda^{AA'}$, $\lambda^{A'B'}$, θ and ϵ_\pm . Considering first the transformation rules under the bosonic symmetries, one finds

$$\delta\tau_\mu^A = \lambda_M\epsilon^A{}_B\tau_\mu^B + \frac{1}{c^2}\lambda^A{}_{A'}e_\mu^{A'}, \quad \delta e_\mu^{A'} = \lambda^{A'}{}_{B'}e_\mu^{B'} - \lambda_A{}^{A'}\tau_\mu^A, \quad (18a) \quad \{\text{eq:LorsymmB0}\}$$

$$\delta b_{\mu\nu} = 2 \partial_{[\mu} \theta_{\nu]} - 2 \epsilon_{AB} \lambda^A{}_{A'} \tau_{[\mu}{}^B{}_{\nu]} e_{\nu]}{}^{A'}, \quad \delta \phi = 0. \quad (18b)$$

for the bosonic fields and

$$\delta \psi_{\mu+} = \frac{1}{4} (\lambda^{A'B'} \Gamma_{A'B'} - 2 \lambda_M) \psi_{\mu+} + \frac{1}{2 c^2} \lambda^{AA'} \Gamma_{AA'} \psi_{\mu-}, \quad (19a)$$

$$\delta \psi_{\mu-} = \frac{1}{4} (\lambda^{A'B'} \Gamma_{A'B'} + 2 \lambda_M) \psi_{\mu-} + \frac{1}{2} \lambda^{AA'} \Gamma_{AA'} \psi_{\mu+}, \quad (19b)$$

$$\delta \lambda_+ = \frac{1}{4} (\lambda^{A'B'} \Gamma_{A'B'} - 2 \lambda_M) \lambda_+ + \frac{1}{2 c^2} \lambda^{AA'} \Gamma_{AA'} \lambda_-, \quad (19c)$$

$$\delta \lambda_- = \frac{1}{4} (\lambda^{A'B'} \Gamma_{A'B'} + 2 \lambda_M) \lambda_- + \frac{1}{2} \lambda^{AA'} \Gamma_{AA'} \lambda_+, \quad (19d)$$

for the fermionic fields. Note that for both sets of transformation rules (18), (19), the limit $c \rightarrow \infty$ is well-defined.

One can similarly find the transformation rules under supersymmetry (with parameters ϵ_{\pm}). For the bosonic fields $\tau_{\mu}{}^A$, $e_{\mu}{}^{A'}$, $b_{\mu\nu}$, ϕ , one finds, upon using that certain bilinears with spinors of definite worldsheet chirality are identically zero (see e.g. (105)), that

$$\delta \tau_{\mu}{}^A = \bar{\epsilon}_+ \Gamma^A \psi_{\mu+} + \frac{1}{c^2} \bar{\epsilon}_- \Gamma^A \psi_{\mu-}, \quad (20a)$$

$$\delta e_{\mu}{}^{A'} = \bar{\epsilon}_+ \Gamma^{A'} \psi_{\mu-} + \bar{\epsilon}_- \Gamma^{A'} \psi_{\mu+}, \quad (20b)$$

$$\delta \phi = \frac{1}{2} (\bar{\epsilon}_+ \lambda_- + \bar{\epsilon}_- \lambda_+), \quad (20c)$$

$$\delta b_{\mu\nu} = 4 \tau_{[\mu}{}^A \bar{\epsilon}_- \Gamma_A \psi_{\nu]-} + 2 \left(e_{[\mu}{}^{A'} \bar{\epsilon}_+ \Gamma_{A'} \psi_{\nu]-} + E_{[\mu}{}^{A'} \bar{\epsilon}_- \Gamma_{A'} \psi_{\nu]+} \right). \quad (20d)$$

As for (18) and (19), the $c \rightarrow \infty$ limit of these transformations is regular. The derivation of the ϵ_{\pm} supersymmetry transformation rules of $\psi_{\mu\pm}$ and λ_{\pm} is straightforward, but leads to more lengthy expressions that involve powers of c^2 , c^0 and c^{-2} . We collect terms with like powers of c as follows:

$$\delta \psi_{\mu\pm} = c^2 \delta^{(2)} \psi_{\mu\pm} + c^0 \delta^{(0)} \psi_{\mu\pm} + c^{-2} \delta^{(-2)} \psi_{\mu\pm}, \quad (21a)$$

$$\delta \lambda_{\pm} = c^2 \delta^{(2)} \lambda_{\pm} + c^0 \delta^{(0)} \lambda_{\pm} + c^{-2} \delta^{(-2)} \lambda_{\pm}. \quad (21b)$$

Explicitly, the terms that appear at order c^2 are given by

$$\delta^{(2)} \psi_{\mu+} = \frac{1}{2} \tau_{\mu}{}^+ \tau^{A'B'} \Gamma_{A'B'} \epsilon_+, \quad (22a)$$

$$\delta^{(2)} \psi_{\mu-} = \frac{1}{2} \tau_{\mu}{}^+ (\tau^{A'B'} \Gamma_{A'B'} \epsilon_- - \tau^{A'-} \Gamma_{-A'} \epsilon_+), \quad (22b)$$

$$\delta^{(2)} \lambda_+ = 0, \quad (22c)$$

$$\delta^{(2)} \lambda_- = -\frac{1}{2} \tau^{A'B'} \Gamma_{A'B'} \epsilon_+, \quad (22d)$$

where $\tau_{\mu\nu}{}^A = \partial_{[\mu} \tau_{\nu]}{}^A$ and we refer to Appendix A.1 for details on how curved μ, ν indices are converted into flat longitudinal and transversal ones and on how flat light-cone indices are used to denote longitudinal directions (as an alternative to $A, B = 0, 1$).

The terms in (21) at order c^0 can be written in terms of composite fields b_{μ} , ω_{μ} , $\omega_{\mu}{}^{AA'}$, $\omega_{\mu}{}^{A'B'}$ that depend on the bosonic fields $\tau_{\mu}{}^A$, $e_{\mu}{}^{A'}$, $b_{\mu\nu}$ and ϕ . Their explicit expressions can be found in Appendix B. These composite fields correspond to the dependent dilatation and spin connections of the Torsional SNC (TSNC) geometry, that was introduced in [6]. In particular, ω_{μ} , $\omega_{\mu}{}^{AA'}$ and $\omega_{\mu}{}^{A'B'}$ will correspond to spin connections for longitudinal $\text{SO}(1,1)$ Lorentz transformations, Galilean boosts and $\text{SO}(8)$ transversal rotations, while b_{μ} will act as a gauge field for an emerging local dilatation symmetry. In terms of these dependent gauge fields, we then find that

$$\delta^{(0)} \psi_{\mu+} = \delta_+ \psi_{\mu+} + \delta_- \psi_{\mu-} + \frac{1}{2} \tau_{\mu}{}^+ \Gamma_+ \eta_-, \quad (23a)$$

$$\delta^{(0)} \psi_{\mu-} = \delta_+ \psi_{\mu-} + \delta_- \psi_{\mu-} + \tau_{\mu}{}^+ \rho_-, \quad (23b)$$

$$\delta^{(0)}\lambda_+ = \delta_+\lambda_+ + \delta_-\lambda_+, \quad (23c)$$

$$\delta^{(0)}\lambda_- = \delta_+\lambda_- + \delta_-\lambda_- + \eta_-, \quad (23d)$$

where

$$\eta_- = \left(\partial_+\phi \Gamma^+ - \frac{1}{4} h^{-A'B'} \Gamma_{-A'B'} \right) \epsilon_+ + 2b^{A'} \Gamma_{A'} \epsilon_- \quad (+ \text{ terms quadratic in } \psi_{\mu\pm}, \lambda_{\pm}), \quad (24a)$$

$$\rho_- = \left(-2\partial_+\phi + \frac{1}{4} h^{-A'B'} \Gamma_{A'B'} \right) \epsilon_- + \frac{1}{2} W_+^{-A'} \Gamma_{-A'} \epsilon_+ \quad (+ \text{ terms quadratic in } \psi_{\mu\pm}, \lambda_{\pm}). \quad (24b)$$

and

$$\delta_+\psi_{\mu+} = \mathcal{D}_\mu \epsilon_+ - \frac{1}{8} e_\mu^{C'} h^{C'A'B'} \Gamma_{A'B'} \epsilon_+ \quad (+ \text{ terms quadratic in } \psi_{\mu\pm}, \lambda_{\pm}), \quad (25a)$$

$$\delta_-\psi_{\mu+} = (e_\mu^{B'} \tau^{B'A'+} + \tau_\mu^{-A'+}) \Gamma_{A'+} \epsilon_- \quad (+ \text{ terms quadratic in } \psi_{\mu\pm}, \lambda_{\pm}), \quad (25b)$$

$$\delta_+\psi_{\mu-} = -\frac{1}{2} \omega_\mu^{-A'} \Gamma_{-A'} \epsilon_+ \quad (+ \text{ terms quadratic in } \psi_{\mu\pm}, \lambda_{\pm}), \quad (25c)$$

$$\delta_-\psi_{\mu-} = \mathcal{D}_\mu \epsilon_- - \frac{1}{8} e_\mu^{C'} h^{C'A'B'} \Gamma_{A'B'} \epsilon_+ \quad (+ \text{ terms quadratic in } \psi_{\mu\pm}, \lambda_{\pm}), \quad (25d)$$

$$\delta_+\lambda_+ = (\partial_{A'}\phi \Gamma^{A'} - b_{A'} \Gamma^{A'} - \frac{1}{12} h^{A'B'C'} \Gamma_{A'B'C'}) \epsilon_+ \quad (+ \text{ terms quadratic in } \psi_{\mu\pm}, \lambda_{\pm}), \quad (25e)$$

$$\delta_-\lambda_+ = \frac{1}{2} \tau^{A'B'+} \Gamma_{A'B'+} \epsilon_- \quad (+ \text{ terms quadratic in } \psi_{\mu\pm}, \lambda_{\pm}), \quad (25f)$$

$$\delta_+\lambda_- = 0 \quad (+ \text{ terms quadratic in } \psi_{\mu\pm}, \lambda_{\pm}), \quad (25g)$$

$$\delta_-\lambda_- = (\partial_{A'}\phi \Gamma^{A'} - b_{A'} \Gamma^{A'} - \frac{1}{12} h^{A'B'C'} \Gamma_{A'B'C'}) \epsilon_- \quad (+ \text{ terms quadratic in } \psi_{\mu\pm}, \lambda_{\pm}), \quad (25h)$$

where $\mathcal{D}_\mu \epsilon_\pm$ is given by

$$\mathcal{D}_\mu \epsilon_\pm = \left(\partial_\mu - \frac{1}{4} \omega_\mu^{A'B'} \Gamma_{A'B'} \pm \frac{1}{2} \omega_\mu \mp \frac{1}{2} b_\mu \right) \epsilon_\pm, \quad (26)$$

and $W_+^{-A'}$ in (24) refers to components of the spin connections that are not fully determined in TSNC geometry, but that do not play a role in the rest of this paper (see also Appendix B and [6]). Note that the redefined supersymmetry transformation rules (21) also contain non-trivial terms at order c^{-2} . We will not give the explicit expressions for $\delta^{(-2)}\psi_{\mu\pm}$, $\delta^{(-2)}\lambda_\pm$ here, as we will not need them in what follows.

Let us finish this section, by commenting on the appearance of terms of order c^2 in the supersymmetry transformation rules of the fermionic fields $\psi_{\mu\pm}$ and λ_\pm . Since we wish to identify τ_μ^A , $e_\mu^{A'}$, $b_{\mu\nu}$, ϕ , $\psi_{\mu\pm}$ and λ_\pm as fields in the NR theory that is obtained after taking $c \rightarrow \infty$, one would hope that the redefinitions (13), (14) and (17) lead to transformation rules for these fields that take the form of an expansion in powers of c^{-2} that starts at order c^0 . That way, the $c \rightarrow \infty$ limit of these transformation rules is well-defined and can be identified with the transformation rules of the NR theory. Clearly, the terms of order c^2 in the supersymmetry transformation rules of $\psi_{\mu\pm}$ and λ_\pm are potentially problematic in this regard. In order to explain how to deal with these divergent terms in the next section, let us make the following useful observations here. One can isolate the divergent terms of order c^2 in the supersymmetry transformation rules of $\psi_{\mu\pm}$, λ_\pm by performing the following field redefinition:

$$\tilde{\psi}_{\mu+} \equiv \psi_{\mu+} - \frac{1}{2} \tau_\mu^+ \Gamma_+ \lambda_-, \quad \tilde{\psi}_{\mu-} \equiv \psi_{\mu-} - \tau_\mu^+ \tilde{\psi}_- \quad \text{with} \quad \tilde{\psi}_- \equiv \tau_+^\mu \psi_{\mu-}. \quad (27) \quad \{\text{eq:NRfermio}\}$$

Using (22), one can then easily see that the parts of the supersymmetry transformation rules of $\tilde{\psi}_{\mu\pm}$, λ_+ , $\tilde{\psi}_-$ and λ_- at order c^2 are given by:

$$\delta^{(2)}\tilde{\psi}_{\mu+} = 0, \quad \delta^{(2)}\tilde{\psi}_{\mu-} = 0, \quad \delta^{(2)}\lambda_+ = 0,$$

$$\begin{aligned}\delta^{(2)}\tilde{\psi}_- &= \frac{1}{2}(\tau^{A'B'}\Gamma_{A'B'}\epsilon_- - \tau^{A'--}\Gamma_{-A'}\epsilon_+), \\ \delta^{(2)}\lambda_- &= -\frac{1}{2}\tau^{A'B'}\Gamma_{A'B'}\epsilon_+.\end{aligned}\tag{28} \quad \{\text{eq:susydiv}\}$$

The supersymmetry rules of $\tilde{\psi}_{\mu\pm}$ and λ_{\pm} thus do not contain any ‘divergent’ terms at order c^2 and their $c \rightarrow \infty$ limit is well-defined. Note furthermore that the parts of the supersymmetry transformations of $\tilde{\psi}_{\mu\pm}$ at order c^0 then also no longer depend on the quantities η_- , ρ_- , defined in (24). We will see the significance of these observations in what follows.

3 The NR Limit of the Action

\{\text{sec:limitact}\}

In the previous section, we reviewed the action and transformation rules of relativistic $D = 10$, $\mathcal{N} = 1$ supergravity, introduced a field redefinition that involves a parameter c and expressed all transformation rules for the redefined fields as expansions in powers of c^{-2} . Starting from this section, we wish to discuss the NR limit $c \rightarrow \infty$, that should lead to NR minimal ten-dimensional supergravity, similar to how NR NS-NS gravity was obtained in [6], by performing the bosonic part of the field redefinition (13) and taking the $c \rightarrow \infty$ limit. In this section, we will apply this limit to the action of ten-dimensional $\mathcal{N} = 1$ supergravity, while the limit of its equations of motion will be discussed in the next section.

Ordinarily, the NR limit of quantities (such as an action or equations of motion) is performed by applying a c -dependent field redefinition to the quantities under consideration, expanding the result in powers of c^{-2} and retaining only the terms at leading order. In case the transformation rules of the redefined fields assume the form of expansions in powers of c^{-2} that start at order c^0 , this procedure guarantees that one ends up with quantities that are invariant or covariant under the NR transformation rules that are given by the $c \rightarrow \infty$ limit of those of the redefined fields.

We wish to apply a similar limit procedure to the action (8) of $D = 10$, $\mathcal{N} = 1$ supergravity. In particular, we still wish to define the NR limit of (8) as the leading order term in the c^{-2} -expansion of (8), after performing the field redefinition (13). Similarly, we still want to identify the transformation rules of the NR theory as the part at order c^0 in the c^{-2} -expansions of the relativistic transformation rules (18), (19), (20) and (21) for the redefined fields. Note however that presently, the c^{-2} -expansion of some of the supersymmetry transformation rules of the redefined fermionic fields starts at order c^2 , instead of at order c^0 . As a consequence, the interpretation of the c^0 part of these transformation rules as NR ones is no longer straightforward. Moreover, invariance or covariance under these NR transformation rules of the leading order of an expansion of a quantity in powers of c^{-2} is also no longer guaranteed. Remarkably, even though some transformation rules diverge in the $c \rightarrow \infty$ limit, it turns out that the $c \rightarrow \infty$ limit can be taken in a smooth way, upon imposition of a constraint. Importantly, after taking the limit, the resulting NR action exhibits invariance under three emerging symmetries: one dilatation symmetry and two fermionic shift symmetries. The emerging dilatation symmetry was already encountered when taking the NR limit of NS-NS gravity [6]. In this paper, we find that it extends to a symmetry of the NR limit of the action of $D = 10$, $\mathcal{N} = 1$ supergravity and that it is accompanied by two fermionic symmetries as supersymmetric counterparts.

The emergence of the two fermionic shift symmetries in the NR limit of (8) can be understood on general grounds. In order to see this, let us first apply the redefinition (14) to the action (8) and expand the result in powers of c^{-2} . This gives a sum of three terms, at orders c^0 , c^{-2} and c^{-4} respectively:

$$S = S^{(0)} + c^{-2}S^{(-2)} + c^{-4}S^{(-4)},\tag{29} \quad \{\text{eq:actionex}\}$$

where each of the $S^{(i)}$ now depends on the fields τ_{μ}^A , $e_{\mu}^{A'}$, $b_{\mu\nu}$, ϕ , $\psi_{\mu\pm}$ and λ_{\pm} . It is important to note that it is non-trivial that the expansion (29) of S starts at order c^0 . Indeed, examining all terms of (8) separately, one finds that some of them can contribute terms at order c^2 in the expansion (29), so that there can in principle be a $c^2S^{(2)}$ term on the right-hand-side of (29). It turns out however that all such contributions cancel identically. For the bosonic part of the action (8), this relies on an order c^2 contribution from the Ricci scalar cancelling against a similar

contribution from the kinetic term of the KR field, as was explained in [6]. One can check that this cancellation of order c^2 terms extends to the full $D = 10$, $\mathcal{N} = 1$ supergravity action (8), so that

$$S^{(2)} = 0 \quad (30) \quad \{\text{eq:S2zero}\}$$

identically.

Note that the $c \rightarrow \infty$ limit of S is then well-defined and gives $S^{(0)}$, which we identify as the action that results from taking the NR limit. Let us now examine how $S^{(0)}$ transforms under the NR symmetry transformation rules, that correspond to the parts at order c^0 of the relativistic transformation rules (18), (19), (20) and (21), after performing the redefinition (13), (14). In order to do this, we will not yet take the $c \rightarrow \infty$ limit, but rather require that the full relativistic action S , written as the c^{-2} -expansion (29), is invariant under the full relativistic transformation rules (18), (19), (20) and (21). Expanding the symmetry variation of S in powers of c^{-2} and requiring that terms at different order in this expansion vanish separately, then indicates how $S^{(0)}$, $S^{(-2)}$ and $S^{(-4)}$ transform into each other under the different c^{-2} orders of the relativistic transformation rules. Let us do this first for the bosonic symmetries (18), (19). The infinitesimal action δ_B of a generic bosonic symmetry B leads to two variations $\delta_B^{(0)}$ and $\delta_B^{(-2)}$, according to

$$\delta_B F = \delta_B^{(0)} F + c^{-2} \delta_B^{(-2)} F, \quad (31)$$

where F is any of the fields τ_μ^A , $e_\mu^{A'}$, $b_{\mu\nu}$, ϕ , $\psi_{\mu\pm}$, λ_\pm . As a consequence

$$\delta_B S = \delta_B^{(0)} S^{(0)} + c^{-2} \left(\delta_B^{(0)} S^{(-2)} + \delta_B^{(-2)} S^{(0)} \right) + \mathcal{O}(c^{-4}). \quad (32) \quad \{\text{eq:bosvarSe}\}$$

The requirement that $\delta_B S = 0$, then imposes that every c^{-2} order in (32) is separately zero. One thus in particular finds that

$$\delta_B^{(0)} S^{(0)} = 0, \quad (33)$$

or in other words, that the NR action $S^{(0)}$ is as expected invariant under the NR bosonic symmetries, whose transformation rules are given by $\delta_B^{(0)}$.

We can apply a similar reasoning to the supersymmetries (20), (21). In this case, the supersymmetry transformations can contain terms at order c^2 , so that the infinitesimal action δ_Q of a generic supersymmetry Q leads to three variations $\delta_Q^{(2)}$, $\delta_Q^{(0)}$ and $\delta_Q^{(-2)}$

$$\delta_Q F = c^2 \delta_Q^{(2)} F + \delta_Q^{(0)} F + c^{-2} \delta_Q^{(-2)} F, \quad (34)$$

where again F is any of the fields τ_μ^A , $e_\mu^{A'}$, $b_{\mu\nu}$, ϕ , $\psi_{\mu\pm}$, λ_\pm . The supersymmetry variation $\delta_Q S$ of the action can then be expanded as

$$\delta_Q S = c^2 \delta_Q^{(2)} S^{(0)} + c^0 \left(\delta_Q^{(0)} S^{(0)} + \delta_Q^{(2)} S^{(-2)} \right) + \mathcal{O}(c^{-2}). \quad (35) \quad \{\text{eq:susyvarS}\}$$

Requiring invariance of S again imposes that every order of c^{-2} in (35) is separately zero. This in particular leads to the following two requirements

$$\delta_Q^{(2)} S^{(0)} = 0 \quad \text{and} \quad \delta_Q^{(0)} S^{(0)} = -\delta_Q^{(2)} S^{(-2)}. \quad (36) \quad \{\text{eq:deltaQco}\}$$

From (28), we see that only $\delta_Q^{(2)} \tilde{\psi}_-$ and $\delta_Q^{(2)} \lambda_-$ are non-zero and that these two variations moreover have the effect of shifting the two fields $\tilde{\psi}_-$ and λ_- independently. The only way, in which the variation $\delta_Q^{(2)} S^{(0)}$ can vanish, is then if $S^{(0)}$ does not depend on $\tilde{\psi}_-$ and λ_- . We can alternatively state this in terms of the fields $\psi_{\mu\pm}$ and λ_\pm . The requirement that $\delta_Q^{(2)} S^{(0)}$ vanishes then boils down to saying that $S^{(0)}$ is invariant under two fermionic shift symmetries, that we call the S - and T -symmetries and whose non-trivial action on $\psi_{\mu\pm}$ and λ_\pm is as follows:

$$\delta_S \psi_{\mu+} = \frac{1}{2} \tau_\mu^+ \Gamma_+ \eta_-, \quad \delta_S \lambda_- = \eta_-,$$

$$\delta_T \psi_{\mu-} = \tau_{\mu}^+ \rho_{-}, \quad (37) \quad \{\text{eq:STsymm}\}$$

where η_{-} and ρ_{-} are the parameters of the S - and T -symmetry respectively. As mentioned above and as can be verified in the explicit expression for $S^{(0)}$ given below, the NR action $S^{(0)}$ is also invariant under an emerging dilatation symmetry that has the following non-trivial action on the fields in $S^{(0)}$:

$$\begin{aligned} \delta_D \phi &= \lambda_D, & \delta_D \tau_{\mu}^A &= \lambda_D \tau_{\mu}^A, \\ \delta_D \psi_{\mu\pm} &= \pm \frac{1}{2} \lambda_D \psi_{\mu\pm}, & \delta_D \lambda_{\pm} &= \pm \frac{1}{2} \lambda_D \lambda_{\pm}. \end{aligned} \quad (38) \quad \{\text{eq:dilsymm}\}$$

Note that the dilatation weights of the NR fields are the same as the exponents of the powers of c in the redefinition (14) of the relativistic fields in terms of the NR ones (for the dilaton, this rule holds when considering, e.g., $\exp(\Phi) = c \exp(\phi)$).

We have seen above that general considerations allow us to conclude that the non-relativistic action $S^{(0)}$ is invariant under the fermionic shift symmetries (37). One might wonder whether a similar, general argument exists for the bosonic dilatation-shift symmetry of the NR action $S^{(0)}$. As we will show now the answer is yes. However, the argument is slightly more subtle than the one for the fermionic Stückelberg shift symmetries. Instead of expanding the supersymmetry invariance of the relativistic action as in eq. (35) we have to consider the supersymmetry algebra acting on the relativistic action—which is of course zero $[\delta_Q(\varepsilon_1), \delta_Q(\varepsilon_2)]S = 0$ —and extract information about the NR action $S^{(0)}$ from the different orders in the expansion. Moreover, we use some information about the algebra, in particular the commutator between S -symmetry and supersymmetry (135b), from which we conclude that

$$[\delta_Q^{(2)}(\varepsilon_1), \delta_Q^{(0)}(\varepsilon_2)] = \delta_D(\lambda'_D) + \delta_M(\lambda'_M) + \dots, \quad (39)$$

where the ellipses denote terms involving symmetries of $S^{(0)}$ that do not play a role in the present discussion, see appendix E for more details. The parameters on the right-hand side are dependent expressions $\lambda'_D = -\lambda'_M = -1/4 \tau^{A'B'-} \bar{\epsilon}_{2+} \Gamma_{A'B'-} \epsilon_{1-}$. Furthermore we can use the fact that the fermionic Stückelberg symmetries commute, see section E.8, to show that $[\delta_Q^{(2)}(\varepsilon_1), \delta_Q^{(2)}(\varepsilon_2)] = 0$. Taking all of the above into account one can then show that the terms in $[\delta_Q(\varepsilon_1), \delta_Q(\varepsilon_2)]S = 0$ at order $\mathcal{O}(c^2)$ vanish if and only if

$$\delta_D S^{(0)} = 0. \quad (40)$$

This shows that the non-relativistic action is indeed dilatation invariant due to the divergence structure in the supersymmetry rules and the particular form of the commutator between supersymmetry and fermionic shift symmetries.

Since we wish to identify $\delta_Q^{(0)}$ as the NR supersymmetry transformation rules, the second requirement of (36) tells us that the NR action $S^{(0)}$ is not necessarily invariant under these NR supersymmetries, but is rather given by the variation $\delta_Q^{(2)} S^{(-2)}$ of the c^{-2} order of the expansion of (8) under the leading c^2 order of the relativistic supersymmetry transformation rules. From (22) we see however that all terms in $\delta_Q^{(2)} S^{(-2)}$ are proportional to $\tau_{A'B'}^-$ or $\tau_{A'+}^-$. We thus find that the variation $\delta_Q^{(0)} S^{(0)}$ of the NR action $S^{(0)}$ under the NR supersymmetry transformation rules gives zero when the following constraints on the torsion $\tau_{\mu\nu}^A$ are imposed

$$\tau_{A'B'}^- = 0, \quad \tau_{A'+}^- = 0. \quad (41) \quad \{\text{eq:DSNC-}\}$$

These constraints are invariant under the dilatation symmetry (38) and we will refer to SNC geometry, in which these constraints are imposed, as ‘self-dual Dilatation invariant SNC geometry’ or self-dual DSNC geometry for short. The constraints (41) are not only invariant under dilatations; they are invariant under all non-relativistic transformation rules and in particular, their variation under NR supersymmetry vanishes identically. One can thus impose the constraints (41) by hand in the theory and still maintain supersymmetry, without having to impose extra constraints. (JR: Add remark that DSNC- is necessary for algebra closure on τ_{μ}^- .)

We can summarize the above discussion as follows. The NR limit $S^{(0)}$ of the ten-dimensional $\mathcal{N} = 1$ supergravity action is obtained as the leading order term in the c^{-2} -expansion of (8), after performing the field redefinition (13). This NR action $S^{(0)}$ is invariant under an emerging dilatation symmetry (38), two emerging fermionic S - and T -shift symmetries (37), as well as under the $c \rightarrow \infty$ limit of the bosonic transformation rules (18), (19). The NR supersymmetry transformation rules are identified as the order c^0 part in the relativistic transformation rules (21). The action $S^{(0)}$ is then only invariant under NR supersymmetry, if one assumes that the self-dual DSNc geometry constraints (41) hold.⁷

Let us finish this section, by giving the explicit expression for the NR action $S_{NR} = S^{(0)}$. It is useful to split S_{NR} into a part S_B that is purely bosonic, a part $S_{\psi\psi}$ that is quadratic in the gravitini $\psi_{\mu\pm}$, a part $S_{\lambda\lambda}$ that is quadratic in the dilatini λ_{\pm} and a remaining quadratic fermion part $S_{\psi\lambda}$ that contains both a gravitino and a dilatino:

$$S_{NR} = S_B + S_{\lambda\lambda} + S_{\lambda\psi} + S_{\psi\psi} + \text{quartic fermion terms}. \quad (42) \quad \{\text{eq:NRAction}\}$$

As mentioned above, we will ignore all quartic fermion terms and only require supersymmetry up to cubic fermion terms. The bosonic part of the action has been given in [6] and reads:

$$S_B = \frac{1}{2\kappa^2} \int d^{10}x e e^{-2\phi} \left(R(J) + 4 \partial_{A'} \phi \partial^{A'} \phi - \frac{1}{12} h_{A'B'C'} h^{A'B'C'} \right. \\ \left. - 4 e^\mu_{A'} (\partial_\mu b^{A'} - \omega_\mu^{A'B'} b_{B'} - \omega_\mu^{AB'} \tau^{A'}_{B'A}) \right. \quad (43) \quad \{\text{eq:BosAction}\}$$

$$\left. - 4 b_{A'} b^{A'} - 4 \tau_{A'\{AB\}} \tau^{A'\{AB\}} \right), \quad (44)$$

where $R(J)$ are other geometric quantities are defined in appendix B, see in particular (115). We refer to [6] for a detailed explanation of the notation. The part of the action that is quadratic in the dilatini reads

$$S_{\lambda\lambda} = \frac{1}{2\kappa^2} \int d^{10}x e e^{-2\phi} \left(2 \bar{\lambda}_\pm \Gamma^{A'} D_{A'} \lambda_\mp + 2 \bar{\lambda}_+ \Gamma^A D_A \lambda_+ \right. \\ \left. - \frac{1}{6} h_{A'B'C'} (\bar{\lambda}_+ \Gamma^{A'B'C'} \lambda_-) + \tau_{B'C'A} (\bar{\lambda}_- \Gamma^{B'C'A} \lambda_-) \right), \quad (45) \quad \{\text{eq:SNR2}\}$$

where the covariant derivatives are covariant with respect to Galilean symmetries and dilatations, see (19) and (112). The notation $\bar{\lambda}_\pm \Gamma \lambda_\mp$ is a shorthand for $\bar{\lambda}_+ \Gamma \lambda_- + \bar{\lambda}_- \Gamma \lambda_+$, and will be used also below. The off-diagonal terms read

$$S_{\lambda\psi} = \frac{1}{2\kappa^2} \int d^{10}x e e^{-2\phi} \left(-4 \bar{\lambda}_\pm \Gamma^{A'B'} e^\mu_{A'} e^\nu_{B'} D_{[\mu} \psi_{\nu]\mp} - 8 \bar{\lambda}_+ \Gamma^{AB'} \tau_{A'}^\mu e_{B'}^\nu D_{[\mu} \psi_{\nu] +} \right. \\ - 4 \bar{\lambda}_\pm \Gamma^{A'B'} \psi_{A'\mp} D_{B'} \phi - 4 \bar{\lambda}_+ \Gamma^{AB'} \psi_{A+} D_{B'} \phi \\ + \frac{1}{6} h_{A'B'C'} (\bar{\lambda}_\pm \Gamma^{A'B'C'D'} \psi_{D'\mp}) + \frac{1}{2} h_{A'B'C'} (\bar{\lambda}_+ \Gamma^{A'B'C'D} \psi_{D+}) \\ - (\eta^{DA} + \epsilon^{DA}) \tau_{B'C'D} (\bar{\lambda}_- \Gamma^{B'C'} \psi_{A+} - \bar{\lambda}_+ \Gamma^{B'C'} \psi_{A-}) \\ + 2 \tau_{B'C'A} \bar{\lambda}_\pm \Gamma^{B'C'} \psi_{A\mp} + 2 \tau^{C'\{AB\}} \bar{\lambda}_+ \Gamma_{C'A} \psi_{B+} \\ \left. - 2 \tau_{B'C'A} \bar{\lambda}_- \Gamma^{AB'C'D'} \psi_{D'-} \right), \quad (46) \quad \{\text{eq:SNR3}\}$$

Finally, let us give the pure gravitino terms

$$S_{\psi\psi} = \frac{1}{2\kappa^2} \int d^{10}x e e^{-2\phi} \left(-2 \bar{\psi}_{A+} \Gamma^{AB'C'} e^\mu_{B'} e^\nu_{C'} D_{[\mu} \psi_{\nu] +} - 4 \bar{\psi}_{A+} \Gamma^{A'B'C} e^\mu_{B'} \tau^\nu_{C'} D_{[\mu} \psi_{\nu] +} \right. \\ \left. - 2 \bar{\psi}_{A'\pm} \Gamma^{A'B'C'} e^\mu_{B'} e^\nu_{C'} D_{[\mu} \psi_{\nu]\mp} + \frac{1}{2} h^{A'B'C'} (\bar{\psi}_{A'\pm} \Gamma_{B'} \psi_{C'\mp}) \right)$$

⁷Note that one needs to treat $S^{(0)}$ as a pseudo-action, when checking its invariance under NR supersymmetry, i.e., one should only impose the constraints (41) after performing a general variation.

$$\begin{aligned}
& -\frac{1}{6}h_{A'B'C'}(\bar{\psi}_{D'+}\Gamma^{A'B'C'D'E}\psi_{E+} + \frac{1}{2}\bar{\psi}_{D'\pm}\Gamma^{A'B'C'D'E'}\psi_{E'\mp}) + \\
& -4(\bar{\psi}_{A'\pm}\Gamma^{A'}\psi_{B'\pm} + \bar{\psi}_{A+}\Gamma^A\psi_{B'+})D^{B'}\phi \\
& -2(\eta^{AD} + \epsilon^{AD})\tau^{B'C'}_D\bar{\psi}_{C'\pm}\Gamma_{B'}\psi_{A\mp} + 2\tau^{B'C'}_A(\bar{\psi}_{B'-}\Gamma_A\psi_{C'-}) \\
& -2(\eta_{BC} - \epsilon_{BC})\tau^{C'\{AB\}}\bar{\psi}^C_+\Gamma_A\psi_{C'+} \\
& + (\eta_{AB} + \epsilon_{AB})\tau_{B'C'}^A\bar{\psi}_{D'\pm}\Gamma^{BB'C'D'E}\psi_{E\mp} \\
& + \tau_{B'C'}^A\bar{\psi}_{D'-}\Gamma_A\Gamma^{B'C'D'E'}\psi_{E'-}), \tag{47} \quad \{\text{eq:SNR4}\}
\end{aligned}$$

Finally, the NR supersymmetry transformation rules that leave S_{NR} invariant (up to cubic fermion terms), upon imposition of the constraints (41), are found in (23) and (25). Note that we can leave out the parts in (23) that involve η_- and ρ_- (whose explicit expressions are given in (24)) from these NR supersymmetry transformation rules as these take the form of S - and T -symmetries.

4 The NR Limit of the Equations of Motion

In the previous section, we discussed the NR limit of the action of ten-dimensional $\mathcal{N} = 1$ supergravity. We saw that the resulting NR action is only invariant under NR supersymmetry after the (supersymmetric) self-dual DSNC geometry constraints (41) have been imposed by hand. Nevertheless, one can derive equations of motion from it, by treating it as a pseudo-action, i.e., by applying the usual unconstrained variational principle and imposing the constraints (41) only after variation. In this section, we will examine these equations of motion in more detail. We will see that they can be derived as NR limits of a subset of the equations of motion of relativistic $D = 10$, $\mathcal{N} = 1$ supergravity. As we will explain, this subset is a proper one, due to the fact that the NR action is invariant under the emerging S -, T - and dilatation symmetries (37), (38). The NR limit of the remaining relativistic equations of motion leads to extra ‘missing equations of motion’. These missing equations of motion consist of two fermionic equations, as well as a bosonic one that can be identified as a supersymmetric generalization of the Poisson equation for the Newton potential of NR gravity. Due to the fact that the NR action is only invariant under NR supersymmetry up to the self-dual DSNC geometry constraints, the equations of motion derived from it do not form a closed set under NR supersymmetry, but can also transform to the missing equations of motion. The full set of missing equations of motion and equations of motion derived from the NR action does however form a supersymmetric set, only if the self-dual DSNC geometry constraints (41) are imposed by hand.

4.1 Equations of motion from the NR action and missing NR equations of motion

Viewing the NR action S_{NR} , given in (42), (43), (45), (46) and (47), as a pseudo-action, we can derive equations of motion for τ_μ^A , $e_\mu^{A'}$, ϕ , $b_{\mu\nu}$, λ_\pm and $\psi_{\mu\pm}$, by computing Euler-Lagrange derivatives, denoted here by $\langle\tau\rangle_{A^\mu}$, $\langle e\rangle_{A'^\mu}$, $\langle\phi\rangle$, $\langle b\rangle^{\mu\nu}$, $\langle\lambda_\pm\rangle$ and $\langle\psi_\pm\rangle^\mu$ respectively, with respect to these fields. Explicitly, we define these functional derivatives as the result of performing an unconstrained variation of S_{NR} as follows:

$$\begin{aligned}
\delta S_{NR} = \frac{1}{2\kappa^2} \int d^{10}x e e^{-2\phi} \Big\{ & \langle\tau\rangle_{A^\mu} \delta\tau_\mu^A + \langle e\rangle_{A'^\mu} \delta e_\mu^{A'} - 8\langle\phi\rangle \delta\phi + \frac{1}{2}\langle b\rangle^{\mu\nu} \delta b_{\mu\nu} \\
& + 4\delta\bar{\lambda}_+ \langle\lambda_- \rangle + 4\delta\bar{\psi}_{\mu+} \langle\psi_- \rangle^\mu + 4\delta\bar{\lambda}_- \langle\lambda_+ \rangle + 4\delta\bar{\psi}_{\mu-} \langle\psi_+ \rangle^\mu \Big\}, \tag{48} \quad \{\text{eq:NReomvar}\}
\end{aligned}$$

where the coefficients of the different terms have been chosen for later convenience. The equations of motion derived from S_{NR} are then given by setting $\langle\tau\rangle_{A^\mu}$, $\langle e\rangle_{A'^\mu}$, $\langle\phi\rangle$, $\langle b\rangle^{\mu\nu}$, $\langle\lambda_\pm\rangle$ and $\langle\psi_\pm\rangle^\mu$ to zero and supplementing the resulting set of equations by hand with the self-dual DSNC geometry constraints (41).

As can be expected, the equations thus found can also be obtained as a NR limit of the equations of motion of relativistic $10D$, $\mathcal{N} = 1$ supergravity. To clarify this, we denote convenient

combinations of the Euler-Lagrange derivatives with respect to $E_\mu^{\hat{A}}$, $B_{\mu\nu}$, Φ , λ and Ψ_μ of the relativistic action (8) by $[\widehat{E}]_{\hat{A}}^\mu$, $[B]^{\mu\nu}$, $[\Phi]$, $[\lambda]$ and $[\Psi]^\mu$ respectively. We define these combinations via the following variation:

$$\delta S = \frac{1}{2\kappa^2} \int d^{10}x E e^{-2\Phi} \left\{ [\widehat{E}]_{\hat{A}}^\mu \delta E_\mu^{\hat{A}} + \frac{1}{2} [B]^{\mu\nu} \delta B_{\mu\nu} - 8[\Phi] \delta \Phi \right. \\ \left. + 4\delta\bar{\lambda} [\lambda] + 4\delta\bar{\Psi}_\mu ([\Psi]^\mu + \Gamma^\mu [\lambda]) \right\}. \quad (49) \quad \{\text{eq:releom}\}$$

Up to the order in fermions we are working in, $[\widehat{E}]_{\hat{A}}^\mu$, $[B]^{\mu\nu}$, $[\Phi]$, $[\lambda]$ and $[\Psi]^\mu$ are explicitly given by

$$[\widehat{E}]_{\hat{A}\mu} \equiv \mathcal{R}_{\hat{A}\mu} + 2\nabla_{\hat{A}} \partial_\mu \Phi - \frac{1}{4} \mathcal{H}_{\hat{A}\rho\sigma} \mathcal{H}_\mu^{\rho\sigma} \quad (+ \text{quadratic fermion terms}), \quad (50a) \quad \{\text{eq:EOM1}\}$$

$$[B]_{\mu\nu} \equiv \nabla^\rho \mathcal{H}_{\rho\mu\nu} - 2(\partial^\rho \Phi) \mathcal{H}_{\rho\mu\nu} \quad (+ \text{quadratic fermion terms}), \quad (50b) \quad \{\text{eq:EOM2}\}$$

$$[\Phi] \equiv \nabla^\mu \partial_\mu \Phi + \frac{1}{4} \mathcal{R} - \partial^\mu \Phi \partial_\mu \Phi - \frac{1}{48} \mathcal{H}_{\mu\nu\rho} \mathcal{H}^{\mu\nu\rho} \quad (+ \text{quadratic fermion terms}), \quad (50c) \quad \{\text{eq:EOM3}\}$$

$$[\lambda] \equiv \not{D}\lambda - \Gamma^{\mu\nu} D_\mu \Psi_\nu - \not{\partial}\Phi \lambda - \Gamma^\mu \not{\partial}\Phi \Psi_\mu - \frac{1}{24} \mathcal{H}^{\mu\nu\rho} (\Gamma^\sigma_{\mu\nu\rho} \Psi_\sigma + \Gamma_{\mu\nu\rho} \lambda), \quad (50d)$$

$$[\Psi]_\mu \equiv 2\Gamma^\nu (D_{[\mu} \Psi_{\nu]} - \frac{1}{8} \mathcal{H}_{\rho\sigma[\mu} \Gamma^{\rho\sigma} \Psi_{\nu]}) - (D_\mu - \frac{1}{8} \mathcal{H}_{\mu\nu\rho} \Gamma^{\nu\rho}) \lambda \\ + \not{\partial}\Phi \Psi_\mu - \frac{1}{12} \Gamma^{\hat{A}\hat{B}\hat{C}} \Psi_\mu \mathcal{H}_{\hat{A}\hat{B}\hat{C}}, \quad (50e)$$

where (as in the relativistic action (8)) the Ricci tensor $\mathcal{R}_{\mu\nu}$ and all covariant derivatives are constructed from the relativistic Levi-Civita (spin) connection. For brevity, we have not explicitly given the quadratic fermion terms in $\widehat{E}_{\hat{A}}^\mu$, $[B]_{\mu\nu}$ and $[\Phi]$.

The NR Euler-Lagrange derivatives $\langle\tau\rangle_{A^\mu}$, $\langle e\rangle_{A'^\mu}$, $\langle\phi\rangle$, $\langle b\rangle^{\mu\nu}$, $\langle\lambda_\pm\rangle$ and $\langle\psi_\pm\rangle^\mu$ can then be obtained from a NR limit, in the sense that they correspond to the leading order terms in a c^{-2} -expansion of particular combinations of their relativistic counterparts $[\widehat{E}]_{\hat{A}}^\mu$, $[B]^{\mu\nu}$, $[\Phi]$, $[\lambda]$ and $[\Psi]^\mu$. To quickly find out which combinations of the relativistic Euler-Lagrange derivatives in this way lead to the NR Euler-Lagrange derivatives, we note that we can use (13), (14) to write

$$\delta E_\mu^{\hat{A}} = c \delta \tau_\mu^{\hat{A}}, \quad \delta E_\mu^{A'} = \delta e_\mu^{A'}, \quad \delta B_{\mu\nu} = -2c \epsilon_{AB} \delta \tau_{[\mu}^{\hat{A}} E_{\nu]}^{\hat{B}} + \delta b_{\mu\nu}, \quad \delta \Phi = \delta \phi, \\ \delta \bar{\lambda} = c^{1/2} \delta \bar{\lambda}_+ + c^{-1/2} \delta \bar{\lambda}_-, \quad \delta \bar{\Psi}_\mu = c^{1/2} \delta \bar{\psi}_{\mu+} + c^{-1/2} \delta \bar{\psi}_{\mu-}. \quad (51)$$

Using this in (49), we can rewrite the variation of the relativistic action S as follows

$$\delta S = \frac{1}{2\kappa^2} \int d^{10}x E e^{-2\Phi} \left\{ c [E]_{A^\mu} \delta \tau_\mu^{\hat{A}} + [E]_{A'^\mu} \delta e_\mu^{A'} + \frac{1}{2} [B]^{\mu\nu} \delta b_{\mu\nu} - 8[\Phi] \delta \phi \right. \\ \left. + 4c^{1/2} \delta \bar{\lambda}_+ \Pi_- [\lambda] + 4c^{1/2} \delta \bar{\psi}_{\mu+} \Pi_- ([\Psi]^\mu + \Gamma^\mu [\lambda]) \right. \\ \left. + 4c^{-1/2} \delta \bar{\lambda}_- \Pi_+ [\lambda] + 4c^{-1/2} \delta \bar{\psi}_{\mu-} \Pi_+ ([\Psi]^\mu + \Gamma^\mu [\lambda]) \right\}, \quad (52) \quad \{\text{eq:releom2}\}$$

where we have introduced the notation⁸

$$[E]_{A^\mu} \equiv [\widehat{E}]_{\hat{A}}^\mu - \epsilon_{AB} E_\nu^{\hat{B}} [B]^{\mu\nu}, \quad [E]_{A'^\mu} \equiv [\widehat{E}]_{A'}^\mu. \quad (53)$$

Note that in (52), the quantities that multiply $\delta \tau_\mu^{\hat{A}}$, $\delta e_\mu^{A'}$, $\delta b_{\mu\nu}$, $\delta \phi$, $\delta \bar{\lambda}_\pm$ and $\delta \bar{\psi}_{\mu\pm}$ have not yet been expanded in powers of c^{-2} and are thus still given in terms of the relativistic fields $E_\mu^{\hat{A}}$, $B_{\mu\nu}$, Φ , λ and Ψ_μ . Performing a c^{-2} -expansion of these quantities in δS , noting that the result should take the form $\delta S = \delta S_{NR} + \mathcal{O}(c^{-2})$ (according to (29) with $S^{(0)} = S_{NR}$) and comparing with (48), we see that

$$\langle\tau\rangle_{A^\mu} = ([E]_{A^\mu})^{(-1)}, \quad \langle\phi\rangle = ([\Phi])^{(0)},$$

⁸The quantities $[E]_{A^\mu}$ and $[E]_{A'^\mu}$ then correspond to the Euler-Lagrange derivatives of the relativistic action (8) with respect to $E_\mu^{\hat{A}}$ and $E_\mu^{A'}$, after viewing (8) as a functional of $b_{\mu\nu} = B_{\mu\nu} + \epsilon_{AB} E_\mu^{\hat{A}} E_\nu^{\hat{B}}$ (instead of as a functional of $B_{\mu\nu}$).

$$\begin{aligned}
\langle e \rangle_{A'\mu} &= ([E]_{A'\mu})^{(0)}, & \langle b \rangle^{\mu\nu} &= ([B]^{\mu\nu})^{(0)}, \\
\langle \psi_- \rangle^\mu &= (\Pi_-([\Psi]^\mu + \Gamma^\mu[\lambda]))^{(-1/2)}, & \langle \lambda_- \rangle &= (\Pi_-[\lambda])^{(-1/2)}, \\
\langle \psi_+ \rangle^\mu &= (\Pi_+([\Psi]^\mu + \Gamma^\mu[\lambda]))^{(+1/2)}, & \langle \lambda_+ \rangle &= (\Pi_+[\lambda])^{(+1/2)},
\end{aligned} \tag{54} \quad \{\text{eq:nrEOM}\}$$

where here and in the following, the notation $(X)^{(n)}$ is used to denote all terms of order c^n in the expression obtained by expanding a relativistic quantity X in powers of c^{-2} , after performing the field redefinition (14). In particular, all quantities on the right-hand-side of the equations in (54) refer to the leading order terms in these c^{-2} -expansions.

The tensors $\langle \tau \rangle_{A\mu}$, $\langle e \rangle_{A'\mu}$, $\langle \phi \rangle$, $\langle b \rangle^{\mu\nu}$, $\langle \lambda_\pm \rangle$ and $\langle \psi_\pm \rangle^\mu$ are not all independent, as there exist various algebraic relations between them. The latter correspond to Noether identities for those local symmetries of the NR action (42), under which none of the fundamental fields τ_μ^A , $e_\mu^{A'}$, ϕ , $b_{\mu\nu}$, λ_\pm and $\psi_{\mu\pm}$ transform as a gauge field. For longitudinal Lorentz transformations, transversal rotations and Galilean boosts, these Noether identities are given by

$$\epsilon^{AB} \langle \tau \rangle_{AB} - 2\bar{\lambda}_+ \langle \lambda_- \rangle + 2\bar{\lambda}_- \langle \lambda_+ \rangle - 2\bar{\psi}_{\mu+} \langle \psi_- \rangle^\mu + 2\bar{\psi}_{\mu-} \langle \psi_+ \rangle^\mu = 0, \tag{55a} \quad \{\text{eq:Noetherid}\}$$

$$\langle e \rangle_{[A'B']} - \bar{\lambda}_+ \Gamma_{A'B'} \langle \lambda_- \rangle - \bar{\lambda}_- \Gamma_{A'B'} \langle \lambda_+ \rangle - \bar{\psi}_{\mu+} \Gamma_{A'B'} \langle \psi_- \rangle^\mu - \bar{\psi}_{\mu-} \Gamma_{A'B'} \langle \psi_+ \rangle^\mu = 0, \tag{55b} \quad \{\text{eq:Noethers}\}$$

$$\langle e \rangle_{A'A} + \epsilon_A^B \langle b \rangle_{BA'} + 2\bar{\lambda}_+ \Gamma_{AA'} \langle \lambda_+ \rangle + 2\bar{\psi}_{\mu+} \Gamma_{AA'} \langle \psi_+ \rangle^\mu = 0. \tag{55c}$$

Note that these 45 Noether identities imply that the 45 components $\langle \tau \rangle_{[AB]}$, $\langle e \rangle_{[A'B']}$ and $\langle e \rangle_{A'A'}$ of the 100 Euler-Lagrange derivatives $\langle \tau \rangle_{A\mu}$ and $\langle e \rangle_{A'\mu}$ can be written in terms of other Euler-Lagrange derivatives. We are thus left with 55 algebraically independent components in $\langle \tau \rangle_{A\mu}$ and $\langle e \rangle_{A'\mu}$, the same number of components that is contained in the relativistic $[G]_{\mu\nu}$.

Naively, one would then say that the Euler-Lagrange derivatives $\langle \tau \rangle_{A\mu}$, $\langle e \rangle_{A'\mu}$, $\langle \phi \rangle$, $\langle b \rangle^{\mu\nu}$, $\langle \lambda_\pm \rangle$ and $\langle \psi_\pm \rangle^\mu$ have as many algebraically independent components as their relativistic counterparts $[G]^{\mu\nu}$, $[B]^{\mu\nu}$, $[\Phi]$, $[\lambda]$ and $[\Psi]^\mu$ and that the NR action (42) thus leads to as many equations of motion as there are relativistic ones. This counting is however not correct as it does not yet take into account extra algebraic Noether identities that are associated to the emergent dilatation and S - and T -symmetries. In particular, the S - and T -symmetries (37) lead to the following relation

$$\langle \lambda_+ \rangle - \frac{1}{2} \Gamma_+ \langle \psi_- \rangle^\mu \tau_\mu^+ = 0, \quad \tau_\mu^+ \langle \psi_+ \rangle^\mu = 0, \tag{56} \quad \{\text{eq:Noethers}\}$$

while the dilatation symmetry (38) implies that

$$\langle \tau \rangle_{A\mu} \tau_\mu^A - 8\langle \phi \rangle + 2\bar{\lambda}_+ \langle \lambda_- \rangle - 2\bar{\lambda}_- \langle \lambda_+ \rangle + 2\bar{\psi}_{\mu+} \langle \psi_- \rangle^\mu - 2\bar{\psi}_{\mu-} \langle \psi_+ \rangle^\mu = 0. \tag{57} \quad \{\text{eq:Noetherd}\}$$

In what follows, it will be useful to simplify this identity, by using the Noether identity (55a) for $\text{SO}(1,1)$ longitudinal Lorentz transformations to eliminate the last four terms. This gives

$$\langle \tau \rangle_{A\mu} \tau_\mu^A + \epsilon^{AB} \langle \tau \rangle_{AB} - 8\langle \phi \rangle = 0 \quad \Leftrightarrow \quad \tau_\mu^- \langle \tau \rangle_-^\mu = 4\langle \phi \rangle. \tag{58} \quad \{\text{eq:Noetherd}\}$$

From (56) and (58), we see that $\langle \psi_+ \rangle^+$ is identically zero and that e.g. $\langle \tau \rangle_-^-$ and $\langle \lambda_+ \rangle$ are not independent.

In (54), we saw that the equations of motion, derived from the action (42), arise from a NR limit that consists of retaining only the leading order terms in the c^{-2} -expansion of the relativistic equations of motion, obtained by setting $[E]_{A\mu}$, $[E]_{A'\mu}$, $[B]^{\mu\nu}$, $[\Phi]$, $[\Psi]^\mu + \Gamma^\mu[\lambda]$ and $[\lambda]$ equal to zero. The Noether identities (56) and (58) then tell us that taking the limit in this way leads to some of the resulting equations being the same or identically zero, so that one is left with less independent NR equations than relativistic ones. It is however also possible to take the NR limit directly at the level of the relativistic equations of motion, in such a way that it preserves the total number of algebraically independent equations. To see how this works, we note that the Noether identity for dilatations says that not all the leading order components in the c^{-2} -expansions of the relativistic $[E]_{A\mu}$ and $[\Phi]$ are linearly independent. Indeed, since $\tau_\mu^- \langle \tau \rangle_-^\mu = (E_\mu^- [E]_-^\mu)^{(0)}$, the identity (58) tells us that the leading (c^0)-order contributions in the expansions of $E_\mu^- [E]_-^\mu$ and $[\Phi]$ are proportional to each other:

$$\tau_\mu^- \langle \tau \rangle_-^\mu = 4\langle \phi \rangle \quad \Leftrightarrow \quad (E_\mu^- [E]_-^\mu)^{(0)} = 4([\Phi])^{(0)}$$

$$\Leftrightarrow \quad E_\mu^- [E]_-^\mu = 4\langle\phi\rangle + \mathcal{O}(c^{-2}) \quad \text{and} \quad [\Phi] = \langle\phi\rangle + \mathcal{O}(c^{-2}). \quad (59) \quad \{\text{eq:rewriteM}\}$$

Similarly, the Noether identities (56) for the S - and T -symmetries are equivalent to saying that the contribution to the c^{-2} -expansion of certain components of $[\Psi]^\mu$ vanishes identically at the order indicated in (54):

$$(\Pi_- E_-^\mu [\Psi]_\mu)^{(1/2)} = 0, \quad \text{and} \quad (\Pi_+ E_-^\mu [\Psi]_\mu)^{(3/2)} = 0. \quad (60) \quad \{\text{eq:rewriteM}\}$$

This then indicates how one can take the NR limit of the equations of motion, such that one ends up with as many NR equations of motion as relativistic ones. The limit of most of the relativistic equations of motion is taken as in (54). As regards the equations $\Pi_\pm E_-^\mu [\Psi]_\mu = 0$ however, one has to take into account that they vanish at the order given in (54) and that one should instead retain the terms at one order lower in the c^{-2} -expansion. Furthermore, instead of applying the NR limit to the equations $\{E_\mu^- [E]_-^\mu = 0, [\Phi] = 0\}$, one should apply it to $\{E_\mu^- [E]_-^\mu = 0, E_\mu^- [E]_-^\mu - 4[\Phi] = 0\}$, so that one ends up with two linearly independent equations. Taking the NR limit of the equations of motion in this way, the set of NR equations of motion, obtained by setting (54) to zero, is then supplemented with the following extra equations:

$$\begin{aligned} \langle\psi_-^{(S)}\rangle &\equiv (\Pi_- E_-^\mu [\Psi]_\mu)^{(-3/2)} = 0, & \langle\psi_+^{(T)}\rangle &\equiv (\Pi_+ E_-^\mu [\Psi]_\mu)^{(-1/2)} = 0, \\ \langle P \rangle &\equiv (E_\mu^- [E]_-^\mu - 4[\Phi])^{(-2)} = 0. \end{aligned} \quad (61) \quad \{\text{eq:missinge}\}$$

We will refer to these as ‘the missing equations of motion’. Although they are not derived from the NR action (42), they are valid NR equations of motion, in the sense that they correspond to the leading order in the c^{-2} -expansion of particular components (of combinations) of the relativistic equations of motion.

Explicitly, the fermionic missing equations of motion are given by:

$$\begin{aligned} \langle\psi_-^{(S)}\rangle &\equiv -D_- \lambda_- + 2\tau_-^\mu e_{A'}^\nu \Gamma^{A'} D_{[\mu} \psi_{\nu]}_- + 2\tau_-^\mu \tau_+^\nu \Gamma^+ D_{[\mu} \psi_{\nu]}_+ \\ &\quad + (\Gamma^{A'} D_{A'} \phi + \frac{1}{24} \Gamma^{A'B'C'} h_{A'B'C'}) \tau_-^\mu \psi_{\mu-} - 2\tau_{A'}^{++} \Gamma^{A'} \tau_+^\mu \psi_{\mu-} \\ &\quad (+ \text{terms of higher order in the fermions}) = 0, \end{aligned} \quad (62a)$$

$$\begin{aligned} \langle\psi_+^{(T)}\rangle &\equiv -D_- \lambda_+ + 2\tau_-^\mu e_{A'}^\nu \Gamma^{A'} D_{[\mu} \psi_{\nu]}_+ + (\Gamma^{A'} D_{A'} \phi + \frac{1}{24} \Gamma^{A'B'C'} h_{A'B'C'}) \tau_-^\mu \psi_{\mu+} \\ &\quad + \tau_{A'}^{++} (\Gamma^{A'-} \lambda_- - \Gamma^{B'} \Gamma^{A'-} \psi_{B'-}) + \frac{1}{2} \tau_{A'B'}^{++} \Gamma^{A'B'-} \tau_-^\mu \psi_{\mu-} \\ &\quad (+ \text{terms of higher order in the fermions}) = 0. \end{aligned} \quad (62b)$$

One can then explicitly check that $\langle\psi_-^{(S)}\rangle$ and $\langle\psi_+^{(T)}\rangle$ are invariant under S - and T -transformations, while they transform covariantly under dilatations, with weights $-3/2$ and $-1/2$ respectively. The explicit expression for the bosonic equation of motion reads:

$$\langle P \rangle \equiv -\tau_A^\mu e_{A'}^\nu R_{\mu\nu}(G)^{AA'} - \epsilon^{AB} \tau_A^\mu \tau_B^\nu R_{\mu\nu}(M) \quad (+ \text{fermionic contributions}) = 0, \quad (63) \quad \{\text{eq:Poisson}\}$$

where $R_{\mu\nu}(G)^{AA'}$ and $R_{\mu\nu}(M)$ are defined in (114). The bosonic part of this equation was identified in [6] as the covariant version of the Poisson equation of NR gravity. The missing bosonic equation can thus be identified with a supersymmetric generalization of this Poisson equation and we will refer to it as ‘the Poisson equation’ in what follows. Under dilatations, $\langle P \rangle$ scales covariantly with weight -2 , while the following transformation rules under the S - and T -symmetries (37) are found (JR: Are the following correct? Do Johannes’ and Luca’s Poisson equations agree in their normalization?)

$$\delta_T \langle P \rangle = 8\bar{\rho}_- \langle\psi_+^{(T)}\rangle, \quad \delta_S \langle P \rangle = 4\bar{\eta}_- \Gamma^- \langle\psi_-^{(S)}\rangle. \quad (64) \quad \{\text{eq:STPoisson}\}$$

The Poisson equation thus transforms to the two fermionic missing equations of motion under the S - and T -symmetries. Since the NR action is invariant under dilatations, S - and T -symmetries, we thus find that the full set of NR field equations, including the missing ones, is invariant under these symmetries.

The above discussion can be made more transparent, by performing a field redefinition in the relativistic action (8). This field redefinition is such that the full set of NR field equations, including the missing ones, is obtained by retaining the leading orders of the c^{-2} -expansions of the equations of motion of the redefined fields (instead of of non-trivial combinations of equations of motion of different fields). We will denote the redefined fields with a tilde, as some of them correspond to rescalings of the NR fields with a tilde, that were defined in (27), with a power of c . This field redefinition is explicitly given by:

$$\begin{aligned}\tilde{E}_\mu^- &\equiv e^{-2\Phi} E_\mu^-, & \tilde{E}_\mu^+ &\equiv E_\mu^+, & \tilde{E}_\mu^{A'} &\equiv E_\mu^{A'}, \\ \tilde{\Phi} &\equiv \Phi, & \tilde{B}_{\mu\nu} &\equiv b_{\mu\nu} \equiv B_{\mu\nu} + \epsilon_{AB} E_\mu^A E_\nu^B, \\ \tilde{\lambda}_\pm &\equiv \lambda_\pm, & \tilde{\Psi}_{\mu+} &\equiv \Pi_+ \Psi_\mu - \frac{1}{2} E_\mu^+ \Gamma_+ \Pi_- \lambda, \\ \tilde{\Psi}_- &\equiv E_+^\mu \Pi_- \Psi_\mu, & \tilde{\Psi}_{\mu-} &\equiv \Pi_- \Psi_\mu - E_\mu^+ E_+^\nu \Pi_- \Psi_\nu.\end{aligned}\tag{65} \quad \{\text{eq:Einstein}\}$$

Note that this field redefinition is invertible. We can then use (14) to write the fields with a tilde as powers of c multiplied with NR fields with a tilde, that can be expressed in terms of the NR fields without a tilde:

$$\begin{aligned}\tilde{E}_\mu^- &\equiv c^{-1} \tilde{\tau}_\mu^- \equiv c^{-1} e^{-2\phi} \tau_\mu^-, & \tilde{E}_\mu^+ &\equiv c \tilde{\tau}_\mu^+ \equiv c \tau_\mu^+, & \tilde{E}_\mu^{A'} &\equiv \tilde{e}_\mu^{A'} \equiv e_\mu^{A'}, \\ \tilde{\Phi} &\equiv \tilde{\phi} + \log c = \phi + \log c, & \tilde{B}_{\mu\nu} &\equiv \tilde{b}_{\mu\nu} \equiv b_{\mu\nu}, \\ \tilde{\lambda}_\pm &\equiv c^{\pm 1/2} \tilde{\lambda}_\pm = c^{\pm 1/2} \lambda_\pm, & \tilde{\Psi}_{\mu+} &\equiv c^{1/2} \tilde{\psi}_{\mu+}, \\ \tilde{\Psi}_- &\equiv c^{-3/2} \tilde{\psi}_-, & \tilde{\Psi}_{\mu-} &\equiv c^{-1/2} \tilde{\psi}_{\mu-}.\end{aligned}\tag{66} \quad \{\text{eq:Einstein}\}$$

The expressions for $\tilde{\psi}_{\mu\pm}$ and $\tilde{\psi}_-$ in terms of NR fields without a tilde are given in (27). As in (14), the dilatation weights of the NR fields with a tilde in the above formulas coincide with the exponents of the powers of c that multiply these fields. Note that (66) contains two types of redefinition that will be used in the following. On the one hand, it expresses how the relativistic fields with a tilde are given in terms of NR fields with a tilde, multiplied with a power of c . On the other hand, it also indicates how the NR fields with a tilde are related to those without a tilde.

By applying (65), we can express the relativistic action (8) in terms of the fields with a tilde and define functional derivatives of the resulting action $S[\tilde{E}_\mu^\pm, \tilde{E}_\mu^{A'}, \tilde{\Phi}, \tilde{B}_{\mu\nu}, \tilde{\Psi}_{\mu\pm}, \tilde{\Psi}_-, \tilde{\lambda}_\pm]$ via the following variation

$$\begin{aligned}\delta S = \frac{1}{2\kappa^2} \int d^{10}x \tilde{E} \Big\{ & [\widetilde{E}]_-^\mu \delta \tilde{E}_\mu^- + [\widetilde{E}]_+^\mu \delta \tilde{E}_\mu^+ + [\widetilde{E}]_{A'}^\mu \delta \tilde{E}_\mu^{A'} + \frac{1}{2} [\widetilde{B}]^{\mu\nu} \delta \tilde{B}_{\mu\nu} - 8 [\widetilde{\Phi}] \delta \tilde{\Phi} \\ & + 4 \delta \tilde{\lambda}_+ [\widetilde{\lambda}_-] + 4 \delta \tilde{\Psi}_{\mu+} [\widetilde{\Psi}_-]^\mu + 4 \delta \tilde{\Psi}_{\mu-} [\widetilde{\Psi}_+]^\mu \\ & + 4 \delta \tilde{\lambda}_- [\widetilde{\lambda}_+] + 4 \delta \tilde{\Psi}_- [\widetilde{\Psi}_+] \Big\},\end{aligned}\tag{67} \quad \{\text{eq:releom3}\}$$

where $\tilde{E} = \det(\tilde{E}_\mu^A)$. Using the rules (65), one finds that the following non-trivial relations hold between the functional derivatives with respect to the fields with tildes and those with respect to the original fields:

$$\begin{aligned}[\widetilde{E}]_-^\mu &= e^{2\Phi} [E]_-^\mu, & [\widetilde{E}]_+^\mu &= [E]_+^\mu + 4 \tilde{\Psi}_- [\widetilde{\Psi}_+]^\mu - 2 \tilde{\lambda}_- \Gamma_+ [\widetilde{\Psi}]^\mu, \\ [\widetilde{\Phi}] &= [\Phi] - \frac{1}{4} E_\mu^- [E]_-^\mu, & [\widetilde{\Psi}_\pm]^\mu &= \Pi_\pm ([\Psi]^\mu + \Gamma^\mu [\lambda]), \\ [\widetilde{\Psi}_+] &= E_\mu^+ \Pi_+ [\Psi]^\mu, & [\widetilde{\lambda}_+] &= -\frac{1}{2} E_\mu^+ \Pi_+ \Gamma_+ [\Psi]^\mu.\end{aligned}\tag{68}$$

For all other functional derivatives, the relation is trivial, e.g., $[\widetilde{E}]_{A'}^\mu = [E]_{A'}^\mu$. From this, we see that the field equations of $\tilde{\Phi}$, $\tilde{\lambda}_-$ and $\tilde{\Psi}_-$ are given by

$$E_\mu^- [E]_-^\mu - 4[\Phi] = 0, \quad \Pi_- E_-^\mu [\Psi]_\mu = 0, \quad \Pi_+ E_-^\mu [\Psi]_\mu = 0,\tag{69}$$

and, according to (61) and the discussion preceding it, thus indeed reproduce the missing equations of motion in their c^{-2} -expansion, as was the goal of the field redefinition (65).

We can then revisit the NR limit of the equations of motion. First, using (66) in (67), we have

$$\begin{aligned} \delta S = \frac{1}{2\kappa^2} \int d^{10}x \tilde{E} \Big\{ c^{-1} [\widetilde{E}]_-{}^\mu \delta \tilde{\tau}_\mu^- + c [\widetilde{E}]_+{}^\mu \delta \tilde{\tau}_\mu^+ + [\widetilde{E}]_{A'}{}^\mu \delta \tilde{e}_\mu^{A'} + \frac{1}{2} [\widetilde{B}]^{\mu\nu} \delta \tilde{b}_{\mu\nu} - 8 [\widetilde{\Phi}] \delta \tilde{\phi} \\ + 4 c^{1/2} \delta \tilde{\lambda}_+ [\widetilde{\lambda}_-] + 4 c^{1/2} \delta \tilde{\psi}_{\mu+} [\widetilde{\Psi}_-]^\mu + 4 c^{-1/2} \delta \tilde{\psi}_{\mu-} [\widetilde{\Psi}_+]^\mu \\ + 4 c^{-1/2} \delta \tilde{\lambda}_- [\widetilde{\lambda}_+] + 4 c^{-3/2} \delta \tilde{\psi}_- [\widetilde{\Psi}_+] \Big\}. \end{aligned} \quad (70) \quad \{\text{eq:releom4}\}$$

As in (52), the quantities that multiply $\delta \tilde{\tau}_\mu^\pm$, $\delta \tilde{e}_\mu^{A'}$, $\delta \tilde{b}_{\mu\nu}$, $\delta \tilde{\phi}$, $\delta \tilde{\lambda}_\pm$, $\delta \tilde{\psi}_{\mu\pm}$ and $\delta \tilde{\psi}_-$ have not yet been expanded in powers of c^{-2} . Requiring compatibility with $\delta S = \delta S_{NR} + c^{-2} \delta S^{(-2)}$ (where S_{NR} and $S^{(-2)}$ are expressed in terms of the NR fields with a tilde) shows that the following expansions hold:

$$\begin{aligned} [\widetilde{E}]_-{}^\mu &= c \langle \widetilde{\tau} \rangle_-{}^\mu + \mathcal{O}(c^{-1}), & [\widetilde{E}]_+{}^\mu &= c^{-1} \langle \widetilde{\tau} \rangle_+{}^\mu + \mathcal{O}(c^{-3}), \\ [\widetilde{E}]_{A'}{}^\mu &= \langle \widetilde{e} \rangle_{A'}{}^\mu + \mathcal{O}(c^{-2}), & [\widetilde{B}]^{\mu\nu} &= \langle \widetilde{b} \rangle^{\mu\nu} + \mathcal{O}(c^{-2}), \\ [\widetilde{\Psi}_\pm]^\mu &= c^{\mp 1/2} \langle \widetilde{\psi}_\pm \rangle^\mu + \mathcal{O}(c^{\mp 1/2-2}), & [\widetilde{\lambda}_-] &= c^{-1/2} \langle \widetilde{\lambda}_- \rangle + \mathcal{O}(c^{-5/2}). \end{aligned} \quad (71) \quad \{\text{eq:bulkeom}\}$$

The quantities $\langle \widetilde{\tau} \rangle_\pm{}^\mu$, $\langle \widetilde{e} \rangle_{A'}{}^\mu$, $\langle \widetilde{b} \rangle^{\mu\nu}$, $\langle \widetilde{\psi}_\pm \rangle^\mu$ and $\langle \widetilde{\lambda}_- \rangle$ then correspond to the functional derivatives of the NR action S_{NR} with respect to $\tilde{\tau}_\mu^\pm$, $\tilde{e}_\mu^{A'}$, $\tilde{b}_{\mu\nu}$, $\tilde{\psi}_{\mu\pm}$ and $\tilde{\lambda}_+$ (after expressing S_{NR} in terms of these tilded NR fields).

The same reasoning would lead one to think that the expansions of $[\widetilde{\Phi}]$, $[\widetilde{\lambda}_+]$ and $[\widetilde{\Psi}_+]$ start at orders c^0 , $c^{1/2}$ and $c^{3/2}$ respectively. This is however not correct. Indeed, if this were true, (70) would imply that S_{NR} depends on $\tilde{\phi}$, $\tilde{\lambda}_-$ and $\tilde{\psi}_-$. This can however not be the case, since these fields shift as Stückelberg fields under dilatations and S - and T -symmetries. Any dependence of S_{NR} on $\tilde{\phi}$, $\tilde{\lambda}_-$ and $\tilde{\psi}_-$ would then imply that S_{NR} is not invariant under these symmetries, contradicting what was found in section 3. We thus conclude that the expansions of $[\widetilde{\Phi}]$, $[\widetilde{\lambda}_+]$ and $[\widetilde{\Psi}_+]$ have to start at one c^{-2} -order higher. This is indeed found explicitly:

$$\begin{aligned} [\widetilde{\Phi}] &= -\frac{1}{4} c^{-2} \langle P \rangle + \mathcal{O}(c^{-4}), & [\widetilde{\lambda}_+] &= \frac{c^{-3/2}}{2} \Gamma_+ \langle \psi_-^{(S)} \rangle + \mathcal{O}(c^{-7/2}), \\ [\widetilde{\Psi}_+] &= c^{-1/2} \langle \psi_+^{(T)} \rangle + \mathcal{O}(c^{-5/2}). \end{aligned} \quad (72) \quad \{\text{eq:misseom}\}$$

From this, we then also see that $\langle P \rangle$, $\langle \psi_-^{(S)} \rangle$ and $\langle \psi_+^{(T)} \rangle$ can be interpreted as functional derivatives of $S^{(-2)}$ with respect to $\tilde{\phi}$, $\tilde{\lambda}_-$ and $\tilde{\psi}_-$. This is similar to what happens when considering the NR expansion of General Relativity, where the Poisson equation of NR gravity is seen to arise from subleading orders in the expansion of the Einstein-Hilbert action [32, 37]. Note also that there is a relation between the dilatation weights of the NR fields with a tilde and the exponent of the power of c in front of their corresponding functional derivatives (either of S_{NR} or $S^{(-2)}$) in (71) and (72). The exponent of the power of c in front of $\langle \widetilde{\tau} \rangle_-{}^\mu$, $\langle \widetilde{\tau} \rangle_+{}^\mu$, $\langle \widetilde{e} \rangle_{A'}{}^\mu$, $\langle \widetilde{b} \rangle^{\mu\nu}$, $\langle \widetilde{\psi}_\pm \rangle^\mu$ and $\langle \widetilde{\lambda}_- \rangle$ in (71) is given by the negative of the dilatation weight of $\tilde{\tau}_\mu^-$, $\tilde{\tau}_\mu^+$, $\tilde{e}_\mu^{A'}$, $\tilde{b}_{\mu\nu}$, $\tilde{\psi}_{\mu\pm}$ and $\tilde{\lambda}_+$ respectively. This rule does not hold for $\langle P \rangle$, $\langle \psi_-^{(S)} \rangle$ and $\langle \psi_+^{(T)} \rangle$: the exponent of the power of c in front of these quantities in (72) is obtained by subtracting two from the negative of the dilatation weight of $\tilde{\phi}$, $\tilde{\lambda}_-$ and $\tilde{\psi}_-$ respectively.

Summarizing: after performing the field redefinition (65), the NR limit of the relativistic equations of motion can more easily be taken in such a way that it preserves the number of algebraically independent equations. The resulting NR field equations are given by

$$\begin{aligned} \langle \widetilde{\tau} \rangle_-{}^\mu &= 0, & \langle \widetilde{\tau} \rangle_+{}^\mu &= 0, & \langle \widetilde{e} \rangle_{A'}{}^\mu &= 0, \\ \langle \widetilde{b} \rangle^{\mu\nu} &= 0, & \langle \widetilde{\psi}_\pm \rangle^\mu &= 0, & \langle \widetilde{\lambda}_- \rangle &= 0, \\ \langle P \rangle &= 0, & \langle \psi_-^{(S)} \rangle &= 0, & \langle \psi_+^{(T)} \rangle &= 0, \end{aligned} \quad (73) \quad \{\text{eq:allNRgeom}\}$$

where each equation corresponds to the leading order terms in the c^{-2} -expansion of a relativistic equation of motion for a redefined field with a tilde. The first two lines correspond to equations that

can be derived from the NR action (42). The equations in the last line are the missing equations of motion, that do not follow from the NR action. The full set of NR field equations (73) is invariant under the emergent dilatation and S - and T -symmetries. In the next subsection, we will address the question whether these equations also transform into each other under NR supersymmetry and Galilean boosts.

4.2 Consistency of all NR equations of motion under supersymmetry and Galilean boosts

Here, we will give a generic argument that the set of NR field equations (73) is invariant under NR supersymmetry and Galilean boosts. As we will see, invariance under NR supersymmetry is not automatically guaranteed: it only holds when the (supersymmetric) self-dual DSNC constraint (41) is imposed by hand.

We will argue exclusively in terms of the fields with tilde (65) and their NR counterparts, defined in (66). It will then be useful to split the tilded fields (65) in two sets, based on whether the limit of their equations of motion can be derived from the NR action (42) or not. The first set of fields is given by $\{\tilde{E}_\mu^+, \tilde{E}_\mu^-, \tilde{E}_\mu^{A'}, \tilde{B}_{\mu\nu}, \tilde{\Psi}_{\mu\pm}, \tilde{\lambda}_+\}$. We will refer to the fields in this set as the (relativistic) ‘bulk fields’ and we will collectively denote them as B_i , with the index i enumerating the different bulk fields. The second set of fields is given by $\{e^{-2\tilde{\Phi}}, \tilde{\Psi}_-, \tilde{\lambda}_-\}$ and their members will be referred to as the (relativistic) ‘missing fields’. We will collectively denote them as M_α , where the index α is used to distinguish the different missing fields. The index I and notation X_I will be used to denote the members of the collection of bulk and missing fields: $\{X_I\} = \{B_i, M_\alpha\}$. As before, the functional derivative of the relativistic action with respect to a field B_i , M_α or X_I will be denoted by $[B_i]$, $[M_\alpha]$, $[X_I]$ respectively. We will split the NR fields with a tilde in a set of NR bulk fields and one of NR missing fields in an analogous manner. The NR bulk fields are given by $\{\tilde{\tau}_\mu^+, \tilde{\tau}_\mu^-, \tilde{e}_\mu^{A'}, \tilde{b}_{\mu\nu}, \tilde{\psi}_{\mu\pm}, \tilde{\lambda}_+\}$ and will be collectively denoted by b_i , whereas the NR missing fields $\{e^{-2\tilde{\phi}}, \tilde{\psi}_-, \tilde{\lambda}_-\}$ will be collectively denoted by m_α . Equations (66) can then be summarized as

$$B_i = c^{b(i)} b_i, \quad M_\alpha = c^{m(\alpha)} m_\alpha, \quad (74) \quad \{\text{eq:expBM}\}$$

where $b(i)$ and $m(\alpha)$ are the dilatation weights of the corresponding NR fields b_i and m_α . According to the remark made below (72), the expansions of the relativistic functional derivatives $[B]^i$ and $[M]^\alpha$ then take the form:

$$\begin{aligned} [B]^i &= c^{-b(i)} \langle b \rangle^i + c^{-b(i)-2} [B]_{SL}^i + \mathcal{O}(c^{-b(i)-4}), \\ [M]^\alpha &= c^{-m(\alpha)-2} \langle m \rangle^\alpha + c^{-m(\alpha)-4} [M]_{SL}^\alpha + \mathcal{O}(c^{-m(\alpha)-6}). \end{aligned} \quad (75) \quad \{\text{eq:expBMeom}\}$$

Here, the collection of $\langle b \rangle^i$ corresponds to $\{\langle \tilde{\tau} \rangle_-^\mu, \langle \tilde{\tau} \rangle_+^\mu, \langle \tilde{e} \rangle_{A'}^\mu, \langle \tilde{b} \rangle^{\mu\nu}, \langle \tilde{\psi}_\pm \rangle^\mu, \langle \tilde{\lambda}_- \rangle\}$, the collection of $\langle m \rangle^\alpha$ to $\{\langle P \rangle, \langle \psi_-^{(S)} \rangle, \langle \psi_+^{(T)} \rangle\}$ and we have denoted the first subleading terms in the expansions of $[B]^i$ and $[M]^\alpha$ by $[B]_{SL}^i$ and $[M]_{SL}^\alpha$.

We then wish to show that the $\langle b \rangle^i$ and $\langle m \rangle^\alpha$ transform into each other under NR supersymmetry and Galilean boosts. To do this, we will rely on a formula, derived in [38], that shows how Euler-Lagrange derivatives, derived from an action, transform into each other under a symmetry of that action. Applied to the $[X]^I$, defined via (67), this formula reads:

$$\delta[X]^I = (\tilde{E} \delta \tilde{E}^{-1}) [X]^I - \frac{\delta(\delta X_J)}{\delta X_I} [X]^J. \quad (76) \quad \{\text{eq:simsekv}\}$$

The last term of (76) is written in the DeWitt notation [39], i.e., the sum over J also entails an integral that is not written out explicitly. Furthermore, δX_I refers to an infinitesimal symmetry transformation of the fields X_I that leaves the relativistic action (8) (expressed in terms of the fields with tilde (65)) invariant. The above formula then specifies that the way, in which the $[X]^I$ transform into each other under the symmetry variation δ , is determined by the functional derivatives $\delta(\delta X_J)/\delta X_I$ of δX_J with respect to X_I .⁹ In the following, we will start from the

⁹In case δX_J involves derivatives of the fields X_I , as is the case for supersymmetry, one can see that the second term of (76) contributes terms that involve derivatives of the symmetry parameters, i.e., non-covariant terms. These non-covariant terms are however still zero on-shell. We refer to [38] for more details.

formula (76), applied to relativistic supersymmetry and boosts, and expand it in powers of c^{-2} . This will allow us to infer how $\langle b \rangle^i$ and $\langle m \rangle^\alpha$ transform into each other under NR supersymmetry and Galilean boosts.

Before showing that all NR field equations (73) transform into each other under NR supersymmetry, we however need to investigate the structure of the NR supersymmetry transformation rules in more detail. First, we note that by writing the c^{-2} -expansions (20), (21) of the relativistic supersymmetry transformation rules (denoted here by δ_Q) in terms of the NR fields with a tilde, the following c^{-2} -expansions are seen to hold:

$$\begin{aligned}\delta_Q b_i &= \delta_Q^{(0)} b_i + c^{-2} \delta_Q^{(-2)} b_i, \\ \delta_Q m_\alpha &= c^2 \delta_Q^{(2)} m_\alpha + \delta_Q^{(0)} m_\alpha + c^{-2} \delta_Q^{(-2)} m_\alpha.\end{aligned}\tag{77} \quad \{\text{eq:cexpdelt}$$

The only non-zero $\delta_Q^{(2)} m^\alpha$ take the form of specific S - and T -transformations and are determined by (28). Note that $\delta_Q^{(2)} m^\alpha$ vanishes when the self-dual DSNC constraint (41) is imposed. The terms $\delta_Q^{(0)} b_i$ and $\delta_Q^{(0)} m_\alpha$ at order c^0 in (77) constitute the NR supersymmetry transformation rules. We will for simplicity ignore the S - and T -transformations (24) that are in principle present in $\delta_Q^{(0)} m_\alpha$. Doing this will not affect our arguments significantly. Using (66), we can then express these NR supersymmetry rules in terms of the fields with a tilde. In what follows, it will turn out to be important that $\delta_Q^{(0)} b_i$ takes the following form

$$\delta_Q^{(0)} b_i = \tilde{\delta} b_i + \delta_{\text{SO}(1,1)} b_i + \delta_{\text{boost}} b_i,\tag{78} \quad \{\text{eq:splitoff}$$

where $\tilde{\delta} b_i$ is independent of the missing fields m_α (so $\delta(\tilde{\delta} b_i)/\delta m_\alpha = 0$) and $\delta_{\text{SO}(1,1)} b_i$ and $\delta_{\text{boost}} b_i$ correspond to a local longitudinal $\text{SO}(1,1)$ transformation and Galilean boost, whose parameters depend on the missing fields m_α . Explicitly, the parameters of $\delta_{\text{SO}(1,1)}$ and δ_{boost} are given (up to bilinear fermion terms in the fermionic transformation rules) by¹⁰

$$\lambda_M = -\bar{\epsilon}_+ \tilde{\lambda}_-, \quad \lambda^{+A'} = 0, \quad \lambda^{-A'} = \bar{\epsilon}_+ \Gamma^{A'} \tilde{\psi}_- + \frac{1}{2} \bar{\epsilon}_- \Gamma^{A'} \Gamma_+ \tilde{\lambda}_-, \tag{79} \quad \{\text{eq:compensa}$$

and the $\tilde{\delta} b_i$ are explicitly found as

$$\begin{aligned}\tilde{\delta} \tilde{\tau}_\mu^+ &= \bar{\epsilon}_+ \Gamma^+ \tilde{\psi}_{\mu+}, & \tilde{\delta} \tilde{\tau}_\mu^- &= -\bar{\epsilon}_- \tilde{\lambda}_+ \tilde{\tau}_\mu^-, \\ \tilde{\delta} \tilde{e}_\mu^{A'} &= \bar{\epsilon}_\pm \Gamma^{A'} \tilde{\psi}_{\mu\mp}, & \tilde{\delta} \tilde{b}_{\mu\nu} &= 4 \tilde{\tau}_{[\mu}^+ \bar{\epsilon}_- \Gamma_+ \tilde{\psi}_{\nu]-} + 2 \tilde{e}_{[\mu}^{A'} \tilde{\delta} \tilde{e}_{\nu]}^{A'}, \\ \tilde{\delta} \tilde{\psi}_{\mu+} &= \tilde{\nabla}_\mu^{(+)} \epsilon_+ + \tilde{T}_\mu \epsilon_- - \frac{1}{2} \tilde{\tau}_\mu^+ \left(\tilde{\not{b}} + \frac{1}{12} \tilde{\not{h}} \right) \Gamma_+ \epsilon_-, & \tilde{\delta} \tilde{\psi}_{\mu-} &= (\delta_\mu^\nu - \tilde{\tau}_\mu^+ \tilde{\tau}_+^\nu) \tilde{\nabla}_\nu^{(+)} \epsilon_-, \\ \tilde{\delta} \tilde{\lambda}_+ &= - \left(\tilde{\not{b}} + \frac{1}{12} \tilde{\not{h}} \right) \epsilon_+ + \frac{1}{2} \tilde{\tau}^{A'B'} \Gamma_{A'B'} \epsilon_-.\end{aligned}\tag{80}$$

Here and in the following, we use the notation that field dependent quantities, such as $\tilde{\tau}_{\mu\nu}^A$, $\tilde{h}_{\mu\nu\rho}$, $\tilde{\omega}_\mu^{A'B'}$, \dots are obtained from their counterparts without a tilde, by replacing all NR fields without a tilde by ones with a tilde. Above, we have then denoted $\tilde{T}_\mu \epsilon_- = (\tilde{e}_\mu^{B'} \tilde{\tau}^{B'A'} + \tilde{\tau}_\mu - \tilde{\tau}^{A'++}) \Gamma_{A'} \epsilon_-$ and the superscript $(+)$ on the covariant derivatives means that these are defined with respect to a modified $\text{SO}(8)$ spin connection $\tilde{\omega}_\mu^{(+A'B')} = \tilde{\omega}_\mu^{A'B'} + 1/2 \tilde{e}_\mu^{C'} \tilde{h}^{C'A'B'}$. We have also used the

¹⁰Note that none of the $\delta_Q^{(0)} b_i$ then depends on the dilaton $\tilde{\phi}$. A priori it is not clear why this is the case, as the dilaton appears explicitly in (25). Moreover, since the spin connections and b_μ in (25) depend on $\tau_\mu^- = e^{2\tilde{\phi}} \tilde{\tau}_\mu^-$, an extra dependence on $\tilde{\phi}$ can be introduced, when writing these connections in terms of the NR fields with a tilde. Explicitly, one however finds that $\omega_\mu = \tilde{\omega}_\mu + \partial_\mu \tilde{\phi}$, $b_\mu = \tilde{b}_\mu + \partial_\mu \tilde{\phi}$, $\omega_\mu^{AA'} = \tilde{\omega}_\mu^{AA'}$ and $\omega_\mu^{A'B'} = \tilde{\omega}_\mu^{A'B'}$. The $\tilde{\phi}$ -dependence in $\delta^{(0)} b_i$ due to the connections then drops out, because ω_μ and b_μ always appear in the combination $b_\mu - \omega_\mu$ in (25). The explicit dilaton dependence in (25) also drops out, when going to the NR fields with a tilde, as this dependence appears in the form $\partial_\mu \phi - b_\mu = \partial_\mu \tilde{\phi} - (\tilde{b}_\mu + \partial_\mu \tilde{\phi}) = -\tilde{b}_\mu$. Alternatively, one can also obtain these results by noting that the redefinition from τ_μ^\pm to $\tilde{\tau}_\mu^\pm$ can be viewed as a symmetry operation, namely the diagonal of an $\text{SO}(1,1)$ transformation with a dilatation with parameters $\lambda_M = \lambda_D = -\phi$. The above explicit expressions for the connections then follow from this observation.

Feynman-slash notation to denote complete contractions with transversal gamma matrices, i.e., $\not{\epsilon} = c_{A'B'...C'}\Gamma^{A'B'...C'}$.

We can now argue that $\langle b \rangle^i$ and $\langle m \rangle^\alpha$ transform into each other under NR supersymmetry $\delta_Q^{(0)}$, when the self-dual DSNC constraint (41)¹¹ is imposed. To do this, we specify (76) to relativistic supersymmetry transformations δ_Q and split the $[X]^I$ into $[B]^i$ and $[M]^\alpha$:

$$\begin{aligned}\delta_Q[B]^i &= (\tilde{E} \delta_Q \tilde{E}^{-1})[B]^i - \frac{\delta(\delta_Q B_j)}{\delta B_i}[B]^j - \frac{\delta(\delta_Q M_\alpha)}{\delta B_i}[M]^\alpha, \\ \delta_Q[M]^\alpha &= (\tilde{E} \delta_Q \tilde{E}^{-1})[M]^\alpha - \frac{\delta(\delta_Q B_i)}{\delta M_\alpha}[B]^i - \frac{\delta(\delta_Q M_\beta)}{\delta M_\alpha}[M]^\beta.\end{aligned}\tag{81}$$

We then expand these equations in powers of c^{-2} , using (74), (75), (77), as well as that generically $\delta_Q[X]^I = c^2 \delta_Q^{(2)}[X]^I + \delta_Q^{(0)}[X]^I + c^{-2} \delta_Q^{(-2)}[X]^I + \mathcal{O}(c^{-4})$. The results on invariance of the NR field equations under supersymmetry will follow from the subleading orders of these expansions. Before discussing this however, let us first check how the leading orders are satisfied and what can be learnt from them.

These leading orders amount to the following equations:

$$\delta_Q^{(2)} \langle b \rangle^i = 0,\tag{82a}$$

$$\delta_Q^{(2)} \langle m \rangle^\alpha = -\frac{\delta(\delta_Q^{(0)} b_i)}{\delta m_\alpha} \langle b \rangle^i - \frac{\delta(\delta_Q^{(2)} m_\beta)}{\delta m_\alpha} \langle m \rangle^\beta.\tag{82b}$$

The second of these equations can be simplified, by making use of the Noether identity for relativistic Lorentz transformations, which states that for a Lorentz transformation δ_L with arbitrary parameters

$$\delta_L X_I [X]^I = 0.\tag{83}$$

This Noether identity can be expanded in powers of c^{-2} , by using (74), (75) and the fact that a generic Lorentz transformation δ_L takes the form $\delta_L = \delta_L^{(0)} + c^{-2} \delta_L^{(-2)}$, with $\delta_L^{(0)}$ corresponding to NR longitudinal SO(1,1) Lorentz transformations, Galilean boosts and transversal rotations. The leading order term in the expansion of (83) then says that

$$\delta_L^{(0)} b_i \langle b \rangle^i = 0,\tag{84}$$

with $\delta_L^{(0)} b_i$ an arbitrary longitudinal SO(1,1) transformation, Galilean boost or transversal rotation of b_i . According to (78), the only dependence of $\delta_Q^{(0)} b_i$ on the fields m_α occurs in field-dependent SO(1,1) and Galilean boost transformations that act on b_i . As a consequence, $\delta(\delta_Q^{(0)} b_i)/\delta m_\alpha$ also corresponds to a longitudinal SO(1,1) or Galilean boost transformation (with m_α stripped from the symmetry parameter), acting on b_i . The first term on the right-hand-side of (82b) is then zero as a consequence of (84), so that we are left with

$$\delta_Q^{(2)} \langle b \rangle^i = 0, \quad \delta_Q^{(2)} \langle m \rangle^\alpha = -\frac{\delta(\delta_Q^{(2)} m_\beta)}{\delta m_\alpha} \langle m \rangle^\beta.\tag{85}$$

Note that the only contributions to $\delta_Q^{(2)} \langle b \rangle^i$ or $\delta_Q^{(2)} \langle m \rangle^\alpha$ can come from their dependence on the m_α , since only these fields transform under $\delta_Q^{(2)}$. The first of (85) is then explained by the fact that $\langle b \rangle^i$ do not depend on the fields m_α , since the NR action (42) from which they are derived is independent of the Stückelberg fields m_α . Furthermore, since $\delta_Q^{(2)} m_\alpha$ is given by field-dependent S - and T -transformations, the second of (85) is consistent with the fact that the missing NR equations of motion transform among themselves under S - and T -symmetries (see, e.g., (64)).

¹¹Note that this constraint assumes the same form in terms of the NR fields with a tilde, i.e., it is given by $\tilde{\tau}_{A'B'} = 0$ and $\tilde{\tau}_{A'+} = 0$.

The subleading terms in the c^{-2} -expansion of (89) give the following equations:

$$\delta_Q^{(0)} \langle b \rangle^i = (\tilde{e} \delta_Q^{(0)} \tilde{e}^{-1}) \langle b \rangle^i - \frac{\delta(\delta_Q^{(0)} b_j)}{\delta b_i} \langle b \rangle^j - \frac{\delta(\delta_Q^{(2)} m_\alpha)}{\delta b_i} \langle m \rangle^\alpha - \delta_Q^{(2)} [B]_{SL}^i, \quad (86a)$$

$$\begin{aligned} \delta_Q^{(0)} \langle m \rangle^\alpha &= (\tilde{e} \delta_Q^{(0)} \tilde{e}^{-1}) \langle m \rangle^\alpha - \frac{\delta(\delta_Q^{(0)} b_j)}{\delta m_\alpha} [B]_{SL}^j - \frac{\delta(\delta_Q^{(-2)} b_i)}{\delta m_\alpha} \langle b \rangle^i \\ &\quad - \frac{\delta(\delta_Q^{(0)} m_\beta)}{\delta m_\alpha} \langle m \rangle^\beta - \frac{\delta(\delta_Q^{(2)} m_\beta)}{\delta m_\alpha} [M]_{SL}^\beta - \delta_Q^{(2)} [M]_{SL}^\alpha. \end{aligned} \quad (86b)$$

These equations tell us how the $\langle b \rangle^i$ and $\langle m \rangle^\alpha$ transform under NR supersymmetry $\delta_Q^{(0)}$.¹² The appearance of the subleading parts $[B]_{SL}^i$ and $[M]_{SL}^\alpha$ of the expansions of $[B]^i$ and $[M]^\alpha$ is worrisome, as it implies that the $\langle b \rangle^i$ and $\langle m \rangle^\alpha$ do not form a closed set under NR supersymmetry. Note however that the second term on the right-hand-side of (86b) is harmless, due to the Noether identity (83). Indeed, the subleading order of the c^{-2} -expansion of (83) implies that

$$\delta_L^{(0)} b_i [B]_{SL}^i = -\delta_L^{(0)} m_\alpha \langle m \rangle^\alpha - \delta_L^{(-2)} b_i \langle b \rangle^i, \quad (87)$$

so that $\delta_L^{(0)} b_i [B]_{SL}^i$, with $\delta_L^{(0)} b_i$ an arbitrary longitudinal SO(1,1) transformation, Galilean boost or transversal rotation of b_i , can be written as a combination of $\langle b \rangle^i$ and $\langle m \rangle^\alpha$. As mentioned below (84), $\delta(\delta_Q^{(0)} b_j)/\delta m_\alpha$ takes the form of a longitudinal SO(1,1) transformation or Galilean boost, acting on b_i , and consequently, the $[B]_{SL}^i$ in the second term on the right-hand-side of (86b) can be traded for $\langle b \rangle^i$ and $\langle m \rangle^\alpha$. The remaining terms that involve $[B]_{SL}^i$ and $[M]_{SL}^\alpha$ in (86) can not necessarily be expressed in terms of $\langle b \rangle^i$ and $\langle m \rangle^\alpha$. They however vanish, when the self-dual DSNC constraint (41) is imposed. For $\delta_Q^{(2)} [B]_{SL}^i$ and $\delta_Q^{(2)} [M]_{SL}^\alpha$, this is because each contribution to these terms is proportional to $\delta_Q^{(2)} m_\alpha$ and thus vanishes when the self-dual DSNC constraint is imposed. From (28), one can see that the functional derivatives $\delta(\delta_Q^{(0)} m_\beta)/\delta m_\alpha$ are zero when the self-dual DSNC constraint holds. The fifth term on the right-hand-side of (86b) then also vanishes, upon imposition of this constraint.

We can thus conclude that the full set of NR field equations (73) is invariant under NR supersymmetry, when the self-dual DSNC constraint (41) is imposed by hand. Let us stress again that this constraint is itself invariant under NR supersymmetry, so that it can be imposed consistently in a supersymmetric fashion, without the need for extra constraints. It is furthermore also interesting to note that from the form of (86a), one can see that the supersymmetry variation of the NR field equations $\langle b \rangle^i = 0$, that follow from the NR action (42), in general gives rise to the missing equations of motion $\langle m \rangle^\alpha = 0$. One can check explicitly that this indeed happens. This phenomenon occurs because the NR action (42) is only invariant under NR supersymmetry up to the self-dual DSNC constraint (41). So, even though the missing equations of motion do not follow directly from the NR action (42), they can be obtained indirectly from it, by varying its equations of motion under NR supersymmetry.

(JR: Please check this argument.) Finally, let us finish this section, by noting that the above arguments can be adapted to show that the $\langle b \rangle^i$ and $\langle m \rangle^\alpha$ also transform into each other under Galilean boosts. First, we note that, by writing the c^{-2} -expansions (18), (19) of the relativistic boost transformation rules (denoted here by δ_B) in terms of the NR fields with a tilde, the following c^{-2} -expansions are seen to hold:

$$\delta_B b_i = \delta_B^{(0)} b_i + c^{-2} \delta_B^{(-2)} b_i, \quad \delta_B m_\alpha = \delta_B^{(0)} m_\alpha + c^{-2} \delta_B^{(-2)} m_\alpha, \quad (88)$$

¹²Note again that we assume that the S - and T -transformations (24) are not included in $\delta_Q^{(0)} m_\alpha$. The effect of including these transformations is that $\delta_Q^{(0)} \langle m \rangle^\alpha$ and $(\delta(\delta_Q^{(0)} m_\beta)/\delta m_\alpha) \langle m \rangle^\beta$ in (86b) can receive extra contributions. These are however always proportional to $\langle m \rangle^\alpha$ (for $\delta_Q^{(0)} \langle m \rangle^\alpha$ this is because the $\langle m \rangle^\alpha$ transform into each other under S - and T -transformations), so that the overall conclusion that all NR field equations transform into each other under supersymmetry is not affected.

where $\delta_B^{(0)}$ corresponds to a Galilean boost transformation. We can then again start from (76), applied to relativistic boosts

$$\delta_B[B]^i = -\frac{\delta(\delta_B B_j)}{\delta B_i}[B]^j - \frac{\delta(\delta_B M_\alpha)}{\delta B_i}[M]^\alpha, \quad \delta_B[M]^\alpha = -\frac{\delta(\delta_B B_i)}{\delta M_\alpha}[B]^i - \frac{\delta(\delta_B M_\beta)}{\delta M_\alpha}[M]^\beta, \quad (89) \quad \{\text{eq:simsekv}\}$$

and expand these equations in powers of c^{-2} , using (74), (75), (88), as well as $\delta_B[X]^I = \delta_B^{(0)}[X]^I + c^{-2}\delta_B^{(-2)}[X]^I + \mathcal{O}(c^{-4})$. The leading order in the c^{-2} -expansion of the first of (89) and the leading and subleading order of the second of (89) then lead to the following equations: \{\text{eq:deltaBbm}\}

$$\delta_B^{(0)}\langle b \rangle^i = -\frac{\delta(\delta_B^{(0)} b_j)}{\delta b_i}\langle b \rangle^j, \quad (90a) \quad \{\text{eq:deltaBb}\}$$

$$\frac{\delta(\delta_B^{(0)} b_i)}{\delta m_\alpha}\langle b \rangle^i = 0, \quad (90b) \quad \{\text{eq:deltaBm1}\}$$

$$\delta_B^{(0)}\langle m \rangle^\alpha = -\frac{\delta(\delta_B^{(0)} b_i)}{\delta m_\alpha}[B]_{SL}^i - \frac{\delta(\delta_B^{(-2)} b_i)}{\delta m_\alpha}\langle b \rangle^i - \frac{\delta(\delta_B^{(0)} m_\beta)}{\delta m_\alpha}\langle m \rangle^\beta. \quad (90c) \quad \{\text{eq:deltaBm2}\}$$

The second of these equations is identically satisfied, as a consequence of (84) (applied to a Galilean boost) and the fact that $\delta(\delta_B^{(0)} b_i)/\delta m_\alpha$ corresponds to a Galilean boost transformation (with the field m_α stripped off its parameter), acting on b_i . Similarly, by using (87) (applied to a Galilean boost), we see that the first term on the right-hand-side of (90c) can be rewritten as a combination of $\langle b \rangle^i$ and $\langle m \rangle^\alpha$. We then see that Galilean boost transformations transform the $\langle b \rangle^i$ and $\langle m \rangle^\alpha$ among themselves. Note that (90) shows that the Galilean boost transformation of the missing equations of motion $\langle m \rangle^\alpha = 0$ includes the equations $\langle b \rangle^i = 0$, while the latter only transform among themselves, since they are derived from the Galilean boost invariant action (42). As a representation of the non-semisimple group that consists of longitudinal $\text{SO}(1,1)$ transformations, Galilean boosts and transversal rotations, the NR field equations (73) are then seen to form a so-called reducible indecomposable representation. Note that the appearance of reducible indecomposable representations is quite common, when discussing finite-dimensional representations of non-semisimple groups [40].

5 Conclusions \{\text{sec:conclusi}\}

In this paper we extended our previous work on taking the NR limit of 10D NS-NS gravity to the supersymmetric case. This leads to a NR minimal supergravity theory that is common to all NR superstring theories. In doing so we encountered two complications that were absent in the bosonic case. First of all, the relation between the string sigma model and the target space effective action is less direct than in the bosonic case. This had the effect that we could not independently check the two fermionic Stueckelberg symmetries that we found in the target space effective action at the level of the sigma model description of the superstring. The second complication is that we found, upon taking the NR limit, divergent terms in the supersymmetry rules of the fermionic fields that were all proportional to the torsion components that define a self-dual DSNC geometry, see eq. (7). These divergent terms could be controlled by (i) using the fact that there are two emergent fermionic Stueckelberg symmetries and (ii) the constraints defining a self-dual DSNC geometry are invariant under all the symmetries of the NR theory and therefore can be imposed by hand without the need to impose more constraints. An important simplifying feature that we used is that, after making a particular field redefinition, see eq. (65), all NR equations of motion occurred as the leading term in the expansion of a corresponding relativistic equation of motion, without the need to take special combinations of equations of motion like we did for the bosonic case. Similar to the bosonic case we found that all divergent terms in the action vanished due to cancellations of different contributions and that the NR action did not give rise to all equations of motion. In particular, the Poisson equation for the Newton potential and two fermionic equations were missing. A difference with the bosonic case is that the so-called missing equations of motion, that do not follow from a variation of the NR action, can be obtained by a supersymmetry variation of the equations of motion that do follow from a variation of the NR action.

In hindsight, it is a good thing that we found divergent terms in the supersymmetry rules. Would such terms have been absent we would have found a NR action that is exactly supersymmetric without the need to impose any constraint. In that case the results of [38] would apply with the consequence that the equations of motion corresponding to the NR action would transform to each other forming a supermultiplet but that none of these equations of motion would transform under supersymmetry to the missing Poisson equation and hence the Poisson equation would not be part of this supermultiplet. Such a situation would be hard to reconcile with the closure of the supersymmetry algebra.

Given the fact that we found an emerging dilatation symmetry and two emerging superconformal symmetries one could wonder in which sense the NR minimal supergravity multiplet we found defines a conformal supergravity multiplet. Apart from a few similarities there are important differences. First of all conformal supergravity is usually presented as an off-shell multiplet whereas the NR minimal supergravity multiplet is on-shell. Secondly, the NR minimal supergravity multiplet lacks the special conformal symmetries. This is a consequence of the fact that, unlike in conformal supergravity, all components of the dilatation gauge field are dependent, see the first equation in (112).

We gave some initial results on the Yang-Mills sector needed to discuss the case of the heterotic superstring. Assuming that in the flat spacetime limit we can define an independent NR Yang-Mills supermultiplet, we showed that, starting from a particular field redefinition, there is a unique NR string limit of the relativistic 10D super-Yang-Mills theory that avoids the occurrence of infinities in the supersymmetry rules. We gave the expressions for the action and supersymmetry rules in a separate Appendix. After a dimensional reduction of the spatial worldsheet direction the bosonic part of this NR 10D super-Yang-Mills theory seems to coincide with the one given in [41, 42] and, more recently, in [13] where it was identified with the low energy dynamics of open strings ending on N coincident D-branes in flat spacetime. The same lower-dimensional bosonic Yang-Mills theory also follows from a null reduction of a relativistic Yang-Mills theory in one dimension higher [43]. These results seem to be consistent with the T-duality of the bosonic NR open string theory as discussed in [14]. Here instead, we are interested in the occurrence of Yang-Mills within the heterotic superstring theory. To construct the Yang-Mills coupled to supergravity system, i.e. the NR heterotic supergravity theory, one may proceed in two ways. Either one couples, via a Noether procedure like in the relativistic case [35], the NR Yang-Mills theory to the minimal supergravity theory that we already constructed in this work or one takes the NR limit of the relativistic Yang-Mills coupled to supergravity theory. Approaching the problem from a sigma model point of view, a new complication, not encountered in the pure supergravity or flat spacetime Yang-Mills theory, occurs, namely the occurrence of a worldsheet anomaly in the sigma model giving rise to a NR Chern-Simons term. We hope to show in a follow-up work how these two different approaches lead to the same NR heterotic supergravity theory with its characteristic NR Yang-Mills Chern-Simons term.

It would be interesting to approach the construction of a NR heterotic supergravity theory from a Double Field Theory point of view where in the bosonic case a construction of the NS-NS gravity theory using Double Field Theory with a degenerate geometry has been given [16]. An intriguing issue arises in the supergravity case. Although we took a NR string limit of the minimal supergravity theory in this paper, it is not clear what the dual null reduction would correspond to. The reason for this is that the null Killing condition corresponds to a constraint on the relativistic supergravity theory that is not invariant under supersymmetry. In fact, we are not aware of any relativistic 10D supergravity theory exhibiting a null isometry direction without breaking supersymmetry.¹³ We hope to come back to this issue in a forthcoming publication.

It is natural to generalize the results of this paper to IIA and IIB supergravity and 11D supergravity corresponding to the IIA and IIB superstring theories and M-theory.¹⁴ In particular, it would be interesting to see what happens with the potential divergent terms in the supersymmetry rules and the occurrence of emergent fermionic Stueckelberg symmetries. Once the finite supersymmetry rules have been constructed one could study compactifications and look for interesting

¹³We stress that this is different from looking to *solutions* of the 10D supergravity theory with a null isometry direction. This leads to supersymmetric Killing spinor conditions that do exist and in general break part of the supersymmetry.

¹⁴For the bosonic part of M-theory, see [44].

NR supersymmetric solutions by analyzing the Killing spinor equations. For the case of minimal supergravity dealt with in this paper we have collected the relevant supersymmetry rules and (bosonic part of) the equations of motion in a separate Appendix. This is a convenient starting point for discussing NR supersymmetric NS-NS solutions. In particular, it would be interesting to see whether one can find NR supersymmetric solutions with a horizon. Such solutions could play an important role in setting up a NR holographic principle independent of the relativistic one. We hope to return to these issues in the nearby future.

Acknowledgements

We would like to thank Jaume Gomis and Ziqi Yan for useful discussions. The work of CŞ is part of the research programme of the Foundation for Fundamental Research on Matter (FOM), which is financially supported by the Netherlands Organisation for Science Research (NWO). The work of LR is supported by the FOM/NWO free program *Scanning New Horizons*.

A Notation and Conventions

A.1 Bosonic Conventions

Ten-dimensional flat Lorentz indices are denoted by \hat{A} and split into (A, A') , where $A = 0, 1$ and $A' = 2, \dots, 9$. We refer to this as a splitting into longitudinal, resp. transversal directions. We use the ‘mostly plus’ form of the Minkowski metric, i.e., $(\eta_{\hat{A}\hat{B}}) = (- + + \dots +)$. The ten-dimensional Levi-Civita epsilon symbol is normalized by $\epsilon_{01\dots 9} = +1$ ($\epsilon^{01\dots 9} = -1$). We also use a two-dimensional longitudinal epsilon symbol ϵ_{AB} that is normalized as $\epsilon_{01} = +1$ ($\epsilon^{01} = -1$). The following identities then hold

$$\epsilon_{AC}\epsilon_{BD} = -\eta_{AB}\eta_{CD} + \eta_{AD}\eta_{BC}, \quad \epsilon_A{}^C\epsilon_{CB} = \eta_{AB}. \quad (91)$$

Instead of writing longitudinal components of tensors with respect to coordinates x^A ($A = 0, 1$), we will also often write them with respect to light-cone coordinates x^\pm that are defined as follows:

$$x^\pm = \frac{1}{\sqrt{2}}(x^0 \pm x^1). \quad (92)$$

The longitudinal η_{AB} -part of the Minkowski metric then has $\eta_{+-} = \eta_{-+} = -1$ as its non-zero components and one similarly has that $\epsilon_{+-} = -\epsilon_{-+} = -1$. It is straightforward to check that

$$X^A\eta_{AB}Y^B = -X^-Y^+ - X^+Y^-, \quad X^A\epsilon_{AB}Y^B = X^-Y^+ - X^+Y^-, \quad (93)$$

$$\frac{1}{2}X^A(\eta_{AB} + \epsilon_{AB})Y^B = X^+Y_+, \quad \frac{1}{2}X^A(\eta_{AB} - \epsilon_{AB})Y^B = X^-Y_-. \quad (94)$$

A curved, lower μ index is turned into flat A or A' indices, using the projective inverse Vielbeine $\tau_A{}^\mu, e_{A'}{}^\mu$ via the rule

$$X_A = \tau_A{}^\mu X_\mu, \quad X_{A'} = e_{A'}{}^\mu X_\mu \quad \Leftrightarrow \quad X_\mu = \tau_\mu{}^A X_A + e_\mu{}^{A'} X_{A'}. \quad (95)$$

A.2 Spinor and Clifford Algebra Conventions

The defining relation for the Clifford algebra that is generated by the ten-dimensional gamma matrices $\Gamma_{\hat{A}}$ is taken to have the following normalization

$$\{\Gamma_{\hat{A}}, \Gamma_{\hat{B}}\} = 2\eta_{\hat{A}\hat{B}}\mathbb{1}. \quad (96)$$

The gamma matrices satisfy the following hermiticity property:

$$\Gamma_{\hat{A}}^\dagger = \Gamma_0\Gamma_{\hat{A}}\Gamma_0. \quad (97)$$

The charge conjugation matrix C is unitary and satisfies

$$C^T = -C, \quad \Gamma_{\hat{A}}^T = -C\Gamma_{\hat{A}}C^{-1}. \quad (98) \quad \{\text{eq:ChargeCo}\}$$

All spinors are assumed to satisfy a Majorana-Weyl condition, so that the conjugate $\bar{\psi}$ of a spinor ψ can be interpreted as $\bar{\psi} = \psi^T C$. Using (98), one can show that spinor bilinears obey the following symmetry property

$$\bar{\chi}_1 \Gamma_{\hat{A}_1 \dots \hat{A}_n} \chi_2 = \begin{cases} +\bar{\chi}_2 \Gamma_{\hat{A}_1 \dots \hat{A}_n} \chi_1 & \text{for } n = 0, 3 \pmod{4}, \\ -\bar{\chi}_2 \Gamma_{\hat{A}_1 \dots \hat{A}_n} \chi_1 & \text{for } n = 1, 2 \pmod{4}. \end{cases} \quad (99) \quad \{\text{eq:flipping}\}$$

Left-, resp. right-handed spinors are obtained by projecting with the Weyl projectors P_L , resp. P_R , where

$$P_L = \frac{1}{2}(\mathbb{1} + \Gamma_*) , \quad P_R = \frac{1}{2}(\mathbb{1} - \Gamma_*) , \quad \text{with } \Gamma_* = -\Gamma_0 \Gamma_1 \dots \Gamma_9. \quad (100) \quad \{\text{eq:Weylproj}\}$$

In this paper, we frequently work with spinors that are projected, using the following ‘worldsheet chirality’ orthogonal projection operators Π_{\pm}

$$\Pi_{\pm} = \frac{1}{2}(\mathbb{1} \pm \Gamma_{01}) \quad (\text{obeying } \Pi_{\pm}^2 = \Pi_{\pm} \text{ and } \Pi_{\pm}\Pi_{\mp} = 0). \quad (101) \quad \{\text{eq:PiProj}\}$$

The projection of a spinor χ with Π_+ , resp. Π_- will be denoted as $\chi_+ = \Pi_+ \chi$, resp. $\chi_- = \Pi_- \chi$. One thus has

$$\chi_{\pm} = \Pi_{\pm} \chi, \quad \Pi_{\pm} \chi_{\mp} = 0, \quad \Gamma_{01} \chi_{\pm} = \pm \chi_{\pm}. \quad (102)$$

Since

$$\Gamma_* \Gamma_{01} = \Gamma_{01} \Gamma_*, \quad \Gamma_A \Gamma_{01} = -\Gamma_{01} \Gamma_A, \quad \Gamma_{A'} \Gamma_{01} = \Gamma_{01} \Gamma_{A'}, \quad \Gamma_{01}^T C = -C \Gamma_{01}, \quad (103)$$

worldsheet chirality projection is compatible with the Majorana-Weyl condition. The conjugate of a worldsheet chirality projected spinor is defined as $\bar{\chi}_{\pm} \equiv \chi_{\pm}^T C$. Note that one then has that

$$\bar{\chi}_{\pm} \Gamma_{01} = \mp \bar{\chi}_{\pm}. \quad (104)$$

As a consequence, bilinears of the form

$$\bar{\chi}_{\pm} \Gamma_{A'_1 \dots A'_m} \psi_{\pm}, \quad \bar{\chi}_{\pm} \Gamma_{A B A'_1 \dots A'_m} \psi_{\pm}, \quad \bar{\chi}_{\pm} \Gamma_{A A'_1 \dots A'_m} \psi_{\mp}, \quad (105) \quad \{\text{eq:projbil}\}$$

are identically zero.

When working with purely longitudinal gamma matrices Γ_A , the following duality relations are often handy:

$$\Gamma_{AB} = \epsilon_{AB} \Gamma_{01}, \quad \Gamma_A = -\epsilon_{AB} \Gamma^B \Gamma_{01}. \quad (106)$$

These can for instance be used to show that

$$\begin{aligned} (\eta_{AB} + \epsilon_{AB}) \chi_+ &= \Gamma_A \Gamma_B \chi_+, & (\eta_{AB} - \epsilon_{AB}) \chi_+ &= \Gamma_B \Gamma_A \chi_+, \\ (\eta_{AB} - \epsilon_{AB}) \chi_- &= \Gamma_A \Gamma_B \chi_-, & (\eta_{AB} + \epsilon_{AB}) \chi_- &= \Gamma_B \Gamma_A \chi_-. \end{aligned} \quad (107) \quad \{\text{eq:etaepsil}\}$$

Instead of using indices $A, B = 0, 1$ for longitudinal gamma matrices, it is sometimes useful to work with longitudinal gamma matrices with light-cone indices

$$\Gamma_{\pm} = \frac{1}{\sqrt{2}}(\Gamma_0 \pm \Gamma_1). \quad (108)$$

These satisfy $\Gamma_{\pm} \Gamma_{01} = \mp \Gamma_{\pm}$ and one thus has

$$\Gamma_+ \chi_+ = 0, \quad \text{and} \quad \Gamma_- \chi_- = 0, \quad (109) \quad \{\text{eq:selfdual}\}$$

implying that e.g. $Y^A \Gamma_A \psi_+ = Y^- \Gamma_- \psi_+$.

B Torsional String Newton-Cartan Geometry

{sec:TSNC}

In this section, we give some details on the non-Lorentzian geometric structures appearing in this paper. We refer to these geometries as ‘torsional string Newton-Cartan’ (TSNC). As the name suggests, these structures have intrinsic torsion and were first introduced in [6]. We refer the reader to appendices B and C in the same paper for more details.

The main novel feature of these structures is the occurrence of a 2-form $b_{\mu\nu}$ and a scalar ϕ as part of the geometric structure, next to the longitudinal τ_μ^A ($A = 0, 1$), and transversal Vielbeinen $e_\mu^{A'}$ ($A' = 2, \dots, 8$). As explained in [6], one can introduce spin connections for local $\text{SO}(1, 1) \times \text{SO}(8)$ –rotations ($\omega_\mu, \omega_\mu^{A'B'}$), Galilean boosts $\omega_\mu^{AA'}$, dilatations b_μ , together with an affine connection $\Gamma_{\mu\nu}^\rho$, by imposing¹⁵

{eq:metcomp}

$$\nabla_\mu \phi \equiv \partial_\mu \phi - b_\mu = e_\mu^{A'} \nabla_{A'} \phi, \quad (110a)$$

$$H_{\mu\nu\rho} \equiv h_{\mu\nu\rho} + 6 \epsilon_{AB} \omega_{[\mu}^{AB'} \tau_{\nu}^B e_{\rho]B'} = \frac{1}{3!} e_\mu^{A'} e_\nu^{B'} e_\rho^{C'} h_{A'B'C'}, \quad (110b)$$

$$\nabla_\mu \tau_\nu^A \equiv \partial_\mu \tau_\nu^A - \omega_\mu^{AB} \tau_{\nu B} - b_\mu \tau_\nu^A - \Gamma_{\mu\nu}^\rho \tau_\rho^A = 0, \quad (110c)$$

{eq:taucomp}

$$\nabla_\mu e_\nu^{A'} \equiv \partial_\mu e_\nu^{A'} - \omega_\mu^{A'B'} e_{\nu B'} + \omega_\mu^{AA'} \tau_{\nu A} - \Gamma_{\mu\nu}^\rho e_\rho^{A'} = 0, \quad (110d)$$

{eq:ecomp}

where $h_{\mu\nu\rho} = 3 \partial_{[\mu} b_{\nu\rho]}$. When considering the antisymmetric part of eqs. (110c) and (110d) we observe that not all components depend on the spin connections and hence cannot be used as conventional constraints. This is different from usual (semi-)Riemannian geometry where the affine connection can be chosen to be symmetric without loss of generality. Here, however, this implies that generic TSNC geometries have intrinsic torsion [45]

$$T_{\mu\nu}^\rho = 2 \Gamma_{[\mu\nu]}^\rho = R_{\mu\nu}(H^A) \tau_A^\rho = (e_\mu^{A'} e_\nu^{B'} \tau_{A'B'}^A + 2 e_{[\mu}^{A'} \tau_{\nu]B} \tau_{A'}^{\{BA\}}) \tau_A^\rho, \quad (111)$$

where we denote $\tau_{\mu\nu}^A = \partial_{[\mu} \tau_{\nu]}^A$, $\tau_{A'B'}^A = e_{A'}^\mu e_{B'}^\nu \tau_{\mu\nu}^A$, etc. as in the main text. Note that the independent components in the torsion tensor are equivalent to what we refer to as the *DSNC*–torsion components in the main text. Phrased differently—setting $T_{\mu\nu}^\rho = 0$ is equivalent to imposing *DSNC* geometry (3). By using the above and solving the remaining conventional constraints in (110), one finds the following explicit expressions for the spin connections

{eq:galspinco}

$$b_\mu = e_\mu^{A'} \tau_{A'A}^A + \tau_\mu^A \partial_A \phi, \quad (112a)$$

$$\omega_\mu = (\tau_\mu^{AB} - \frac{1}{2} \tau_\mu^C \tau^{AB}_C) \epsilon_{AB} - \tau_\mu^A \epsilon_{AB} \partial^B \phi, \quad (112b)$$

$$\omega_\mu^{AA'} = -e_\mu^{AA'} + e_{\mu B'} e^{AA'B'} + \frac{1}{2} \epsilon^A_B h_\mu^{BA'} + \tau_{\mu B} W^{BAA'}, \quad (112c)$$

$$\omega_\mu^{A'B'} = -2 e_\mu^{[A'B']} + e_{\mu C'} e^{A'B'C'} - \frac{1}{2} \tau_\mu^A \epsilon_{AB} h^{BA'B'}, \quad (112d)$$

where $e_{\mu\nu}^{A'} = \partial_{[\mu} e_{\nu]}^{A'}$. However, not all components can be solved for, which is reflected by the undetermined $W^{ABA'}$ which is traceless symmetric in the (A, B) indices, but otherwise arbitrary. We have shown in [6] that no relevant quantity depends on W . The conventional constraints (110) can furthermore be used to give the explicit expression for the affine connection

$$\Gamma_{\mu\nu}^\rho = \tau_A^\rho (\partial_\mu \tau_\nu^A - \omega_\mu^{AB} \tau_{\nu B} - b_\mu \tau_\nu^A) + e_{A'}^\rho (\partial_\mu e_\nu^{A'} - \omega_\mu^{A'B'} e_{\nu B'} + \omega_\mu^{AA'} \tau_{\nu A}). \quad (113)$$

Finally, let us give the full covariant curvatures corresponding to the gauge field of Dilatations (D), $\text{SO}(1, 1)$ –rotations (M), Galilean boosts (G), and $\text{SO}(8)$ –rotations (J). These are the expressions that appear in the bosonic equations of motion:

{eq:curv2form}

$$R_{\mu\nu}(D) = 2 \partial_{[\mu} b_{\nu]} + 2 e_{[\mu}^{A'} \omega_{\nu]}^{BB'} \tau_{A'B'B} + 2 \tau_{[\mu}^A \omega_{\nu]A}^{A'} \nabla_{A'} \phi, \quad (114a)$$

¹⁵Note that it is straightforward to derive supercovariant versions of the above expressions by adding the appropriate gravitino bilinears to the anholonomy coefficients $\tau_{\mu\nu}^A / e_{\mu\nu}^{A'} / h_{\mu\nu\rho} / \partial_\mu \phi$, for example $\hat{\tau}_{\mu\nu}^A = \tau_{\mu\nu}^A - 1/2 \bar{\psi}_\mu \Gamma^A \psi_\nu$. In this work we choose to write out the fermion bilinears explicitly when necessary, and not use supercovariant expressions.

$$\begin{aligned} R_{\mu\nu}(M) = & 2\partial_{[\mu}\omega_{\nu]} + 2\epsilon_{AB}e_{[\mu}^{A'}\omega_{\nu]}^{AB'}\tau_{A'B'}^B \\ & - 4\tau_{[\mu}^A\omega_{\nu]}^{BB'}\epsilon_B^C\tau_{B'\{AC\}} + 2\epsilon_{AB}\tau_{[\mu}^A\omega_{\nu]}^{BB'}\nabla_{B'}\phi, \end{aligned} \quad (114b)$$

$$\begin{aligned} R_{\mu\nu}(G)^{AA'} = & 2\partial_{[\mu}\omega_{\nu]}^{AA'} - 2\epsilon^A{}_B\omega_{[\mu}\omega_{\nu]}^{BA'} - 2\omega_{[\mu}^{A'B'}\omega_{\nu]}^A{}_{B'} + 2b_{[\mu}\omega_{\nu]}^{AA'} \\ & - 4e_{[\mu}^{B'}(\omega_{\nu]}B^{[A'}\tau^{B']\{AB\}} - \frac{1}{4}\epsilon^{AB}\omega_{\nu]}{}_{BC'}h^{A'B'C'}), \end{aligned} \quad (114c)$$

$$\begin{aligned} R_{\mu\nu}(J)^{A'B'} = & 2\partial_{[\mu}\omega_{\nu]}^{A'B'} + 2\omega_{[\mu}^{A'C'}\omega_{\nu]}^{B'C'} \\ & + 2e_{[\mu}^{C'}(2\omega_{\nu]}^{C[A'}\tau^{B']}_{C'C}-\omega_{\nu]}{}_{CC'}\tau^{A'B'C'}) \\ & + 8\tau_{[\mu}^A(\omega_{\nu]}^{B[A'}\tau^{B']\{AB\}} - \frac{1}{8}\epsilon^A{}_B\omega_{\nu]}{}_{BC'}h^{A'B'C'}). \end{aligned} \quad (114d)$$

These curvatures satisfy a number of non-trivial Bianchi identities, some of which can be found in [6]. There are several $\text{SO}(1, 1) \times \text{SO}(8)$ -singlets that one can define from the above, for example

$$R(J) \equiv -e_{A'}{}^\mu e_{B'}{}^\nu R_{\mu\nu}(J)^{A'B'}, \quad (115) \quad \{\text{eq:RJ}\}$$

which appears explicitly in the pseudo-action (43).

C Bosonic Equations of Motion and Killing Spinor Equations

Here, we provide the bosonic truncation of the NR bosonic field equations and the transformation rules of the fermionic fields. The truncated bosonic field equations comprise of the field equations followed from the variation of NR action in (29),

$$\langle\phi\rangle = \nabla^{A'}\nabla_{A'}\phi - (\nabla_{A'}\phi)^2 + \frac{1}{4}R(J) - \frac{1}{48}h_{A'B'C'}h^{A'B'C'} - \tau_{A'\{AB\}}\tau^{A'\{AB\}}, \quad (116a) \quad \{\text{eq:phi}\}$$

$$\langle\tau\rangle_{\{AB\}} = 4(\nabla_{B'} - 2(\nabla_{B'}\phi))\tau^{B'}_{\{AB\}}, \quad (116b) \quad \{\text{eq:STL}\}$$

$$\langle\tau\rangle_{AA'} = 2R_{AC'}(J)_{A'}{}^{C'} - 4\nabla_A\nabla_{A'}\phi - 4\nabla^B\tau_{A'\{AB\}}, \quad (116c) \quad \{\text{eq:V-}\}$$

$$\begin{aligned} \langle e\rangle_{A'B'} = & -2R_{C'(A'}(J)_{B')}{}^{C'} - 4\nabla_{(A'}\nabla_{B')}\phi + \frac{1}{2}h_{A'C'D'}h_{B'}{}^{C'D'} \\ & + 8\tau_{A'\{AB\}}\tau_{B'}{}^{\{AB\}} + 4\delta_{A'B'}\langle\phi\rangle, \end{aligned} \quad (116d) \quad \{\text{eq:GA'B'}\}$$

$$\langle e\rangle_{A'A} = 2(\nabla_{B'} - 2(\nabla_{B'}\phi))\tau^{B'}_{A'A} + 4\tau^{B'}_{\{AB\}}\tau_{B'A'}{}^B + \epsilon_{AB}h_{A'B'C'}\tau^{B'C'B}, \quad (116e) \quad \{\text{eq:V+}\}$$

$$\langle b\rangle_{AB} = 2\epsilon_{AB}\tau_{A'B'C'}\tau^{A'B'C'}, \quad (116f) \quad \{\text{eq:S-}\}$$

$$\langle b\rangle_{A'B'} = (\nabla_{C'} - 2(\nabla_{C'}\phi))h^{C'}_{A'B'} + 2\epsilon^{AB}\nabla_A\tau_{A'B'B}, \quad (116g) \quad \{\text{eq:BA'B'}\}$$

and the Poisson equation, which follows from the supersymmetry transformation of the missing fermionic field equations as explained in section ***,

$$\langle P\rangle \equiv -R_{AA'}(G)^{AA'} - \epsilon^{AB}R_{AB}(M) = 0, \quad (117) \quad \{\text{eq:Poisson}\}$$

where ∇_μ is a covariant derivative for dilatations, $\text{SO}(1, 1) \times \text{SO}(8)$ and Galilean boosts and the covariant curvatures with respect to these bosonic symmetries are defined in Appendix B.

The Killing spinor equations are given by

$$\delta\lambda_- = \nabla_{A'}\phi\Gamma^{A'}\epsilon_- - \frac{1}{12}h_{A'B'C'}\Gamma^{A'B'C'}\epsilon_- + \eta_- = 0, \quad (118a)$$

$$\delta\lambda_+ = \nabla_{A'}\phi\Gamma^{A'}\epsilon_+ - \frac{1}{12}h_{A'B'C'}\Gamma^{A'B'C'}\epsilon_+ + \frac{1}{2}\tau^{A'B'+}\Gamma_{A'B'+}\epsilon_- = 0, \quad (118b)$$

$$\delta\psi_{\mu-} = \mathcal{D}_\mu\epsilon_- - \frac{1}{2}\omega^{\mu-A'}\Gamma_{-A'}\epsilon_+ - \frac{1}{8}e_\mu{}^{C'}h_{A'B'C'}\Gamma^{A'B'}\epsilon_- + \tau_\mu{}^+\rho_- = 0, \quad (118c)$$

$$\delta\psi_{\mu+} = \mathcal{D}_\mu\epsilon_+ - \frac{1}{8}e_\mu{}^{C'}h_{A'B'C'}\Gamma^{A'B'}\epsilon_+ + (e_\mu{}^{B'}\tau_{B'}{}^{A'+} + \tau_\mu{}^-\tau^{A'+})\Gamma_{A'+}\epsilon_- + \frac{1}{2}\tau_\mu{}^+\Gamma_+\eta_- = 0. \quad (118d)$$

The fermionic completion of the bosonic field equations in (116) can be easily obtained from the NR action and does not play a role in defining background solutions. However, the complete Poisson equation can only be derived from the NR action indirectly. Thus, we give terms quadratic in fermions in the Poisson equation for completeness:

$$\langle P \rangle \Big|_{\lambda\lambda} = -4\bar{\lambda}_-\Gamma^A D_A \lambda_- \quad (119a)$$

$$\begin{aligned} \langle P \rangle \Big|_{\lambda\psi} &= 8\bar{\lambda}_-\Gamma^{AB'}\tau^\mu{}_A e^\nu{}_{B'} D_{[\mu}\psi_{\nu]-} + 6\bar{\lambda}_-\Gamma^{AB}\tau^\mu{}_A \tau^\nu{}_{B'} D_{[\mu}\psi_{\nu]+} + 2\bar{\lambda}_+\Gamma^{AB}\tau^\mu{}_A \tau^\nu{}_{B'} D_{[\mu}\psi_{\nu]-} + \\ &+ 4\bar{\psi}_{B'}\Gamma^{AB'} D_A \lambda_- - 4\bar{\psi}_{A-}\Gamma^{AB'} D_{B'} \lambda_- + 6\bar{\psi}_{B-}\Gamma^{AB} D_A \lambda_+ + 2\bar{\psi}_{B+}\Gamma^{AB} D_A \lambda_- \\ &+ 2D_A(\bar{\lambda}_\pm\psi_{A\mp}) + \frac{1}{3}h_{A'B'C'}(\bar{\psi}_{D-}\Gamma^{DA'B'C'}\lambda_-) \end{aligned} \quad (119b)$$

$$\begin{aligned} \langle P \rangle \Big|_{\psi\psi} &= 2D_A(\bar{\psi}_{B'\pm}\Gamma^{B'}\psi_{A\mp} + 2\bar{\psi}_{B+}\Gamma^B\psi_{A+}) + 2D_{C'}(2\bar{\psi}_{A-}\Gamma^A\psi_{C'-} + \bar{\psi}_{A+}\Gamma^{C'AB}\psi_{B-}) + \\ &+ 4\bar{\psi}_{A-}\Gamma^{AB'C'}e^\mu{}_{B'}e^\nu{}_{C'}D_{[\mu}\psi_{\nu]-} + 8\bar{\psi}_{C'-}\Gamma^{AB'C'}\tau^\mu{}_A e^\nu{}_{B'}D_{[\mu}\psi_{\nu]-} + \\ &+ 2\bar{\psi}_{C'+}\Gamma^{ABC'}\tau^\mu{}_A \tau^\nu{}_{B'}D_{[\mu}\psi_{\nu]-} + 4\bar{\psi}_{B+}\Gamma^{ABC'}e^\mu{}_{C'}\tau^\nu{}_A D_{[\mu}\psi_{\nu]-} + \\ &+ 6\bar{\psi}_{C'-}\Gamma^{ABC'}\tau^\mu{}_A \tau^\nu{}_{B'}D_{[\mu}\psi_{\nu]+} + 12\bar{\psi}_{A-}\Gamma^{ABC'}\tau^\mu{}_{B'}e^\nu{}_{C'}D_{[\mu}\psi_{\nu]+} + \\ &+ 8D_{C'}\phi\epsilon^{AB}(\bar{\psi}_{A+}\Gamma^{C'}\psi_{B-}) - 8\tau_{C'}{}^{AB}(\bar{\psi}_{C'-}\Gamma_A\psi_{B-}) \\ &+ \frac{1}{2}h_{A'B'C'}(\bar{\psi}_{D'-}\Gamma^{A'B'C'D'E}\psi_{E-}) + \frac{1}{3}h_{A'B'C'}(\bar{\psi}_{D+}\Gamma^{A'B'C'DE}\psi_{E-}) \end{aligned} \quad (119c)$$

D Non-Relativistic $D = 10$ Super-Yang-Mills

In this section, we present a consistent non-relativistic string limit of ten-dimensional super Yang-Mills (SYM) theory [46] in flat space. Moreover, we will show that the multiplet structure is compatible with the soft algebra derived in section E. Thus it is suggestive that one can, in principle, couple this theory via a Noether procedure to $\mathcal{N} = 1$ supergravity along the lines of [35]. Largely, this section is independent of the rest of the paper. We do, however, use the same notation and conventions as explained in A.

We will show that there is one unique re-scaling Ansatz giving rise to a regular limit $c \rightarrow \infty$. Remarkably, this requires an an-isotropic re-scaling of the gauge field. Typically, such an Ansatz would break the spacetime symmetries of the theory. However, since we are working with a 2-foliation structure, we can introduce an-isotropic re-scalings in the longitudinal lightcone directions, which is consistent with the diagonal part $SO(1,1) \times SO(8)$ of Lorentzian symmetries. We show that this leads to a theory with 16 supercharges, defined on flat space with a two-dimensional foliation. We give the re-scaling Ansatz (122)/(123), the non-relativistic multiplet structure (128), and the action for NR SYM (126).

Let us start by reviewing the relativistic theory [46]. The ten-dimensional SYM on-shell multiplet is described by a gauge field $A_\mu{}^I$ and a gaugino \mathfrak{X}^I , both in the adjoint representation of some gauge group \mathcal{G} . The gaugino is a right-handed Majorana-Weyl fermion. The action

$$S = \frac{1}{g^2} \int d^{10}X \left\{ -\frac{1}{4} F_{\mu\nu}{}^I F^{\mu\nu I} - \bar{\mathfrak{X}}^I \not{D} \mathfrak{X}^I \right\} \quad (120) \quad \{\text{eq:SYMaction}\}$$

is invariant under 16 supercharges with parameter ε

$$\delta A_\mu{}^I = \bar{\varepsilon} \Gamma_\mu \mathfrak{X}^I, \quad \delta \mathfrak{X}^I = -\frac{1}{4} \Gamma^{\mu\nu} \varepsilon F_{\mu\nu}{}^I, \quad (121)$$

where $D_\mu \mathfrak{X}^I = \partial_\mu \mathfrak{X}^I + f_{JK}^I A_\mu{}^J \mathfrak{X}^K$, and $F_{\mu\nu}{}^I = 2\partial_{[\mu} A_{\nu]}{}^I + f_{JK}^I A_\mu{}^J A_\nu{}^K$.

Let us now define a non-relativistic limit by introducing a (dimensionless) contraction parameter

c. We choose this Ansatz such that the limit $c \rightarrow \infty$ is well-defined both in the symmetry transformation rules and the action. In order to get Galilean boost symmetries with parameter $\lambda^{AA'} = c^{-1} \Lambda^{AA'}$ we rescale the flat coordinates $X^\mu = (X^0, X^1, X^{A'=2, \dots, 9})$ as $X^\pm = (X^0 \pm X^1)/\sqrt{2} = c x^\pm$ and $X^{A'} = x^{A'}$. (This can be seen as the flat space limit of (13).) It is not a priori clear how to choose a consistent Ansatz for the fields $(A_\mu^I, \mathfrak{X}^I)$ —hence we parametrize different choices with four to-be determined parameters $\alpha/\beta/\gamma/\delta$

$$A_\mu^I dX^\mu = c^\alpha a^I dx^+ + c^\beta b^I dx^- + c^\gamma c_{A'}^I dx^{A'}, \quad \mathfrak{X}^I = c^{\delta+1/2} \chi_+^I + c^{\delta-1/2} \chi_-^I, \quad (122) \quad \{\text{eq:genericA}\}$$

which is equivalent to expressing $A_\mu^I = (A_+^I, A_-^I, A_{A'}^I)$ as $a^I = c^{1-\alpha} A_+^I$, $b = c^{1-\beta} A_-^I$, $c_{A'}^I = c^{-\gamma} A_{A'}^I$, and $\chi_\pm^I = c^{-\delta \mp 1/2} \Pi_\pm \mathfrak{X}^I$. Note that the re-scaling of χ_+^I and χ_-^I differ by a relative power of c^1 . This has been chosen so that the fermions transform appropriately under Galilean boosts, see eqs. (127).

Let us now constrain the parameters $\alpha/\beta/\gamma/\delta$ by requiring that the limit $c \rightarrow \infty$ is well-defined. In other words, we choose the parameters such that there are no positive powers of c in the symmetry rules. This gives a number of constraints—such as $\beta \geq 2$ from $\delta b^I = c^{2-\beta} \bar{\epsilon}_+ \Gamma_- \chi_+^I$ —that are enough to uniquely fix

$$\alpha = 0, \quad \beta = 2, \quad \gamma = 0, \quad \delta = 0, \quad (123) \quad \{\text{eq:parchoic}\}$$

It is remarkable that we are lead to an an-isotropic limit, where the gauge field in the X^-/x^- direction $b^I = c^{-1} A_-^I$ is re-scaled differently from the rest—see eq. (122). Note, that this is only compatible with the bosonic symmetries of the theory due to the lightlike nature of this direction since under longitudinal $\text{SO}(1,1)$ rotations $\delta A_\pm^I = \pm \Lambda_M A_\pm^I$. It would be interesting to see whether similar limits can be taken for other foliation structures too.

Let us now consider the non-relativistic symmetries that follow after taking the limit $c \rightarrow \infty$. First of all, the an-isotropic rescaling implies that the $b^I = c^{-1} A_-^I$ field becomes a matter field under gauge transformations

$$\begin{aligned} \delta_\theta a^I &= \partial_+ \theta^I - f_{JK}^I \theta^J a^K, & \delta_\theta c_{A'} &= \partial_{A'} \theta^I - f_{JK}^I \theta^J c_{A'}^K, \\ \delta_\theta b^I &= -f_{JK}^I \theta^J a^K, & \delta_\theta \chi_\pm^I &= -f_{JK}^I \theta^J \chi_\pm^K, \end{aligned} \quad (124) \quad \{\text{eq:nrymtraf}\}$$

which motivates the introduction of the following covariant expressions

$$f_{A'B'}^I \equiv 2 \partial_{[A'} c_{B']}^I + f_{JK}^I c_{A'}^J c_{B'}^K \quad (125a)$$

$$f_{A'}^I \equiv \partial_+ c_{A'}^I - \partial_{A'} a^I + f_{JK}^I c_{A'}^J a^K, \quad (125b)$$

$$D_{A'} b^I \equiv \partial_{A'} b^I + f_{JK}^I c_{A'}^J b^K, \quad (125c)$$

$$D_+ b^I \equiv \partial_+ b^I + f_{JK}^I a^J b^K, \quad (125d)$$

and similarly for the fermions. Here and in the following we take $\partial_{A'} = \partial/\partial x^{A'}$ and $\partial_\pm = \partial/\partial x^\pm$. Similarly, it is not hard to see that the limit of the action (120) is well defined with $S_{NR} = \lim_{c \rightarrow \infty} c^{-2} S$,¹⁶ and explicitly given by

$$\begin{aligned} S_{NR} &= \frac{1}{g^2} \int d^{10}x \left\{ -\frac{1}{4} f_{A'B'}^I f^{A'B'I} - f_{A'}^I D_{A'} b^I + D_+ b^I D_+ b^I \right. \\ &\quad \left. - 2 \bar{\chi}_-^I \gamma^{A'} D_{A'} \chi_+^I - \bar{\chi}_+^I \Gamma^+ D_+ \chi_+^I + f_{IJK} b^J \bar{\chi}_-^I \Gamma^- \chi_-^K \right\}. \end{aligned} \quad (126) \quad \{\text{eq:NRSYM}\}$$

Remarkably this action (and, relatedly, none of the equations of motion) contains a derivative in the x^- direction. This makes it tempting to perform a dimensional reduction along the $\partial/\partial x^1$ direction leading to a theory in nine dimensions with a one-dimensional foliation structure. Some aspects of such Galilean gauge theories have been studied in [13, 42, 43, 47, 48].

Let us now study the symmetries of the non-relativistic SYM action (126). The theory is manifestly invariant under Yang-Mills transformations (124) and Galilean symmetries. The action of the

¹⁶Alternatively one could define an effective non-relativistic coupling $g_{YM} = c^{-1} g$.

diagonal part $\text{SO}(1,1) \times \text{SO}(8)$ is as expected from the index structure. The Galilean boosts, however, act non-trivially

$$\begin{aligned}\delta_G a^I &= -\lambda^{-A'} c_{A'}^I, & \delta_G c_{A'}^I &= -\lambda^{-A'} b^I, & \delta_G b^I &= 0, \\ \delta_G \chi_-^I &= \frac{1}{2} \lambda^{-A'} \Gamma_{-A'} \chi_+^I, & \delta_G \chi_+^I &= 0,\end{aligned}\tag{127} \quad \{\text{eq:ymboost}\}$$

showing that the multiplet forms a reducible, yet in-decomposable representation of the Galilei algebra. Furthermore, just as the relativistic parent theory, the non-relativistic action (126) is invariant under 16 supercharges with parameters (ϵ_+, ϵ_-) , and transformation rules

$$\delta a^I = \bar{\epsilon}_- \Gamma_+ \chi_-^I, \tag{128a}$$

$$\delta b^I = \bar{\epsilon}_+ \Gamma_- \chi_+^I, \tag{128b}$$

$$\delta c_{A'}^I = \bar{\epsilon}_+ \Gamma_{A'} \chi_-^I + \bar{\epsilon}_- \Gamma_{A'} \chi_+^I, \tag{128c}$$

$$\delta \chi_+^I = -\frac{1}{4} \Gamma^{A'B'} \epsilon_+ f_{A'B'}^I + \frac{1}{2} \Gamma^{-A'} \epsilon_- D_{A'} b^I + \frac{1}{2} \epsilon_+ D_+ b^I, \tag{128d}$$

$$\delta \chi_-^I = -\frac{1}{4} \Gamma^{A'B'} \epsilon_- f_{A'B'}^I - \frac{1}{2} \Gamma^{+A'} \epsilon_+ f_{A'}^I - \frac{1}{2} \epsilon_- D_+ b^I. \tag{128e}$$

It is not hard to see that these transformations close on the symmetries of the theory. Boost commute with supersymmetry as follows

$$[\delta_G(\lambda^{AA'}), \delta(\epsilon_+)] = \delta(\epsilon'_- = \frac{1}{2} \lambda^{-A'} \Gamma_{-A'} \epsilon_+), \quad [\delta_G(\lambda^{AA'}), \delta(\epsilon_-)] = 0, \tag{129}$$

which shows that one can in principle truncate to a theory with 8 supercharges by setting $\epsilon_+ = 0$. However, this contraction does not allow for an interesting superalgebra closing on spacetime translations, as can be seen from

$$[\delta(\epsilon_+), \delta(\eta_+)] = \xi_{(++)} \partial_+ + \delta_\theta(-\xi_{(++)} a^I), \tag{130a}$$

$$[\delta(\epsilon_-), \delta(\eta_-)] = \delta_\theta(-\xi_{(--) } b^I), \tag{130b}$$

$$[\delta(\epsilon_+), \delta(\eta_-)] = \xi_{(+-)}^{A'} \partial_{A'} + \delta_\theta(-\xi_{(+-)}^{A'} c_{A'}^I), \tag{130c}$$

where $\xi_{(++)} = \bar{\epsilon}_+ \Gamma^+ \eta_+$, $\xi_{(--) } = \bar{\epsilon}_- \Gamma^- \eta_-$, and $\xi_{(+-)}^{A'} = \bar{\epsilon}_+ \Gamma^{A'} \eta_-$. Just as in the relativistic theory these commutators close off-shell on the bosonic fields and on-shell on the fermionic ones. Remarkably this superalgebra is compatible with the $\mathcal{N} = 1$ superalgebra derived in section E. As mentioned above, this is an encouraging observation suggesting that the non-relativistic SYM multiplet can be coupled to the $\mathcal{N} = 1$ supergravity theory studied in this paper along the lines of [35]. We hope to return to these questions soon.

E Closure of the Non-Relativistic Super-Algebra

This section gives some details on how the algebra is realized on fields by describing some commutation relations involving the zeroth order of relativistic supersymmetry rules and some commutators involving S and T-symmetries. It is to be understood that the realizations below hold provided the self-dual DSNC constraint is imposed.

E.1 Boost- $\delta^{(0)}$

The commutator between boosts and supersymmetry closes on symmetries as follow

$$[\delta_B(\lambda^{AB'}), \delta_-^{(0)}(\epsilon_-)] = \delta_T(\rho_-'), \tag{131a}$$

$$[\delta_B(\lambda^{AB'}), \delta_+^{(0)}(\epsilon_+)] = \delta_-^{(0)}(\epsilon'_-) + \delta_S(\eta_-') + \delta_T(\rho_-''), \tag{131b}$$

with parameters

$$\epsilon'_- = \frac{1}{2} \lambda^{A'-} \Gamma_{-A'} \epsilon_+, \tag{132a}$$

$$\rho'_- = \frac{1}{2} \left(\partial_{A'} \lambda_{B'}^- + \lambda_{C'}^- e_{A'B'}^{C'} + 2\lambda_{-A'}^- \tau_{-B'}^- - \lambda^{-C'} \bar{\psi}_{A'-} \Gamma_{C'} \psi_{B'+} \right) \Gamma^{A'B'} \epsilon_- , \quad (132b)$$

$$\rho''_- = \frac{1}{2} \left(\partial^- \lambda^{A'-} + 2\lambda^{B'-} e_+^{A'B'} + 2\lambda^{A'-} \tau_{+-}^- + 2\lambda^{-B'} \tau_+^\nu + e^{\rho A'} \bar{\psi}_{[\nu-} \Gamma_{B'} \psi_{\rho]+} \right) \Gamma_{-A'} \epsilon_+ , \quad (132c)$$

$$\eta'_- = \frac{1}{2} \left(\partial_{A'} \lambda_{B'}^- + \lambda_{C'}^- e_{A'B'}^{C'} + 2\lambda_{-A'}^- \tau_{-B'}^- - \lambda^{-C'} \bar{\psi}_{A'-} \Gamma_{C'} \psi_{B'+} \right) \Gamma^{+A'B'} \epsilon_+ . \quad (132d)$$

E.2 Dilatation- $\delta^{(0)}$

The commutator between the emerging dilatation symmetry and supersymmetry closes as follows

$$[\delta_D(\lambda_D), \delta_+^{(0)}(\epsilon_+)] = \delta_+^{(0)}(\epsilon'_+) + \delta_S(\eta'_-) , \quad (133a)$$

$$[\delta_D(\lambda_D), \delta_-^{(0)}(\epsilon_-)] = \delta_-^{(0)}(\epsilon'_-) + \delta_S(\eta''_+) + \delta_T(\rho_-) , \quad (133b)$$

with parameters

$$\epsilon'_\pm = \mp \frac{1}{2} \lambda_D \epsilon_\pm , \quad (134a)$$

$$\rho_- = -\partial_+ \lambda_D \epsilon_- , \quad (134b)$$

$$\eta'_- = \Gamma^A \tau^\mu_{A-} \partial_\mu \lambda_D \epsilon_+ , \quad (134c)$$

$$\eta''_- = 2\Gamma^{A'} \epsilon_- \partial_{A'} \lambda_D . \quad (134d)$$

E.3 S-symmetry- $\delta^{(0)}$

The commutator between S-symmetry and supersymmetry gives

$$[\delta_S(\eta_-), \delta_-^{(0)}(\epsilon_-)] = \delta_G(\lambda_{(S-)}^{AB'}) + \delta_S(\eta_{(S-)} + \delta_T(\rho_{(S-)})) , \quad (135a)$$

$$[\delta_S(\eta_-), \delta_+^{(+)}(\epsilon_+)] = \delta_L(\lambda_{(S+)}^{AB}) + \delta_D(\lambda_{D(S+)} + \delta_S(\eta_{(S+)} + \delta_T(\rho_{(S+)})) , \quad (135b) \quad \{\text{eq:Ssusy}\}$$

with parameters

$$\lambda_{(S-)}^{A'+} = 0 , \quad (136a)$$

$$\lambda_{(S-)}^{A'-} = \frac{1}{2} \bar{\epsilon}_- \Gamma^{A'-} \eta_- , \quad (136b)$$

$$\eta_{(S-)} = \frac{1}{4} \Gamma^+ [2(\bar{\psi}_{A'+} \eta_-) \Gamma^{-A'} \epsilon_- + (\bar{\epsilon}_- \lambda_+) \Gamma_+ \eta_- + (\bar{\epsilon}_- \Gamma_{A'+} \eta_-) \Gamma^{A'} \lambda_+] , \quad (136c)$$

$$\rho_{(S-)} = \frac{1}{4} [(\bar{\psi}_{A'-} \Gamma_{B'+} \eta_-) \Gamma^{A'B'} \epsilon_- + 2\tau_+^\nu (\bar{\psi}_{\nu+} \eta_-) \epsilon_- + (\bar{\eta}_- \Gamma_+ \lambda_-) \epsilon_- + (\bar{\epsilon}_- \Gamma_{A'+} \eta_-) \Gamma^{A'} \lambda_-] , \quad (136d)$$

$$\lambda_{(S+)}^{AB} = -\frac{1}{2} \epsilon^{AB} (\bar{\epsilon}_+ \eta_-) , \quad (136e)$$

$$\lambda_{D(S+)} = \frac{1}{2} (\bar{\epsilon}_+ \eta_-) , \quad (136f)$$

$$\begin{aligned} \eta_{(S+)} = & -\frac{1}{2} \tau^\mu_A (\bar{\psi}_{\mu+} \eta_-) \Gamma^A \epsilon_+ + \frac{1}{4} (\bar{\psi}_{B'-} \Gamma_{C'A'} \eta_-) \Gamma^{AB'C'} \epsilon_+ - \frac{1}{4} (\bar{\lambda}_- \Gamma_B \eta_-) \Gamma^B \epsilon_+ + \\ & + \frac{3}{4 \cdot 4!} \Gamma^{AB'C'} \epsilon_+ (\bar{\lambda}_- \Gamma_{AB'C'} \eta_-) + \frac{1}{2} (\bar{\epsilon}_+ \eta_-) \lambda_- , \end{aligned} \quad (136g)$$

$$\rho_{(S+)} = \frac{1}{4} (\bar{\psi}_{A-} \Gamma_{B'+} \eta_-) \Gamma^{AB'} \epsilon_+ . \quad (136h)$$

E.4 T-symmetry- $\delta^{(0)}$

Commutation relations between T-symmetry and supersymmetry closes as follow

$$[\delta_T(\rho_-), \delta_+^{(0)}(\epsilon_+)] = \delta_G(\lambda_{(T+)}^{AB'}) + \delta_S(\eta_{(T+)-}) + \delta_T(\rho_{(T+)-}) , \quad (137a)$$

$$[\delta_T(\rho_-), \delta_-^{(0)}(\epsilon_-)] = \delta_T(\rho_{(T-)-}) , \quad (137b)$$

with parameters

$$\lambda_{(T+)}^{A'+} = 0 , \quad (138a)$$

$$\lambda_{(T+)}^{A'-} = -\bar{\epsilon}_+ \Gamma^{A'} \rho_- , \quad (138b)$$

$$\eta_{(T+)-} = \frac{1}{2} \left[\bar{\psi}_{A'+} \Gamma_{B'} \rho_- + \frac{1}{4} (\bar{\lambda}_+ \Gamma_{A'B'} \rho_-) \right] \Gamma^{+A'B'} \epsilon_+ , \quad (138c)$$

$$\rho_{(T-)-} = (\bar{\psi}_{A'+} \Gamma_{B'} \rho_-) \Gamma^{A'B'} \epsilon_- - (\bar{\lambda}_+ \rho_-) \epsilon_- + (\bar{\epsilon}_- \lambda_+) \rho_- + (\bar{\epsilon}_- \Gamma_{A'} \rho_-) \Gamma^A \lambda_+ , \quad (138d)$$

$$\rho_{(T+)-} = \frac{1}{2} [(\bar{\psi}_{B'-} \Gamma^- \rho_-) \Gamma^{+B'} \epsilon_+ + (\bar{\epsilon}_+ \lambda_-) \rho_- + (\bar{\epsilon}_+ \Gamma_{A'} \rho_-) \Gamma^{A'} \lambda_- + (\bar{\psi}_{A'+} \Gamma_{B'} \rho_-) \Gamma^{AB'} \epsilon_+] . \quad (138e)$$

E.5 $\delta^{(0)}\text{-}\delta^{(0)}$

The commutators between two supersymmetry transformations closes on the algebra as

$$[\delta_+^{(0)}(\eta_+), \delta_+^{(0)}(\epsilon_+)] = \mathcal{L}_{\zeta_{(++)}} + \delta_+^{(0)}(\epsilon_{(++)+}) + \delta_-^{(0)}(\epsilon_{(++)-}) + \delta_L(\lambda_{(++)}) + \delta_M(\theta_{(++)}) + [EOM] , \quad (139a) \quad \{\text{eq:commutat}$$

$$[\delta_+^{(0)}(\eta_+), \delta_-^{(0)}(\epsilon_-)] = \mathcal{L}_{\zeta_{(+-)}} + \delta_+^{(0)}(\epsilon_{(+-)+}) + \delta_-^{(0)}(\epsilon_{(+-)-}) + \delta_L(\lambda_{(+-)}) + \delta_M(\theta_{(+-)}) + \delta_T(\rho_{(+-)-}) + \delta_S(\eta_{(+-)-}) + [EOM] , \quad (139b) \quad \{\text{eq:commutat}$$

$$[\delta_-^{(0)}(\eta_-), \delta_-^{(0)}(\epsilon_-)] = \delta_-^{(0)}(\epsilon_{(-)-}) + \delta_L(\lambda_{(-)}) + \delta_M(\theta_{(-)}) + \delta_T(\rho_{(-)-}) + [EOM] , \quad (139c) \quad \{\text{eq:commutat}$$

where δ_L denotes longitudinal Lorentz, transverse Lorentz and boost transformations. The parameters are

$$\zeta_{(++)}^\mu = \xi_{(++)}^A \tau^\mu_A , \quad (140a)$$

$$\lambda_{(++)}^{AB} = -\xi_{(++)}^C \tau^\nu_{C\omega_\nu^{(0)(-)AB}} , \quad (140b)$$

$$\epsilon_{(++)+} = -\xi_{(++)}^B \tau^\nu_{B\bar{\psi}_{\nu+}} + \frac{7}{16} \xi_{(++)}^B \Gamma_B \lambda_- - \frac{1}{32 \cdot 4!} \xi_{(++)}^{B(4')} \Gamma_{(4')B} \lambda_- , \quad (140c)$$

$$\epsilon_{(++)-} = -\xi_{(++)}^B \tau^\nu_{B\psi_{\nu-}} , \quad (140d)$$

$$\lambda_{(++)}^{A'B'} = -\xi_{(++)}^C \tau^\nu_{C\omega_\nu^{(0)(-)A'B'}} , \quad (140e)$$

$$\lambda_{(++)}^{A'B} = -\xi_{(++)}^C \tau^\nu_{C\omega_\nu^{(1)(-)A'B}} , \quad (140f)$$

$$\lambda_{(-)-}^{AB} = 0 , \quad (140g)$$

$$\lambda_{(-)-}^{A'B'} = -\xi_{(-)-}^C \tau^\nu_{C\omega_\nu^{(2)(-)A'B'}} , \quad (140h)$$

$$\lambda_{(-)-}^{A'B} = -\xi_{(-)-}^C \tau^\nu_{C\omega_\nu^{(1)(-)A'B}} , \quad (140i)$$

$$\epsilon_{(-)-} = \frac{7}{16} \xi_{(-)-}^B \Gamma_B \lambda_+ + -\frac{1}{32 \cdot 4!} \xi_{(-)-}^{(4')A} \Gamma_{(4')A} \lambda_+ , \quad (140j)$$

$$\zeta_{(+-)}^\nu = \xi_{(+-)}^{A'} e^\nu_{A'} , \quad (140k)$$

$$\lambda_{(+-)}^{AB} = -\xi_{(+-)}^{C'} e^\nu_{C'\omega_\nu^{(0)(-)AB}} , \quad (140l)$$

$$\lambda_{(+-)}^{AB'} = -\xi_{(+-)}^{C'} e^\nu_{C'\omega_\nu^{(1)(-)AB'}} , \quad (140m)$$

$$\epsilon_{(+-)+} = -\xi_{(+-)}^{C'} e^\nu_{C'\psi_{\nu+}} + \frac{7}{16} \xi_{(+-)}^{C'} \Gamma_{C'} \lambda_+ - \frac{1}{4 \cdot 4!} \xi_{(+-)}^{A'B'C'} \Gamma_{A'B'C'} \lambda_+ , \quad (140n)$$

$$\epsilon_{(+-)-} = -\xi_{(+-)}^{C'} e^\nu_{C'\psi_{\nu-}} + \frac{7}{16} \xi_{(+-)}^{A'} \Gamma_{A'} \lambda_- + \frac{1}{4 \cdot 4!} \xi_{(+-)}^{A'B'C'} \Gamma_{A'B'C'} \lambda_- , \quad (140o)$$

$$\theta_{(++)\mu} = \xi_{(++)}^A \tau^\rho_A b_{\mu\rho} , \quad (140p)$$

$$\theta_{(-)\mu} = -2\xi_{(-)}^A \tau_{\mu A} , \quad (140q)$$

$$\theta_{(+ -)\mu} = -\xi_{(+ -)}^{A'} e_{\mu A'} + \xi_{(+ -)}^{A'} e^{\nu}_{A'} b_{\mu\nu} , \quad (140r)$$

$$\eta_{(- +)-} = -\frac{1}{2} [e^{\mu}_{A'} e^{\rho}_{B'} \partial_{[\nu} (\bar{\eta}_- \Gamma^- \psi_{\rho]-}) + 2\tau_{B'-}^- (\bar{\eta}_- \Gamma^- \psi_{A'-})] \Gamma^{+A'B'} \epsilon_+ , \quad (140s)$$

$$\begin{aligned} \rho_{(- +)-} = & -[\tau^{\nu}_{A'} e^{\rho}_{B'} \partial_{[\nu} (\bar{\eta}_- \Gamma^- \psi_{\rho]-}) + \tau_{B'-}^- (\bar{\eta}_- \Gamma^- \psi_{A-}) + \\ & - \tau_{A-}^- (\bar{\eta}_- \Gamma^- \psi_{B'-})] \Gamma^{AB'} \epsilon_+ , \end{aligned} \quad (140t)$$

$$\begin{aligned} \rho_{(- -)-} = & -\frac{1}{2} [e^{\nu}_{A'} e^{\rho}_{B'} \partial_{[\nu} (\bar{\eta}_- \Gamma^- \psi_{\rho]-}) + 2\tau_{B'-}^- (\bar{\eta}_- \Gamma^- \psi_{A'-})] \Gamma^{AB'} \epsilon_- + \\ & + \frac{1}{2} [e^{\nu}_{A'} e^{\rho}_{B'} \partial_{[\nu} (\bar{\epsilon}_- \Gamma^- \psi_{\rho]-}) + 2\tau_{B'-}^- (\bar{\epsilon}_- \Gamma^- \psi_{A'-})] \Gamma^{AB'} \eta_- , \end{aligned} \quad (140u)$$

and we have used \mathcal{L}_X to denote Lie derivative, $[EOM]$ to define terms proportional to the equations of motion, the notation

$$\xi_{(\pm\pm)}^{\hat{A}_1 \dots \hat{A}_n} = \bar{\epsilon}_{\pm} \Gamma^{\hat{A}_1 \dots \hat{A}_n} \eta_{\pm} , \quad (141)$$

$$\xi_{(\pm\mp)}^{\hat{A}_1 \dots \hat{A}_n} = \bar{\epsilon}_{\pm} \Gamma^{\hat{A}_1 \dots \hat{A}_n} \eta_{\mp} , \quad (142)$$

$$\Gamma^{(n')(m)} = \Gamma^{A'_1 \dots A'_n B_1 \dots B_m} , \quad (143)$$

and

$$\omega_{\mu}^{(0)(-)}{}_{A'B'} = \omega_{\mu}^{(0)}{}^{A'B'} - \frac{1}{2} h_{\mu A'B'} , \quad (144a)$$

$$\omega_{\mu}^{(-1)(-)}{}_{AB'} = \omega_{\mu}^{(-1)}{}^{AB'} - \frac{1}{2} h_{\mu AB'} , \quad (144b)$$

$$\omega_{\mu}^{(0)(-)}{}_{AB} = \omega_{\mu}^{(0)}{}^{AB} - \epsilon_{CD} \tau_{\mu}^C \tau_{AB}^D + \epsilon_{AD} \tau_{\mu B}^D - \epsilon_{BD} \tau_{\mu A}^D , \quad (144c)$$

$$\omega_{\mu}^{(2)(-)}{}_{A'B'} = \omega_{\mu}^{(2)}{}^{A'B'} - \epsilon_{CD} \tau_{\mu}^C \tau_{AB'}^D , \quad (144d)$$

$$\omega_{\mu}^{(1)(-)}{}_{AB'} = \omega_{\mu}^{(1)}{}^{AB} - \epsilon_{CD} \tau_{\mu}^C \tau_{AB'}^D + \epsilon_{AD} \tau_{\mu B'}^D . \quad (144e)$$

E.6 Dilatation-S/T Symmetry

The commutation rules between dilatation and S- and T-Symmetries are

$$[\delta_D(\lambda_D), \delta_T(\rho_-)] = \delta_T \left(\frac{3}{2} \lambda_D \rho_- \right) , \quad (145a)$$

$$[\delta_D(\lambda_D), \delta_S(\eta_-)] = \delta_S \left(\frac{1}{2} \lambda_D \eta_- \right) . \quad (145b)$$

E.7 Boost-S/T Symmetry

The commutation rules between boost and S and T-symmetries are

$$[\delta_G(\lambda_{AB'}), \delta_T(\rho_-)] = 0 , \quad (146a)$$

$$[\delta_G(\lambda_{AB'}), \delta_S(\eta_-)] = \delta_T \left(-\frac{1}{4} \lambda_{AB'} \Gamma^{AB'} \Gamma_+ \eta_- \right) . \quad (146b)$$

E.8 S/T Symmetry-S/T Symmetry

S and T-symmetries define an Abelian algebra:

$$[\delta_T(\rho_-), \delta_T(\rho'_-)] = 0 , \quad (147a)$$

$$[\delta_S(\eta_-), \delta_S(\eta'_-)] = 0 , \quad (147b)$$

$$[\delta_T(\rho_-), \delta_S(\eta_-)] = 0 . \quad (147c)$$

{sec:algstst}

References

- [1] J. Gomis and H. Ooguri, *Nonrelativistic closed string theory*, *J. Math. Phys.* **42** (2001) 3127 [[hep-th/0009181](#)].
- [2] U. H. Danielsson, A. Guijosa and M. Kruczenski, *IIA/B, wound and wrapped*, *JHEP* **10** (2000) 020 [[hep-th/0009182](#)].
- [3] J. Gomis, J. Gomis and K. Kamimura, *Non-relativistic superstrings: A New soluble sector of $AdS_5 \times S^5$* , *JHEP* **12** (2005) 024 [[hep-th/0507036](#)].
- [4] E. Bergshoeff, J. Gomis and Z. Yan, *Nonrelativistic String Theory and T-Duality*, *JHEP* **11** (2018) 133 [[1806.06071](#)].
- [5] E. A. Bergshoeff, J. Gomis, J. Rosseel, C. Şimşek and Z. Yan, *String Theory and String Newton-Cartan Geometry*, *J. Phys. A* **53** (2020) 014001 [[1907.10668](#)].
- [6] E. A. Bergshoeff, J. Lahnsteiner, L. Romano, J. Rosseel and C. Şimşek, *A non-relativistic limit of NS-NS gravity*, *JHEP* **06** (2021) 021 [[2102.06974](#)].
- [7] L. Bidussi, T. Harmark, J. Hartong, N. A. Obers and G. Oling, *Torsional string Newton-Cartan geometry for non-relativistic strings*, [2107.00642](#).
- [8] T. Harmark, J. Hartong and N. A. Obers, *Nonrelativistic strings and limits of the AdS/CFT correspondence*, *Phys. Rev. D* **96** (2017) 086019 [[1705.03535](#)].
- [9] T. Harmark, J. Hartong, L. Menculini, N. A. Obers and Z. Yan, *Strings with Non-Relativistic Conformal Symmetry and Limits of the AdS/CFT Correspondence*, *JHEP* **11** (2018) 190 [[1810.05560](#)].
- [10] T. Harmark, J. Hartong, L. Menculini, N. A. Obers and G. Oling, *Relating non-relativistic string theories*, *JHEP* **11** (2019) 071 [[1907.01663](#)].
- [11] J. Gomis, J. Oh and Z. Yan, *Nonrelativistic String Theory in Background Fields*, *JHEP* **10** (2019) 101 [[1905.07315](#)].
- [12] Z. Yan and M. Yu, *Background Field Method for Nonlinear Sigma Models in Nonrelativistic String Theory*, *JHEP* **03** (2020) 181 [[1912.03181](#)].
- [13] J. Gomis, Z. Yan and M. Yu, *Nonrelativistic Open String and Yang-Mills Theory*, *JHEP* **03** (2021) 269 [[2007.01886](#)].
- [14] J. Gomis, Z. Yan and M. Yu, *T-Duality in Nonrelativistic Open String Theory*, *JHEP* **02** (2021) 087 [[2008.05493](#)].
- [15] A. D. Gallegos, U. Gürsoy and N. Zinnato, *Torsional Newton Cartan gravity from non-relativistic strings*, *JHEP* **09** (2020) 172 [[1906.01607](#)].
- [16] A. D. Gallegos, U. Gürsoy, S. Verma and N. Zinnato, *Non-Riemannian gravity actions from double field theory*, *JHEP* **06** (2021) 173 [[2012.07765](#)].
- [17] K. Cho and J.-H. Park, *Remarks on the non-Riemannian sector in Double Field Theory*, *Eur. Phys. J. C* **80** (2020) 101 [[1909.10711](#)].
- [18] S. M. Ko, C. Melby-Thompson, R. Meyer and J.-H. Park, *Dynamics of Perturbations in Double Field Theory & Non-Relativistic String Theory*, *JHEP* **12** (2015) 144 [[1508.01121](#)].
- [19] K. Morand and J.-H. Park, *Classification of non-Riemannian doubled-yet-gauged spacetime*, *Eur. Phys. J. C* **77** (2017) 685 [[1707.03713](#)].
- [20] C. D. A. Blair, G. Oling and J.-H. Park, *Non-Riemannian isometries from double field theory*, *JHEP* **04** (2021) 072 [[2012.07766](#)].

- [21] J.-H. Park and S. Sugimoto, *String Theory and non-Riemannian Geometry*, *Phys. Rev. Lett.* **125** (2020) 211601 [2008.03084].
- [22] J. Klusoň, *Remark About Non-Relativistic String in Newton-Cartan Background and Null Reduction*, *JHEP* **05** (2018) 041 [1803.07336].
- [23] J. Klusoň, *Note About T-duality of Non-Relativistic String*, *JHEP* **08** (2019) 074 [1811.12658].
- [24] J. Klusoň, *(m,n)-String and D1-Brane in Stringy Newton-Cartan Background*, *JHEP* **04** (2019) 163 [1901.11292].
- [25] D. Roychowdhury, *Probing tachyon kinks in Newton-Cartan background*, *Phys. Lett. B* **795** (2019) 225 [1903.05890].
- [26] J. Hartong and E. Have, *On the Non-Relativistic Expansion of Closed Bosonic Strings*, 2107.00023.
- [27] R. Andringa, E. Bergshoeff, J. Gomis and M. de Roo, *‘Stringy’ Newton-Cartan Gravity*, *Class. Quant. Grav.* **29** (2012) 235020 [1206.5176].
- [28] J. Bruges, T. Curtright, J. Gomis and L. Mezincescu, *Non-relativistic strings and branes as non-linear realizations of Galilei groups*, *Phys. Lett. B* **594** (2004) 227 [hep-th/0404175].
- [29] J. Bruges, J. Gomis and K. Kamimura, *Newton-Hooke algebras, non-relativistic branes and generalized pp-wave metrics*, *Phys. Rev. D* **73** (2006) 085011 [hep-th/0603023].
- [30] T. Frankel, *The Geometry of Physics: An Introduction*. Cambridge University Press, 3 ed., 2011, 10.1017/CBO9781139061377.
- [31] Z. Yan, *Torsional Deformation of Nonrelativistic String Theory*, 2106.10021.
- [32] D. Hansen, J. Hartong and N. A. Obers, *Action Principle for Newtonian Gravity*, *Phys. Rev. Lett.* **122** (2019) 061106 [1807.04765].
- [33] J. Gomis, K. Kamimura and P. K. Townsend, *Non-relativistic superbranes*, *JHEP* **11** (2004) 051 [hep-th/0409219].
- [34] C. D. Blair, *A worldsheet supersymmetric Newton-Cartan string*, *JHEP* **10** (2019) 266 [1908.00074].
- [35] E. Bergshoeff, M. de Roo, B. de Wit and P. van Nieuwenhuizen, *Ten-Dimensional Maxwell-Einstein Supergravity, Its Currents, and the Issue of Its Auxiliary Fields*, *Nucl. Phys. B* **195** (1982) 97.
- [36] A. H. Chamseddine, *N=4 Supergravity Coupled to N=4 Matter*, *Nucl. Phys. B* **185** (1981) 403.
- [37] D. Hansen, J. Hartong and N. A. Obers, *Non-Relativistic Gravity and its Coupling to Matter*, *JHEP* **06** (2020) 145 [2001.10277].
- [38] B. Vanhecke and A. Van Proeyen, *Covariant field equations in supergravity*, *Fortsch. Phys.* **65** (2017) 1700071 [1705.06675].
- [39] B. S. DeWitt, *Quantum Theory of Gravity. 2. The Manifestly Covariant Theory*, *Phys. Rev.* **162** (1967) 1195.
- [40] C. George and M. Lévy-Nahas, *Finite-dimensional representations of some non-semisimple Lie algebras*, *Journal of Mathematical Physics* **7** (1966) 980.
- [41] E. S. Santos, M. de Montigny, F. C. Khanna and A. E. Santana, *Galilean covariant Lagrangian models*, *J. Phys. A* **37** (2004) 9771.

- [42] E. Bergshoeff, J. Rosseel and T. Zojer, *Non-relativistic fields from arbitrary contracting backgrounds*, *Class. Quant. Grav.* **33** (2016) 175010 [[1512.06064](#)].
- [43] G. Festuccia, D. Hansen, J. Hartong and N. A. Obers, *Symmetries and Couplings of Non-Relativistic Electrodynamics*, *JHEP* **11** (2016) 037 [[1607.01753](#)].
- [44] C. D. A. Blair, D. Gallegos and N. Zinnato, *A non-relativistic limit of M-theory and 11-dimensional membrane Newton-Cartan*, [2104.07579](#).
- [45] J. Figueroa-O'Farrill, *On the intrinsic torsion of spacetime structures*, [2009.01948](#).
- [46] L. Brink, J. H. Schwarz and J. Scherk, *Supersymmetric Yang-Mills Theories*, *Nucl. Phys. B* **121** (1977) 77.
- [47] S. Chapman, L. Di Pietro, K. T. Grosvenor and Z. Yan, *Renormalization of Galilean Electrodynamics*, *JHEP* **10** (2020) 195 [[2007.03033](#)].
- [48] N. Lambert, A. Lipstein and P. Richmond, *Non-Lorentzian M5-brane Theories from Holography*, *JHEP* **08** (2019) 060 [[1904.07547](#)].