

⑦

Variation of Parameters

This method only for the general second order linear differential equation with variable coefficients

$$a_0(x) \frac{d^2 y}{dx^2} + a_1(x) \frac{dy}{dx} + a_2(x)y = b(x). \quad (1)$$

Suppose that y_1 and y_2 are linearly independent solutions of the corresponding homogeneous equation

$$a_0(x) \frac{d^2 y}{dx^2} + a_1(x) \frac{dy}{dx} + a_2(x)y = 0 \quad (2)$$

We assume a solution of the form

$$y_p = v_1 y_1 + v_2 y_2$$

where v_1 and v_2 are functions of x . Then

$$y_p' = v_1 y_1' + v_2 y_2' + v_1' y_1 + v_2' y_2.$$

If we assume $v_1' y_1 + v_2' y_2 = 0$, then

$$y_p' = v_1 y_1' + v_2 y_2' \quad (3)$$

Again differentiating (3), we obtain

$$y_p'' = v_1 y_1'' + v_2 y_2'' + v_1' y_1' + v_2' y_2'. \quad (4)$$

We impose the basic condition that y_p be a solution of (1).

Thus we substitute (4), (3) in y_p respectively.

$$v_1 [a_0 y_1'' + a_1 y_1' + a_2 y_1] + v_2 [a_0 y_2'' + a_1 y_2' + a_2 y_2] + a_0 [v_1' y_1' + v_2' y_2'] = b.$$

$= 0$ $= 0$ which are solutions of (2)

Thus $v_1' y_1' + v_2' y_2' = \frac{b}{a_0}$. Thus we have.

$$y_1 v_1' + y_2 v_2' = 0 \quad (\text{by assumption})$$

$$y_1' v_1' + y_2' v_2' = \frac{b}{a_0}.$$

$$\text{Then } v_1' = \frac{\begin{vmatrix} 0 & y_2 \\ \frac{b}{a_0} & y_2' \end{vmatrix}}{\begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}} = \frac{-b y_2}{a_0 w(y_1, y_2)}$$

$$\text{Similarly } v_2' = \frac{b y_1}{a_0 w(y_1, y_2)}.$$

⑧

Examples $y'' + y = \tan x$

$$m^2 + 1 = 0 \quad m = \pm i \quad y_c = C_1 \sin x + C_2 \cos x \quad (y_1 = \sin x, y_2 = \cos x)$$

$$W(y_1, y_2) = \begin{vmatrix} \sin x & \cos x \\ \cos x & -\sin x \end{vmatrix} = -\sin^2 x - \cos^2 x = -1 //$$

$$V_1' = \frac{-b y_2}{a_0 W(y_1, y_2)} = \frac{-\tan x \cdot \cos x}{1 \cdot (-1)} = +\sin x \Rightarrow V_1(x) = -\cos x + C_3.$$

$$V_2' = \frac{b y_1}{a_0 W(y_1, y_2)} = \frac{+\tan x \cdot \sin x}{-1} = \frac{-\sin^2 x}{\cos x} = \frac{\cos^2 x - 1}{\cos x} = \cos x - \sec x$$

$$\Rightarrow V_2 = \sin x - \ln |\sec x + \tan x| + C_4.$$

$$y_p = C_3 \sin x + C_4 \cos x - (\cos x) (\ln |\sec x + \tan x|)$$

The general solution is

$$y = C_1 \sin x + C_2 \cos x - (\cos x) (\ln |\sec x + \tan x|)$$

$$(C_1 = C_1 + C_3, C_2 = C_2 + C_4).$$

$$(2) \quad y''' - 6y'' + 11y' - 6y = e^x$$

$$m^3 - 6m^2 + 11m - 6 = (m-1)(m-2)(m-3) \quad y_1 = e^x \quad y_2 = e^{2x} \quad y_3 = e^{3x}$$

$$y_c = C_1 e^x + C_2 e^{2x} + C_3 e^{3x}$$

$$y_p = V_1 e^x + V_2 e^{2x} + V_3 e^{3x} \quad (\text{condition first}).$$

Since we have three conditions (because we have three functions).

$$y_p' = V_1 e^x + 2V_2 e^{2x} + 3V_3 e^{3x} + \underbrace{V_1' e^x + V_2' e^{2x} + V_3' e^{3x}}_{=0} \quad (\text{condition second}).$$

$$y_p' = V_1 e^x + 2V_2 e^{2x} + 3V_3 e^{3x}$$

$$y_p'' = V_1 e^x + 4V_2 e^{2x} + 9V_3 e^{3x} + \underbrace{V_1' e^x + 2V_2' e^{2x} + 3V_3' e^{3x}}_{=0} \quad (\text{condition third}).$$

$$y_p'' = V_1 e^x + 4V_2 e^{2x} + 9V_3 e^{3x}$$

$$y_p''' = V_1 e^x + 8V_2 e^{2x} + 27V_3 e^{3x} + V_1' e^x + 4V_2' e^{2x} + 9V_3' e^{3x}$$

Substituting $V_1' e^x + 4V_2' e^{2x} + 9V_3' e^{3x} = e^x$

Thus we have

$$\begin{cases} W_1' e^x + V_2' e^{2x} + V_3' e^{3x} = 0 \\ V_1' e^x + 2V_2' e^{2x} + 3V_3' e^{3x} = 0 \\ V_1' e^x + 4V_2' e^{2x} + 9V_3' e^{3x} = e^x \end{cases}$$

$$W(y_1, y_2, y_3) = e^{6x} \begin{vmatrix} 1 & 1 & 1 \\ 1 & 2 & 3 \\ 1 & 4 & 9 \end{vmatrix} = 2e^{6x}$$

$$V_1' = \frac{\begin{vmatrix} 0 & e^{2x} & e^{3x} \\ 0 & 2e^{2x} & 3e^{3x} \\ e^x & 4e^{2x} & 9e^{3x} \end{vmatrix}}{2e^{6x}} = \frac{1}{2}$$

Similarly $V_2' = -e^{-x}$ $V_3' = \frac{1}{2}e^{2x}$

$$V_1 = \frac{x}{2} \quad V_2 = -e^{-x} \quad V_3 = -\frac{1}{4}e^{-2x}$$

$$y_p = \frac{x}{2}e^x + e^{-x} \cdot e^{2x} - \frac{1}{4}e^{-2x}e^{3x} = \frac{1}{2}xe^x + \frac{3}{4}e^x$$

$$y = C_1 e^x + C_2 e^{2x} + C_3 e^{3x} + \frac{1}{2}xe^x + \frac{3}{4}e^x \quad \text{or}$$

$$= C_1 e^x + C_2 e^{2x} + C_3 e^{3x} + \frac{1}{2}xe^x$$

(3) $(x^2+1) \frac{d^2y}{dx^2} - 2x \frac{dy}{dx} + 2y = 6(x^2+1)$

We solved that $y_c = C_1 x + C_2 (x^2-1)$ is the general solution of the corresponding homogeneous equation.

$$y_p = V_1 x + V_2 (x^2-1)$$

$$y_p' = V_1 + V_2 2x + V_1' x + V_2' (x^2-1)$$

By the assumption $V_1' x + V_2' (x^2-1) = 0$,

$$y_p' = V_1 + 2x V_2$$

$$y_p'' = V_1' + 2V_2 + 2x V_2'$$

By substituting,

$$(x^2+1)(V_1' + 2V_2 + 2xV_2') - 2x(V_1 + 2xV_2) + 2(V_1 x + V_2(x^2-1))$$

$$(x^2+1)V_1' + 2x(x^2+1)V_2' + (-2x+2x)V_1 + (x^2+2-4x^2+2x^2-2)V_2$$

$$= (x^2+1)V_1' + 2x(x^2+1)V_2' = 6(x^2+1)^2$$

$$x v_1' + (x^2 - 1) v_2' = 0$$

$$(x^2 + 1) v_1' + 2x(x^2 + 1) v_2' = 6(x^2 + 1)^2 \quad \text{or}$$

$$x v_1' + (x^2 - 1) v_2' = 0$$

$$v_1' + 2x v_2' = 6(x^2 + 1)$$

$$W = \begin{vmatrix} x & x^2 - 1 \\ 1 & 2x \end{vmatrix} = 2x^2 - x^2 + 1 = x^2 + 1$$

$$W_1' = \frac{\begin{vmatrix} 0 & x^2 - 1 \\ 6(x^2 + 1) & 2x \end{vmatrix}}{x^2 + 1} = -6(x^2 - 1) = -6x^2 + 6 \Rightarrow \underline{v_1' = -2x^3 + 6x}$$

$$v_2' = \frac{\begin{vmatrix} x & 0 \\ 1 & 6(x^2 + 1) \end{vmatrix}}{(x^2 + 1)} = 6x \Rightarrow \underline{v_2' = 3x^2}$$

$$y = C_1 x + C_2 (x^2 - 1) + x^4 + 3x^2$$

Theorem Let f be a particular integral of

$$a_0(x) \frac{d^4 y}{dx^4} + a_1(x) \frac{d^3 y}{dx^3} + \dots + a_{n-1}(x) \frac{dy}{dx} + a_n(x) y = f(x)$$

Let g be a particular integral of

$$a_0(x) \frac{d^4 y}{dx^4} + \dots + a_n(x) y = G(x).$$

Then $k_1 f + k_2 g$ is a particular integral of

$$a_0(x) \frac{d^4 y}{dx^4} + \dots + a_n(x) y = k_1 f + k_2 G$$

where k_1, k_2 are constants.

Example Find a particular integral of $y'' + y = 3e^x + 5 \tan x$.

We found $y_{p1} = \frac{1}{2} e^x$ is a particular solution of $y'' + y = 3e^x$ and

$y_{p2} = -(\cos x) \ln |\sec x + \tan x|$ is a particular solution of $y'' + y = \tan x$.

Thus $y_p = \frac{3}{2} e^x - 5(\cos x) \ln |\sec x + \tan x|$ is the particular integral of $y'' + y = 3e^x + 5 \tan x$.

Exercises

Find the general solution of each of the followings

1) $y'' + y = \cot x$ ($y_p = \sin x \cdot \ln|\cos x - \cot x|$) 2) $y'' + y = \tan 2x$

3) $y'' + y = \sec x$ ($y_p = \cos x \cdot \ln|\cos x| + x \sin x$) 4) $y'' + y = \sec^3 x$

5) $y'' + 4y = \sec^2 2x$

Solution: $m^2 + 4 = 0 \Rightarrow y_c = c_1 \sin 2x + c_2 \cos 2x$

$$y_p = v_1 \sin 2x + v_2 \cos 2x$$

$$y_p' = 2v_1 \cos 2x - 2v_2 \sin 2x + \frac{\sin 2x \cdot v_1' + \cos 2x \cdot v_2'}{= 0}$$

$$y_p'' = -4v_1 \sin 2x - 4v_2 \cos 2x + 2\cos 2x \cdot v_1' - 2\sin 2x \cdot v_2'$$

$$\cancel{-4v_1 \sin 2x} - \cancel{4v_2 \cos 2x} + 2\cos 2x \cdot v_1' - 2\sin 2x \cdot v_2' + \cancel{4v_1 \sin 2x} + \cancel{4v_2 \cos 2x}$$

$$= 2\cos 2x \cdot v_1' - 2\sin 2x \cdot v_2' = \sec^2 2x$$

$$\sin 2x \cdot v_1' + \cos 2x \cdot v_2' = 0$$

$$2\cos 2x \cdot v_1' - 2\sin 2x \cdot v_2' = \sec^2 2x$$

$$\Rightarrow W = \begin{vmatrix} \sin 2x & \cos 2x \\ 2\cos 2x & -2\sin 2x \end{vmatrix} = -2$$

$$v_1' = \frac{\begin{vmatrix} 0 & \cos 2x \\ \sec^2 2x & -2\sin 2x \end{vmatrix}}{-2} = \frac{\sec 2x}{-2} \Rightarrow v_1 = +\frac{1}{4} \ln|\sec x + \tan x|$$

$$v_2' = \frac{\begin{vmatrix} \sin 2x & 0 \\ 2\cos 2x & \sec^2 2x \end{vmatrix}}{-2} = \frac{\sin 2x}{-2\cos^2 2x} \Rightarrow v_2 = -\frac{1}{4} \frac{1}{\cos 2x}$$

$$y_p = \frac{\sin 2x}{4} \ln|\sec x + \tan x| - \frac{1}{4}$$

6) $y'' - 2y' + y = e^x \ln x \quad (x > 0)$

7) $y'' - 2y' + y = x e^x \ln x$

Solution

$$m^2 - 2m + 1 = (m-1)^2 = 0 \quad m = 1, 1$$

$$y_c = c_1 e^x + c_2 x e^x$$

$$y_p = v_1 e^x + v_2 x e^x$$

$$y_p' = v_1 e^x + v_2 (e^x + x e^x) + \frac{v_1' e^x + v_2' x e^x}{= 0}$$

$$y_p' = e^x v_1 + (x+1) e^x v_2$$

$$y_p'' = e^x v_1 + (x+2) e^x v_2 + e^x v_1' + (x+1) e^x v_2'$$

$$\cancel{e^x v_1' + (x+2)e^x v_2' + e^x v_1' + (x+1)e^x v_2' - 2e^x v_1 - 2(x+1)e^x v_2 + v_1 e^x + v_2 x e^x}$$

$$= e^x v_1' + (x+1)e^x v_2' = x e^x \ln x$$

$$w = \begin{vmatrix} e^x & (x+1)e^x \\ e^x & x e^x \end{vmatrix} = \underline{\underline{-e^{2x}}}$$

$$e^x v_1' + x e^x v_2' = 0$$

$$v_1' = \frac{\begin{vmatrix} x e^x \ln x & (x+1)e^x \\ 0 & x e^x \end{vmatrix}}{-e^{2x}} = -x^2 \ln x \Rightarrow v_1 = -\frac{1}{3} (x^3 \ln x - x^3)$$

$$v_2' = \frac{\begin{vmatrix} e^x & x e^x \ln x \\ e^x & 0 \end{vmatrix}}{-e^{2x}} = x \ln x \Rightarrow v_2 = \frac{1}{2} (x^2 \ln x - x^2)$$

$$y_c = -\frac{e^x}{3} (x^3 \ln x - x^3) + \frac{e^x}{2} (x^2 \ln x - x^2)$$

$$= -x^3 e^x \left(\frac{\ln x - 1}{3} - \frac{\ln x - 1}{2} \right) = \frac{x^3 e^x}{6} (\ln x - 1) \text{ Yalnız}$$

$$y_c = \frac{-5x^3 e^x}{36} + \frac{x^3 e^x \ln x}{6} \quad (\text{Doğrudur}).$$

$$9) y'' - 2y + y = (\ln x)^2$$

$$8) y'' - 2y + x = e^x \sin^{-1} x$$

$$10) y'' + 3y' + 2y = \frac{e^{-x}}{x}$$

$$11) y'' + 3y' + 2y = \frac{1}{1+e^x}$$

$$12) y'' + 3y' + 2y = \frac{1}{1+e^{2x}}$$

$$13) y'' + y = \frac{1}{1+\sin x}$$

Let us apply this theorem to an equation of the form (4.64), where the coefficients a_0, \dots, a_n are constants. Suppose that F is a simple UC function but that G is not. Then we could *not* apply the method of undetermined coefficients to find a particular integral of Equation (4.64). However, we could find a particular integral of Equation (4.62) by the method of undetermined coefficients, and then use variation of parameters to find a particular integral of Equation (4.63). Then applying Theorem 4.12, the appropriate linear combination of these two particular integrals, found by different methods, is a particular integral of Equation (4.64).

Example 4.37. Find a particular integral of

$$(4.65) \quad \frac{d^2y}{dx^2} + y = 3e^x + 5\tan x.$$

We consider the two equations

$$(4.66) \quad \frac{d^2y}{dx^2} + y = e^x$$

and

$$(4.67) \quad \frac{d^2y}{dx^2} + y = \tan x.$$

Since the function defined by e^x is a UC function, a particular integral of Equation (4.66) may be found by the method of undetermined coefficients. Letting $y_p = Ae^x$, we find at once that $A = \frac{1}{2}$; hence a particular integral of (4.66) is

$$y_p = \frac{1}{2}e^x.$$

Since the function defined by $\tan x$ is *not* a UC function, we turn to variation of parameters to find a particular integral of Equation (4.67). We have already solved this problem in Example 4.34; we found there that a particular integral of (4.67) is

$$y_p = -(\cos x) \ln |\sec x + \tan x|.$$

Thus, applying Theorem 4.12, a particular integral of Equation (4.65) is

$$y_p = \frac{3}{2}e^x - 5(\cos x) \ln |\sec x + \tan x|.$$

Exercises

Find the general solution of each of the differential equations in Exercises 1 through 13.

$$1. \quad \frac{d^2y}{dx^2} + y = \cot x.$$

$$2. \quad \frac{d^2y}{dx^2} + y = \tan^2 x.$$

$$3. \quad \frac{d^2y}{dx^2} + y = \sec x.$$

$$4. \frac{d^2y}{dx^2} + y = \sec^3 x.$$

$$\sqrt{5}. \frac{d^2y}{dx^2} + 4y = \sec^2 2x.$$

$$6. \frac{d^2y}{dx^2} - 2\frac{dy}{dx} + y = e^x \ln x, \quad (x > 0).$$

$$7. \frac{d^2y}{dx^2} - 2\frac{dy}{dx} + y = xe^x \ln x, \quad (x > 0).$$

$$8. \frac{d^2y}{dx^2} - 2\frac{dy}{dx} + y = (\ln x)^2, \quad (x > 0).$$

$$9. \frac{d^2y}{dx^2} - 2\frac{dy}{dx} + y = e^x \sin^{-1} x.$$

$$10. \frac{d^2y}{dx^2} + 3\frac{dy}{dx} + 2y = \frac{e^{-x}}{x}.$$

$$11. \frac{d^2y}{dx^2} + 3\frac{dy}{dx} + 2y = \frac{1}{1 + e^x}.$$

$$12. \frac{d^2y}{dx^2} + 3\frac{dy}{dx} + 2y = \frac{1}{1 + e^{2x}}.$$

$$13. \frac{d^2y}{dx^2} + y = \frac{1}{1 + \sin x}.$$

14. Find the general solution by two methods:

$$\frac{d^3y}{dx^3} - 3\frac{d^2y}{dx^2} - \frac{dy}{dx} + 3y = x^2 e^x.$$

15. Given that $y = x$ and $y = \frac{1}{x+1}$ are linearly independent solutions of

$$(2x+1)(x+1)\frac{d^2y}{dx^2} + 2x\frac{dy}{dx} - 2y = 0,$$

find the general solution of

$$(2x+1)(x+1)\frac{d^2y}{dx^2} + 2x\frac{dy}{dx} - 2y = (2x+1)^2.$$

16. Given that $y = \sin x$ and $y = x \sin x$ are linearly independent solutions of

$$(\sin^2 x)\frac{d^2y}{dx^2} - 2\sin x \cos x \frac{dy}{dx} + (\cos^2 x + 1)y = 0,$$

find the general solution of

$$(\sin^2 x)\frac{d^2y}{dx^2} - 2\sin x \cos x \frac{dy}{dx} + (\cos^2 x + 1)y = \sin^3 x.$$