



and Nadeau, 2021; Saranathan and Finet, 2021; Finet, 2024) and research papers on this topic highlights an active field of research. Our understanding of the spatial control of chitin assembly at the subcellular level has progressed (Pesch et al., 2017; Adler, 2019; Sviben et al., 2020; De Giorgio et al., 2023; Ghosh and Treisman, 2024; Ikeda et al., 2024; Inagaki et al., 2024; Itakura et al., 2024). Single-cell gene expression atlases of developing scales in butterflies (Prakash et al., 2024; Loh et al., 2025) and bristles in *Drosophila* (Hopkins et al., 2023) have identified gene networks involved in hair-like structure morphogenesis. Progress has been made to understand the formation of different nanostructures in butterfly wing scales such as the laminae (Thayer et al., 2020; Prakash et al., 2022; Chatterjee et al., 2023; Balakrishnan et al., 2024), the ridges (Brien et al., 2022; Ficarro et al., 2022; Lloyd et al., 2024; Totz et al., 2024), the luminal gyroid (Wilts et al., 2017), and the trabeculae (Ru et al., 2024). Work on cuticular proteins has identified their roles and spatial distributions in the development of beetle elytra (Noh et al., 2014; Murata et al., 2022; Bao et al., 2024) and butterfly scales (Liu et al., 2021).

In particular, many studies feed into the biomimetic and bioengineering fields of research. For example, the natural world has inspired multiple solutions for material adhesion (Li et al., 2024). The morphology of shark scales has led to various applications from membrane antifouling to hydrodynamics (Ghimire et al., 2024), and chameleons and cephalopods have inspired color-changing hydrogels for multiple functions including sensors (Lu et al., 2022; Zhang et al., 2023). However, beyond extracting design principles from biological materials, the ultimate goal would be to develop bio-inspired manufacturing processes, which require an in-depth understanding of the biological processes themselves.

This topic explores the genetics and cellular mechanisms underlying the development of integument and associated structures in animals and plants. With this Research Topic, we wish to direct readers to the emerging field of bio-inspired manufacturing and hope to acknowledge the need for more studies in understanding biological processes that can produce precise and compositionally driven micro- and nano-structures.

Plant cell walls are made up of cellulose microfibrils embedded in a matrix of glycoproteins, and pectic and hemicellulosic polysaccharides. Glycosylphosphatidylinositol (GPI) is a common eukaryotic lipid modification that helps proteins adhere to the membrane lipid bilayer. Zhou reviews our knowledge on GPI-anchored proteins involved in cell wall regulation in the plant model *Arabidopsis*. The author proposes that these proteins might act as structural components of the cell wall by organizing cellulose microfibrils at the cell surface.

In reptiles, a mutation in the TFEC gene leads to a piebald phenotype with white patches in the ball python while it causes reduced coloration in the brown anole lizard due to the loss of iridophores (Garcia-Elfring et al., 2023). Using comparative histology, Tzika demonstrates that TFEC mutants produce similar phenotypes of reduced coloration via different mechanisms. In the anole, TFEC is necessary for the development of iridophores, whereas in the ball python, which lacks iridophores, TFEC is important for the development of melanophores and

xanthophores. By pointing out that the same transcription factor can function differently within the same taxon, this study emphasizes that the phenotypic mutant approach is insufficient for elucidating the underlying molecular mechanisms.

Like reptiles, the ribbontail stingray exhibits structurally colored blue spots produced by dermal iridophores. These iridophores are unique by having numerous fingerlike protuberances (Surapaneni et al., 2024) and contain spherical iridosomes enclosing guanine nanocrystals. Blumer et al. provide a detailed ultrastructural description of the ribbontail ray's novel iridophore. They found that intermediate filaments form an intracellular scaffold that spaces the iridosomes within the iridophores.

In crickets and grasshoppers, males have evolved cuticular structures on their forewings to produce sound via stridulation. Turchyn and Popadic identify the POU homeodomain gene nubbin as a regulator of the development of sound resonators on the wings of the house cricket. They propose that nubbin, a key player in the wing development network, has been recruited in the course of evolution of Orthoptera to evolve these new cuticular nanostructures.

Banerjee et al. investigate the interplay between nanomorphology and pigmentation during the development of butterfly scales. They show that the loss-of-function mutations in Optix result in both pigmentation and nanomorphology defects. By comparing these effects with mutants in melanin and/or ommochrome pathways, they propose that Optix regulates nanomorphology via its effects on pigmentation, complementing earlier studies on melanized (Matsuoka and Monteiro, 2018) and silver scales (Prakash et al., 2022).

## Author contributions

CF: Writing  $\frac{1}{2}$ original draft, Writing  $\frac{1}{2}$ review & editing. AP: Writing  $\frac{1}{2}$ original draft, Writing  $\frac{1}{2}$ review & editing.

## Conflict of interest

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

## Generative AI statement

The author(s) declare that no Generative AI was used in the creation of this manuscript.

## Publisher's note

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

## References

- Adler, P. N. (2019). The localization of chitin synthase mediates the patterned deposition of chitin in developing *Drosophila* bristles. *bioRxiv* [Preprint]. doi: 10.1101/718841
- Airolidi, C. A., Ferria, J., and Glover, B. J. (2019). The cellular and genetic basis of structural colour in plants. *Curr. Opin. Plant Biol.* 47, 81–87. doi: 10.1016/j.cpb.2018.10.002
- Akat, E., Yenmis, M., Pombal, M. A., Molist, P., Megias, M., Arman, S., et al. (2022). Comparison of vertebrate skin structure at class level: A review. *Anat. Rec.* 305, 3543–3608. doi: 10.1002/ar.24908
- Balakrishnan, D., Prakash, A., Daurer, B. J., Finet, C., Chen Lim, Y., Shen, Z., et al. (2024). Nanoscale cuticle mass density variations in uenced by pigmentation in butterfly wing scales. *arXiv* [Preprint]. doi: 10.48550/arXiv.2305.16628
- Bao, H., Liu, Y., Duan, Y., Chen, L., and Yang, Q. (2024). The beetle's structural protein CPCFC making elytra tough and rigid. *Insect Sci* [Preprint]. doi: 10.1111/1744-7917.13443. Online ahead of print.
- Barthlott, W., Mail, M., and Neinhuis, C. (2016). Superhydrophobic hierarchically structured surfaces in biology: Evolution, structural principles and biomimetic applications. *Philos. Trans. R. Soc. A Math. Phys. Eng. Sci.* 374, 20160191. doi: 10.1098/rsta.2016.0191
- Brien, M. N., Enciso-Romero, J., Lloyd, V. J., Curran, E. V., Parnell, A. J., Moroch, C., et al. (2022). The genetic basis of structural colour variation in mimetic *Heliconius* butterflies. *Philos. Trans. R. Soc. B Biol. Sci.* 377, 20200505. doi: 10.1098/rstb.2020.0505
- Carroll, S., Amsbury, S., Durney, C. H., Smith, R. S., Morris, R. J., Gray, J. E., et al. (2022). Altering arabinans increases *Arabidopsis* guard cell exibility and stomatal opening. *Curr. Biol.* 32, 3170–3179.e4. doi: 10.1016/j.cub.2022.05.042
- Chatterjee, M., Siegel, K. J., Loh, L. S., McDonald, J. M. C., and Reed, R. D. (2023). Butterfly wing iridescence is regulated by araucan, a direct target of optix and spalt. *bioRxiv* [Preprint]. doi: 10.1101/2023.11.21.568172
- Cosgrove, D. J. (2024). Structure and growth of plant cell walls. *Nat. Rev. Mol. Cell Biol.* 25, 340–358. doi: 10.1038/s41580-023-00691-y
- De Giorgio, E., Giannios, P., Espinosa, M. L., and Llimargas, M. (2023). A dynamic interplay between chitin synthase and the proteins Expansion/Rebuc reveals that chitin polymerisation and translocation are uncoupled in *Drosophila*. *PLoS Biol.* 21, e3001978. doi: 10.1371/journal.pbio.3001978
- Ficarrotta, V., Hanly, J. J., Loh, L. S., Francescutti, C. M., Ren, A., Tunstir, K., et al. (2022). A genetic switch for male UV iridescence in an incipient species pair of sulphur butterflies. *Proc. Natl. Acad. Sci. U. S. A.* 119, e2109255118. doi: 10.1073/pnas.2109255118
- Finet, C. (2024). Developmental genetics of cuticular micro- and nano-structures in insects. *Curr. Opin. Insect Sci.* 65, 101254. doi: 10.1016/j.cois.2024.101254
- Finet, C., Decaras, A., Armisen, D., and Khila, A. (2018). The achaete-scute complex contains a single gene that controls bristle development in the semi-aquatic bugs. *Proc. R. Soc. B Biol. Sci.* 285, 20182387. doi: 10.1098/rspb.2018.2387
- Finet, C., Decaras, A., Rutkowska, M., Roux, P., Collaudin, S., Joncour, P., et al. (2022). Leg length and bristle density, both necessary for water surface locomotion, are genetically correlated in water striders. *Proc. Natl. Acad. Sci. U. S. A.* 119, e2119210119. doi: 10.1073/pnas.2119210119
- Finet, C., Ruan, Q., Bei, Y. Y., You En Chan, J., Saranathan, V., Yang, J. K. W., et al. (2023). Multi-scale dissection of wing transparency in the clearwing butterfly *Phanus vitreus*. *J. R. Soc. Interface* 20, 20230135. doi: 10.1098/rsif.2023.0135
- Garcia-Elfring, A., Sabin, C. E., Iouchmanov, A. L., Roffey, H. L., Samudra, S. P., Alcalá, A. J., et al. (2023). Piebaldism and chromatophore development in reptiles are linked to the tfec gene. *Curr. Biol.* 33, 755–763.e3. doi: 10.1016/j.cub.2023.01.004
- Ghimire, A., Dahl, R. B., Shen, S. F., and Chen, P. Y. (2024). Shark skin denticles: from morphological diversity to multi-functional adaptations and applications. *Adv. Funct. Mater.* 34, 2307121. doi: 10.1002/adfm.202307121
- Ghosh, N., and Treisman, J. E. (2024). Apical cell expansion maintained by Dusky-like establishes a scaffold for corneal lens morphogenesis. *Sci. Adv.* 10, ead04167. doi: 10.1126/sciadv.ad04167
- Gow, N. A. R., and Lenardon, M. D. (2023). Architecture of the dynamic fungal cell wall. *Nat. Rev. Microbiol.* 21, 248–259. doi: 10.1038/s41579-022-00796-9
- Hopkins, B. R., Barmina, O., and Kopp, A. (2023). A single-cell atlas of the sexually dimorphic *Drosophila* foreleg and its sensory organs during development. *PLoS Biol.* 21, e3002148. doi: 10.1371/journal.pbio.3002148
- Hsiung, B. K., Shawkey, M. D., and Blackledge, T. A. (2019). Color production mechanisms in spiders. *J. Arachnol.* 47, 165–180. doi: 10.1636/JoA-S-18-022
- Ikeda, K. N., Belevich, I., Zelaya-Lainez, L., Orel, L., Fussl, J., Gumulec, J., et al. (2024). Dynamic microvilli sculpt bristles at nanometric scale. *Nat. Commun.* 15, 3733. doi: 10.1038/s41467-024-48044-3
- Inagaki, S., Wada, H., Itabashi, T., Itakura, Y., Nakagawa, R., Chen, L., et al. (2024). Endoplasmic reticulum patterns insect cuticle nanostructure. *bioRxiv* [Preprint]. doi: 10.1101/2024.08.20.608717
- Itakura, Y., Wada, H., Inagaki, S., and Hayashi, S. (2024). Mechanical control of the insect extracellular matrix nanostructure. *bioRxiv* [Preprint]. doi: 10.1101/2024.08.20.608778
- Li, W., Zhou, R., Ouyang, Y., Guan, Q., Shen, Y., Saiz, E., et al. (2024). Harnessing biomimicry for controlled adhesion on material surfaces. *Small* 20, 2401859. doi: 10.1002/sml.202401859
- Liu, J., Chen, Z., Xiao, Y., Asano, T., Li, S., Peng, L., et al. (2021). Lepidopteran wing scales contain abundant cross-linked lm-forming histidine-rich cuticular proteins. *Commun. Biol.* 4, 491. doi: 10.1038/s42003-021-01996-4
- Lloyd, V. J., Burg, S. L., Harizanov, J., Garcia, E., Hill, O., Enciso-Romero, J., et al. (2024). The actin cytoskeleton plays multiple roles in structural color formation in butterfly wing scales. *Nat. Commun.* 15, 4073. doi: 10.1038/s41467-024-48060-3
- Lloyd, V. J., and Nadeau, N. J. (2021). The evolution of structural colour in butterflies. *Curr. Opin. Genet. Dev.* 69, 28–34. doi: 10.1016/j.gde.2021.01.004
- Loh, L. S., DeMarr, K. A., Tsimba, M., Heryanto, C., Berrio, A., Patel, N. H., et al. (2025). Lepidopteran scale cells derive from sensory organ precursors through a canonical lineage. *Development* 152, DEV204501. doi: 10.1242/dev.204501
- Lu, W., Si, M., Le, X., and Chen, T. (2022). Mimicking color-changing organisms to enable the multicolors and multifunctions of smart fluorescent polymeric hydrogels. *Acc. Chem. Res.* 55, 2291–2303. doi: 10.1021/acs.accounts.2c00320
- Matsuoka, Y., and Monteiro, A. (2018). Melanin pathway genes regulate color and morphology of butterfly wing scales. *Cell Rep.* 24, 56–65. doi: 10.1016/j.celrep.2018.05.092
- McPhedran, R. C., Nicorovici, N. A., McKenzie, D. R., Botten, L. C., Parker, A. R., and Rouse, G. W. (2001). The sea mouse and the photonic crystal. *Aust. J. Chem.* 54, 241–244. doi: 10.1071/CH01054
- Middleton, R., Sinnott-Armstrong, M., Ogawa, Y., Jacucci, G., Moyroud, E., Rudall, P. J., et al. (2020). *Viburnum tinus* fruits use lipids to produce metallic blue structural color. *Curr. Biol.* 30, 3804–3810.e2. doi: 10.1016/j.cub.2020.07.005
- Miserez, A., Schneberk, T., Sun, C., Zok, F. W., and Waite, J. H. (2008). The transition from stiff to compliant materials in squid beaks. *Science* 319, 1816–1819. doi: 10.1126/science.1154117
- Murata, S., Rivera, J., Noh, M. Y., Hiyoshi, N., Yang, W., Parkinson, D. Y., et al. (2022). Unveiling characteristic proteins for the structural development of beetle elytra. *Acta Biomater.* 140, 467–480. doi: 10.1016/j.actbio.2021.12.021
- Muthukrishnan, S., Arakane, Y., Noh, M. Y., Mun, S., Merzendorfer, H., Boehringer, C., et al. (2022). Chitin in insect cuticle. *Adv. Insect Physiol.* 62, 1–110. doi: 10.1016/b.s.aip.2022.03.001
- Noh, M. Y., Kramer, K. J., Muthukrishnan, S., Kanost, M. R., Beeman, R. W., and Arakane, Y. (2014). Two major cuticular proteins are required for assembly of horizontal laminae and vertical pore canals in rigid cuticle of *Tribolium castaneum*. *Insect Biochem. Mol. Biol.* 53, 22–29. doi: 10.1016/j.ibmb.2014.07.005
- Pasquina-Lemonche, L., Burns, J., Turner, R. D., Kumar, S., Tank, R., Mullin, N., et al. (2020). The architecture of the Gram-positive bacterial cell wall. *Nature* 582, 294–297. doi: 10.1038/s41586-020-2236-6
- Pesch, Y. Y., Riedel, D., and Behr, M. (2017). *Drosophila* Chitinase 2 is expressed in chitin producing organs for cuticle formation. *Arthropod Struct. Dev.* 46, 4–12. doi: 10.1016/j.asd.2016.11.002
- Prakash, A., Dion, E., Banerjee, T. D., and Monteiro, A. (2024). The molecular basis of scale development highlighted by a single-cell atlas of *Bicyclus anynana* butterfly pupal forewings. *Cell Rep.* 43, 114147. doi: 10.1016/j.celrep.2024.114147
- Prakash, A., Finet, C., Banerjee, T. D., Saranathan, V., and Monteiro, A. (2022). Antennapedia and optix regulate metallic silver wing scale development and cell shape in *Bicyclus anynana* butterflies. *Cell Rep.* 40, 111052. doi: 10.1016/j.celrep.2022.111052
- Ru, H., Finet, C., and Monteiro, A. (2024). Calreticulin is required for cuticle deposition and trabeculae formation inside butterfly wing scale cells. *bioRxiv* [Preprint]. doi: 10.1101/2024.09.12.612608
- Saranathan, V., and Finet, C. (2021). Cellular and developmental basis of avian structural coloration. *Curr. Opin. Genet. Dev.* 69, 56–64. doi: 10.1016/j.gde.2021.02.004
- Seago, A. E., Brady, P., Vigneron, J. P., and Schultz, T. D. (2009). Gold bugs and beyond: A review of iridescence and structural colour mechanisms in beetles (Coleoptera). *J. R. Soc. Interface* 6, S165–S184. doi: 10.1098/rsif.2008.0354.focus
- Seale, M., Cummins, C., Viola, I. M., Mastropalo, E., and Nakayama, N. (2018). Design principles of hair-like structures as biological machines. *J. R. Soc. Interface* 15, 20180206. doi: 10.1098/rsif.2018.0206
- Sun, J., and Bhushan, B. (2012). Hierarchical structure and mechanical properties of nacre: A review. *RSC Adv.* 2, 7617–7632. doi: 10.1039/c2ra20218b
- Surapaneni, V. A., Blumer, M. J., Tadayon, K., McIvor, A. J., Redl, S., Honis, H. R., et al. (2024). Ribbontail stingray skin employs a core-shell photonic glass ultrastructure to make blue structural color. *Adv. Opt. Mater.* 12, 2301909. doi: 10.1002/adom.202301909
- Sviben, S., Spaeker, O., Bennet, M., Alberic, M., Alberic, M., Dirks, J. H., et al. (2020). Epidermal cell surface structure and chitin-protein co-assembly determine fiber architecture in the locust cuticle. *ACS Appl. Mater. Interfaces* 12, 25581–25590. doi: 10.1021/acsami.0c04572

Thayer, R., Allen, F., and Patel, N. H. (2020). Structural color in *Junonia* butterflies evolves by tuning scale lamina thickness. *Elife* 9, e52187. doi: 10.7554/eLife.52187

Thayer, R., and Patel, N. H. (2023). A meta-analysis of butterfly structural colors: their color range, distribution and biological production. *J. Exp. Biol.* 226, jeb245940. doi: 10.1242/jeb.245940

Totz, J. F., McDougal, A. D., Wagner, L., Kang, S., So, P. T. C., Dunkel, J., et al. (2024). Cell membrane buckling governs early-stage ridge formation in butterfly wing scales. *Cell Rep. Phys. Sci.* 5, 102063. doi: 10.1016/j.xcrp.2024.102063

Watson, G. S., Watson, J. A., and Cribb, B. W. (2017). Diversity of cuticular micro- and nanostructures on insects: properties, functions, and potential applications. *Annu. Rev. Entomol.* 62, 185–205. doi: 10.1146/annurev-ento-031616-035020

Wilts, B. D., Zubiri, B. A., Klatt, M. A., Butz, B., Fischer, M. G., Kelly, S. T., et al. (2017). Butterfly gyroid nanostructures as a time-frozen glimpse of intracellular membrane development. *Sci. Adv.* 3, e1603119. doi: 10.1126/sciadv.1603119

Zhang, W., Tian, H., Liu, T., Liu, H., Zhao, F., Li, X., et al. (2023). Chameleon-inspired active tunable structural color based on smart skin with multi-functions of structural color, sensing and actuation. *Mater. Horizons* 10, 2024–2034. doi: 10.1039/d3mh00070b