

# On the Effects of Interference in BSM Production and Detection of diTaus at the LHC

Cristian Fernando Rodríguez Cruz

Authors:

A. Flórez<sup>1</sup>, **C. Rodríguez**<sup>1</sup>, J. Reyes-Vega<sup>1</sup>,  
J. Jones-Pérez<sup>2</sup>.

<sup>1</sup>Universidad de los Andes

<sup>2</sup>Pontificia Universidad Católica del Perú

November 3, 2024

# Outline

## 1 Introduction

- Motivation
- BSM Signatures
- Interference Phenomena in the SM

## 2 Example: The 4321-Model


- The model
- Sensitivity Reach of the  $U_1$  Leptoquark
- Interference with a  $Z'$  vector boson

## 3 Final Remarks

- Conclusions
- Future Work


# The Standard Model of Particle Physics

Weak bosons mix the different generations of quarks via the CKM matrix, but this does not happen for leptons. This property of the model is known as **lepton flavor universality (LFU)**.




# The Standard Model of Particle Physics



Weak bosons mix the different generations of quarks via the CKM matrix, but this does not happen for leptons. This property of the model is known as **lepton flavor universality (LFU)**.





However, recent measurements of the  $R(D)$  and  $R(D^*)$  ratios show a deviation from the SM predictions. This could be a hint of **lepton flavor violation (LFV)** and then **new physics beyond the SM**.



# Large Hadron Collider



# Large Hadron Collider





- Feasibility Studies is needed.
- Take Care on the dependence on the different parameters.
- Take care on the content of particles.
- Take care of the signal composition.
- Take care on interference effects.



# BSM Signatures on the Di-Tau Channel at the LHC

LFV and  $\tau$  lepton as window to new physics

In the different models, we can have different production mechanisms.  
For example, resonant ones



or non-resonant ones




Different Models could have one or several contributions to the di-tau channel, and the **interference between them could be relevant.**

# Interference Phenomena in the SM

Photon and Z-boson interference,  $q\bar{q} \longrightarrow \tau^+\tau^-$

The squared matrix element can be written as

$$\begin{aligned} |\mathcal{M}|^2 &= |\mathcal{M}_{\gamma^*} + \mathcal{M}_Z|^2 \\ &= |\mathcal{M}_{\gamma^*}|^2 + |\mathcal{M}_Z|^2 + 2 \operatorname{Re} (\mathcal{M}_{\gamma^*}^* \mathcal{M}_Z) . \end{aligned}$$






# Interference Phenomena in the SM

Photon and Z-boson interference,  $q\bar{q} \longrightarrow \tau^+\tau^-$


The squared matrix element can be written as

$$\begin{aligned} |\mathcal{M}|^2 &= |\mathcal{M}_{\gamma^*} + \mathcal{M}_Z|^2 \\ &= |\mathcal{M}_{\gamma^*}|^2 + |\mathcal{M}_Z|^2 + 2 \operatorname{Re} (\mathcal{M}_{\gamma^*}^* \mathcal{M}_Z) . \end{aligned}$$



For the case  $q_R \bar{q}_L \longrightarrow \tau_L^+ \tau_R^-$ , the amplitudes are

$$\begin{aligned} |\mathcal{M}_{\gamma^*}|^2 &= e^4 \left[ Q^{(f)} Q^{(q)} \right]^2 [1 + \cos \theta]^2 \\ |\mathcal{M}_Z|^2 &= \frac{s^2 g_Z^4 \left[ g_R^{(f)} g_R^{(q)} \right]^2}{(s - m_Z^2)^2 + (m_Z \Gamma_Z)^2} [1 + \cos \theta]^2 \\ \mathcal{M}_{\gamma^*}^* \mathcal{M}_Z &= \frac{g_Z^2 e^2 Q^{(f)} Q^{(q)} g_R^{(f)} g_R^{(q)}}{(s - m_Z^2 + i \Gamma_Z m_Z)} s (1 + \cos \theta)^2 \end{aligned}$$




# Interference Phenomena in the SM

Photon and Z-boson interference,  $q\bar{q} \longrightarrow \tau^+\tau^-$


The squared matrix element can be written as

$$\begin{aligned} |\mathcal{M}|^2 &= |\mathcal{M}_{\gamma^*} + \mathcal{M}_Z|^2 \\ &= |\mathcal{M}_{\gamma^*}|^2 + |\mathcal{M}_Z|^2 + 2 \operatorname{Re} (\mathcal{M}_{\gamma^*}^* \mathcal{M}_Z) . \end{aligned}$$



For the case  $q_R \bar{q}_L \longrightarrow \tau_L^+ \tau_R^-$ , the amplitudes are


$$\begin{aligned} |\mathcal{M}_{\gamma^*}|^2 &= e^4 \left[ Q^{(f)} Q^{(q)} \right]^2 [1 + \cos \theta]^2 \\ |\mathcal{M}_Z|^2 &= \frac{s^2 g_Z^4 \left[ g_R^{(f)} g_R^{(q)} \right]^2}{(s - m_Z^2)^2 + (m_Z \Gamma_Z)^2} [1 + \cos \theta]^2 \\ \mathcal{M}_{\gamma^*}^* \mathcal{M}_Z &= \frac{g_Z^2 e^2 Q^{(f)} Q^{(q)} g_R^{(f)} g_R^{(q)}}{(s - m_Z^2 + i \Gamma_Z m_Z)} s (1 + \cos \theta)^2 \end{aligned}$$



Always that you have two or more contributions to a process, the interference between near to the resonances could be relevant.

# Example: The Vector Leptoquark Model

A leptoquark is defined as a particle with a vertex that mix vectors and quarks.



If  $U_1$  is a vector leptoquark that preserves the chirality on the vertex, we expect an interaction term like

$$\sim U_1^\mu \bar{q}_L \gamma_\mu \ell_L,$$

and these allows a similar interaction term for the right handed currents

$$\sim U_1^\mu \bar{d}_R \gamma_\mu e_R.$$

Where the SM charges for the leptoquark, in the  $Y = 2(Q - T_3)$  convention, are

	$\bar{q}_L$	$\ell_L^i$	$\bar{q}_L \gamma_\mu \ell_L$	$U_1^\mu$
$U(1)$	$-1/3$	$-1$	$-4/3$	$+4/3$
$SU(2)$	$\mathbf{\bar{2}}$	$\mathbf{2}$	$\mathbf{1}$	$\mathbf{1}$
$SU(3)$	$\mathbf{\bar{3}}$	$\mathbf{1}$	$\mathbf{\bar{3}}$	$\mathbf{3}$

Then, the leptoquark  $U_1 \sim (\mathbf{3}_C, \mathbf{1}_I, 4/3_Y)$ , and its covariant derivative is

$$\mathcal{D}_\mu U_\nu = \left( \partial_\mu + ig_s T^a G_\mu^a + i \frac{2}{3} g' B_\mu \right) U_\nu.$$

# The Vector Leptoquark Lagrangian

The full Lagrangian for the vector leptoquark is

$$\begin{aligned}\mathcal{L}_U = & -\frac{1}{2}U_{\mu\nu}^\dagger U^{\mu\nu} + M_U^2 U_\mu^\dagger U^\mu \\ & -ig_s(1-\kappa_c)U_\mu^\dagger T^a U_\nu G_a^{\mu\nu} - \frac{2i}{3}g'(1-\kappa_Y)U_\mu^\dagger U_\nu B^{\mu\nu} \\ & + \frac{g_U}{\sqrt{2}} \left[ U_1^\mu \left( \beta_L^{ij} \bar{q}_L^i \gamma_\mu e_L^j + \beta_R^{ij} \bar{d}_R^i \gamma_\mu e_R^j \right) + \text{h.c.} \right]\end{aligned}$$

where  $U_{\mu\nu} = \mathcal{D}_\mu U_\nu - \mathcal{D}_\nu U_\mu$ ,  $\mathcal{D}_\mu = \partial_\mu - ig_s G_\mu^a T^a - i\frac{2}{3}g_Y B_\mu$ , and the couplings  $\beta_L$  and  $\beta_R$  are complex  $3 \times 3$  matrices in flavor space.

# The Vector Leptoquark Lagrangian

The full Lagrangian for the vector leptoquark is

$$\begin{aligned}\mathcal{L}_U = & -\frac{1}{2}U_{\mu\nu}^\dagger U^{\mu\nu} + M_U^2 U_\mu^\dagger U^\mu \\ & -ig_s(1-\kappa_c)U_\mu^\dagger T^a U_\nu G_a^{\mu\nu} - \frac{2i}{3}g'(1-\kappa_Y)U_\mu^\dagger U_\nu B^{\mu\nu} \\ & + \frac{g_U}{\sqrt{2}} \left[ U_1^\mu \left( \beta_L^{ij} \bar{q}_L^i \gamma_\mu e_L^j + \beta_R^{ij} \bar{d}_R^i \gamma_\mu e_R^j \right) + \text{h.c.} \right]\end{aligned}$$


where  $U_{\mu\nu} = \mathcal{D}_\mu U_\nu - \mathcal{D}_\nu U_\mu$ ,  $\mathcal{D}_\mu = \partial_\mu - ig_s G_\mu^a T^a - i\frac{2}{3}g_Y B_\mu$ , and the couplings  $\beta_L$  and  $\beta_R$  are complex  $3 \times 3$  matrices in flavor space.

The  $\Delta F = 2$  and lepton flavor violating processes indicates an structure as

$$\beta_L = \begin{pmatrix} 0 & 0 & \beta_L^{13} \\ 0 & 0 & \beta_L^{23} \\ 0 & \beta_L^{32} & \beta_L^{33} \end{pmatrix}, \quad \beta_R = \text{diag}(0, 0, \beta_R^{33})$$



If  $U_1$  has a gauge origin  $\kappa_c = \kappa_Y = 0$ . We choose  $U(2)$  in quark and leptons space, in a way that you have an hierarchy,  $|\beta_L^{31}| \ll |\beta_L^{23}|, |\beta_L^{32}| \ll |\beta_R^{33}|, |\beta_L^{33}| = \mathcal{O}(1)$ .

# Feasibility Studies Workflow



$$S_T^{\text{meT}} = \text{met} + \sum_i |p_T^i|$$

# Sensitivity Reach of the $U_1$ Leptoquark



$$S_T^{\text{meT}} = \text{met} + \sum_i |p_T^i|$$

Non-resonant production is highly dependent on the couplings, so it dominates the regions of high coupling constants at all masses.



# Take care, you could need a $Z'$ boson

The generator  $T_{B-L}$  is associated with the  $U(1)_{B-L}$  symmetry with a  $Z'$  boson. The interaction terms for the  $Z'$  boson have the form

$$\begin{aligned}\mathcal{L}_{\text{int}} &\sim Z'_\mu \left( \bar{\psi}_L^{\text{SM}} \gamma^\mu (3T_{B-L}) \psi_L^{\text{SM}} \right) \\ &\sim Z'_\mu \left( \bar{q}_L \gamma^\mu q_L - 3\bar{\ell}_L \gamma^\mu \ell_L \right) .\end{aligned}$$

# Take care, you could need a $Z'$ boson

The generator  $T_{B-L}$  is associated with the  $U(1)_{B-L}$  symmetry with a  $Z'$  boson. The interaction terms for the  $Z'$  boson have the form

$$\begin{aligned}\mathcal{L}_{\text{int}} &\sim Z'_\mu \left( \bar{\psi}_L^{\text{SM}} \gamma^\mu (3T_{B-L}) \psi_L^{\text{SM}} \right) \\ &\sim Z'_\mu \left( \bar{q}_L \gamma^\mu q_L - 3\bar{\ell}_L \gamma^\mu \ell_L \right).\end{aligned}$$

so, the full Lagrangian for the  $Z'$  boson is

$$\begin{aligned}\mathcal{L}_{Z'} &= -\frac{1}{4}Z'_{\mu\nu}Z'^{\mu\nu} + \frac{1}{2}M_{Z'}^2 Z'_\mu Z'^\mu \\ &\quad + \frac{g_{Z'}}{2\sqrt{6}} Z'^\mu \left( \zeta_q^{ij} \bar{q}_L^i \gamma_\mu q_L^j + \zeta_u^{ij} \bar{u}_R^i \gamma_\mu u_R^j + \zeta_d^{ij} \bar{d}_R^i \gamma_\mu d_R^j - 3\zeta_\ell^{ij} \bar{\ell}_L^i \gamma_\mu \ell_L^j - 3\zeta_e^{ij} \bar{e}_R^i \gamma_\mu e_R^j \right),\end{aligned}\tag{1}$$

where the couplings  $\zeta$  are complex  $3 \times 3$  matrices in flavor space.

# Take care, you could need a $Z'$ boson

The generator  $T_{B-L}$  is associated with the  $U(1)_{B-L}$  symmetry with a  $Z'$  boson. The interaction terms for the  $Z'$  boson have the form

$$\begin{aligned}\mathcal{L}_{\text{int}} &\sim Z'_\mu \left( \bar{\psi}_L^{\text{SM}} \gamma^\mu (3T_{B-L}) \psi_L^{\text{SM}} \right) \\ &\sim Z'_\mu \left( \bar{q}_L \gamma^\mu q_L - 3\bar{\ell}_L \gamma^\mu \ell_L \right).\end{aligned}$$

so, the full Lagrangian for the  $Z'$  boson is


$$\begin{aligned}\mathcal{L}_{Z'} &= -\frac{1}{4}Z'_{\mu\nu}Z'^{\mu\nu} + \frac{1}{2}M_{Z'}^2 Z'_\mu Z'^\mu \\ &\quad + \frac{g_{Z'}}{2\sqrt{6}} Z'^\mu \left( \zeta_q^{ij} \bar{q}_L^i \gamma_\mu q_L^j + \zeta_u^{ij} \bar{u}_R^i \gamma_\mu u_R^j + \zeta_d^{ij} \bar{d}_R^i \gamma_\mu d_R^j - 3\zeta_\ell^{ij} \bar{\ell}_L^i \gamma_\mu \ell_L^j - 3\zeta_e^{ij} \bar{e}_R^i \gamma_\mu e_R^j \right),\end{aligned}\tag{1}$$

where the couplings  $\zeta$  are complex  $3 \times 3$  matrices in flavor space.


We assume that both, the  $Z'$  and the vector leptoquark  $U_1$ , have preferential couplings to third generation fermions, so  $\beta^{33} \gg \beta^{ij}$  and  $\zeta^{33} \gg \zeta^{ij}$ .

# Interference with a $Z'$ vector boson

Non-Resonant Production (leptoquarks)    Resonant Production (neutral bosons)



$$\mathcal{M}_{U_1} \sim \frac{1}{t - m_{U_1}^2 + im_{U_1}\Gamma_{U_1}}, \quad (2)$$



$$\mathcal{M}_{Z'} \sim \frac{1}{s - m_{Z'}^2 + im_{Z'}\Gamma_{Z'}}, \quad (3)$$


so

the interference has the form

$$\sim \frac{m_{LQ}m_{Z'}\Gamma_{LQ}\Gamma_{Z'} - (t - m_{LQ}^2)(s - m_{Z'}^2)}{\left[(t - m_{LQ}^2)^2 + m_{LQ}^2\Gamma_{LQ}^2\right] \left[(s - m_{Z'}^2)^2 + m_{Z'}^2\Gamma_{Z'}^2\right]}.$$

# Interference with a $Z'$ vector boson

$$\frac{d}{dm} \left[ \sigma_{LQ+Z'} - (\sigma_{LQ} + \sigma_{Z'}) \right] \sim \frac{g_{Z'} g_U}{s} \frac{m_{LQ} m_{Z'} \Gamma_{LQ} \Gamma_{Z'} - (t - m_{LQ}^2)(s - m_{Z'}^2)}{\left[ (t - m_{LQ}^2)^2 + m_{LQ}^2 \Gamma_{LQ}^2 \right] \left[ (s - m_{Z'}^2)^2 + m_{Z'}^2 \Gamma_{Z'}^2 \right]}.$$



# Conclusions

# Future Work