

T1DES in SIDM: A New Class of SIDM Subhalo Simulations

Charles Gannon,^{*} Anna Nierenberg, and Ryan Keeley
University of California, Merced, 5200 N Lake Road, Merced, CA, 95341, USA[†]

Andrew Benson
Carnegie Observatories, 813 Santa Barbara Street, Pasadena, CA 91101, USA

Xiaolong Du
Department of Physics and Astronomy, University of California, Los Angeles, CA 90095, USA

Daniel Gilman
Department of Astronomy & Astrophysics, University of Chicago, Chicago, IL 60637 USA

Ethan Nadler
Department of Astronomy & Astrophysics, University of California, San Diego, La Jolla, CA 92093, USA

Andrew Robertson
*The Observatories of the Carnegie Institution for Science,
813 Santa Barbara Street, Pasadena, 91101, CA, USA
(Dated: September 15, 2026)*

Dark matter models with self interactions have gained to resolve apparent shortcomings of the Λ CDM cosmology. Testing these models of Self Interacting Dark Matter (SIDM) requires accurate modeling the gravothermal evolution of halos under accelerated heat transport facilitated by these interactions. The picture becomes more complicated for subhalos (halos evolving within the tidal field of the host), as they have their gravothermal evolution altered by tidal physics in addition to any potential dark matter self interaction; many probes of SIDM such as gravitational lensing sample from a halo population that includes subhalos. Existing methods of simulating SIDM subhalos are very computationally expensive or make restrictive assumptions on tidal physics. To address these shortcomings, we present T1DES (Tidal 1D Evolution of SIDM subhalos), a new class of 1D SIDM subhalo simulations incorporating realistic tidal physics while maintaining the many orders of magnitude speed up over full 3D SIDM simulations. We find that under no evaporation the core collapse is accelerated, with nearly 4 times as many subhalos collapsing compared to isolated counterparts.

I. INTRODUCTION

Cold Dark Matter or CDM has had a long string of successes explaining the largest scales of cosmology, from the cosmic microwave background Planck Collaboration [1] to the spatial and mass function distributions of galaxies [2–5]. However, recent inferences on the sub-galactic from direct gravitational imaging the potential of lenses such as the jackpot lens (SDSSJ0946+1006) [6, 7], B1938+666 and J0946+1006 [8] have provided evidence for anomalously dense (high concentration) dark matter halos compared to CDM predictions. Although these results have been hotly debated and solutions within the CDM framework have been proposed [9], the intriguing possibility remains that these results hint at physics beyond CDM. A class of alternate dark matter models with non-negligible self-interaction on the sub-galactic scales, called self-interacting dark matter (SIDM) has

been proposed as an explanation of anomalously dense halos inferred in these systems [10]. In these models, dark matter self-interaction facilitates the conduction of heat within the halo; due to the negative heat capacity of self-gravitating systems, runaway heat transfer can occur leading to the halo gravothermally collapsing, leading to extremely dense collapsed remnants at the end of their evolution (see Refs. [11, 12] for reviews).

In this paper we focus on studying core collapse rates of SIDM subhalos; lensing and stellar streams results sample from a population that includes subhalos. To infer SIDM constraints, it is important to understand how fast gravothermal collapse may occur given an SIDM model; tidal forces from the host are known to alter core collapse rates [13–17]. Previously there have been 2 main methods to study SIDM subhalos: 3-D N-body simulations (both cosmological [18–25] and idealized [10, 13, 26–28]) and gravothermal fluid modeling [14, 15]; while these have all been fruitful ways of studying SIDM substructure in their own right, they each have significant drawbacks. 3-D N-body simulations, which are traditionally the primary method of studying dark matter halos are computationally expensive especially in the core collapse regime;

^{*} University of California, Merced, 5200 N Lake Road, Merced, CA, 95341, USA; cgannon@ucmerced.edu

[†] annanierenberg@ucmerced.edu

tidal mass loss amplifies this challenge by requiring a larger number of particles to resolve the halo initially, often requiring tens of thousands of core hours to properly resolve core collapse even in idealized simulations. Gravothermal fluid models, while much computationally cheaper than 3-D N-body simulations use restrictive assumptions on tidal physics namely approximating tidal stripping as one or more truncations event where mass is abruptly removed from the halo. Because of these limitations, accurately predicting the core collapse rates for large samples of halos across different SIDM models remains a challenge.

In this work, we develop **T1DES** (Tidal 1D Evolution of Subhalos): a pipeline to simulate the evolution of idealized SIDM subhalos in 1-D with realistic tidal evolution. By simulating subhalos evolution in 1-D, we are able to simulate SIDM physics much more computationally efficiently than the 3D case, resulting in over an order of magnitude speed up in some case enabling previously infeasible exploration of SIDM model space. We begin by explaining our methods in Section II. First, we discuss generating a realization of **galacticus** [29] CDM halos in Section II A to set the initial conditions for our 1-D simulations and provide a target mass loss histories. Next, in Section II B we discuss how we modify the dynamics of the 1-D simulation code **NSphere** Kamionkowski *et al.* [30], Kamionkowski and Sigurdson [31, 32] to include an effective tidal field calculated from a target mass loss history. Then in Section II C we discuss the SIDM physics implemented in **NSphere** along with the several modifications we make for our use case including a dynamic timestep and core collapse termination condition. We then discuss the results of our simulations in Section III and provide a summary and concluding remarks in Section IV.

II. METHODS

We aim to explore the population level statistics expected of subhalos systems hosted by strong gravitational lenses. To accomplish this we develop the **T1DES** pipeline to simulate realizations of SIDM subhalos across various tidal fields and under the influence of a tidal field. We use a modified version of the 1-D N-body **NSphere** code, this provides a large speed up over traditional 3-D N-body simulations; the slowest halos in our pipeline complete in ~ 600 core hours while a comparable idealized 3-D SIDM simulation can take $> 10,000$ core hours to complete. In this section, we give an overview of the **T1DES** pipeline with a visual diagram given in Figure 1. Our pipeline requires the the mass history, infall properties (i.e scale radius r_s , virial radius r_v , and time since infall t_{inf}) and tidal tracks (i.e $R_{\max}(t)$, $V_{\max}(t)$) [33]. In Section II A we discuss generating generating realizations of lens like hosts and their subhalos in **galacticus** to obtain these quantities. After generating the realizations in **galacticus** we proceed to evolve the halos in CDM

mode in our modified version of **NSphere**. While 1-D N-body simulations provide a significant speed up of full 3-D N-body simulations they provide a unique challenge: simulating asymmetric dynamics of a tidal force in 1-D. Instead of calculating a tidal field directly from the subhalo's orbit, we do a dynamic "first pass" to calculate the force at every timestep required to match a subhalo's mass history; then we rerun the subhalo with a smoothed version of this tidal field. We discuss this as well as the dynamics implemented in our simulations in Section II B. Finally, we run **NSphere** in SIDM mode using a smoothed tidal field; we give an overview of the SIDM implementation in Section II C.

A. Subhalo Realizations

We aim to simulate realizations of lensed subhalos in a $10^{13} M_{\odot}$ host in SIDM; our **T1DES** pipeline requires subhalo properties (mass history, infall properties) as a starting point. To obtain these quantities we simulate subhalo populations we use the semi-analytic galaxy formation model, **galacticus** to generate subhalo populations. Semi-analytic are faster than traditional 3D N-body simulations and allow us to realize subhalo populations quickly. The **galacticus** model is particularly well suited for our use case because of it's recent calibration to high resolution idealized simulations [34] and it's extensive and well tested library of dark matter physics. However, we note that our model is not limited to only using subhalo properties obtained using the **galacticus** galaxy formation model. It is possible to use evolutionary histories obtained from other sources such as N-body simulations, which we explore later in this paper.

We give a brief summary of the **galacticus** dark matter models used and then discuss the simulation configurations used and selection criteria for subhalos. First we discuss the discuss the procedure **galacticus** uses to simulate subhalo populations: first, merger trees are generated and then evolved forward in time using a set of analytic and semi-analytic models. Finally, we discuss our simulation parameters and the selection criteria for lensing like subhalos.

Merger tree generation — We build our **galacticus** merger trees algorithms presented in Cole *et al.* [35] and Parkinson *et al.* [36]. The algorithm is works by calculating branching probabilities: step back in time and at each time step calculating branching probabilities using the extended Press and Schechter [37] formalism. Using the branching probabilities, a merger tree is then built backwards in time that includes both merger events (infalling halos) and smooth accretion. Halo scale radii are set using the Ludlow *et al.* [38] and concentrations are set using the Diemer and Joyce [39] model.

Node evolution — After merger trees are built, our **galacticus** model tracks infalling (subhalos) halos and forward evolves them in time; first subhalo's orbits are




FIG. 1. A diagram of the **T1DES** pipeline simulating a single subhalo. The **T1DES** pipeline begins with extracting infall properties (r_s , r_v , z_{infall}) as well as the bound mass of the subhalo as a function of time. Our main results use the **galacticus** galaxy formation model to extract the infall mass properties of the subhalo, however we also include results using properties extracted from N-body simulations. These are used to set the initial conditions of the halo and begin running a first pass of NSphere in “dynamic” tidal field mode. During this run, NSphere calculates a tidal field “in flight” that results in a mass history approximating that of the galacticus halo. After the run, the tidal field is then smoothed on the scale of the dynamical time of the subhalo to remove the rapid variations in the initial calculation of the tidal field. For comparison, we rerun NSphere in CDM mode to get a “replay” of the subhalo’s evolution using the smoothed tidal field. Then we run NSphere in SIDM mode with the smoothed tidal field. Finally, for comparison we run a NSphere SIDM on an isolated halo corresponding to the infall properties of the subhalo.

initialized, then their orbits and profiles are evolved using a set of analytic models. When a subhalo infalls (a merger event happens) it’s orbit is initialized according to the potential of the host. After initialization the subhalo’s orbit is evolved forward in time using models for dynamical friction, tidal heating and tidal stripping. We give a high level overview of these models here:

- **Dynamical friction** — Dynamical friction causes subhalo orbits to decay; our configuration uses the [40] model with a Columb logarithm $\log \lambda = ?$ tuned to the Griffen *et al.* [41] simulations.
- **Tidal Heating** — Interactions with the tidal field of the host dynamically heat the host’s dark matter density causing the profile to expand. Our **galacticus** models use the Gnedin *et al.* [42] tidal heating model tuned to high resolution idealized simulations provided by Du *et al.* [34].
- **Tidal Stripping** — Mass outside of the tidal radius is removed by tidal forces. Our **galacticus** models use the Zentner *et al.* [43] tidal stripping model with and is tuned to high resolution idealized simulations provided by Du *et al.* [34].

Simulation Configuration — Using the `galacticus` model we simulate the subhalo population of 10 lensing like $10^{13} M_\odot$ halos. In this work, we focus on subhalos in the mass range $10^9 \leq m/M_\odot \leq 10^{10}$. Our `galacticus` models have an infall mass resolution of $8.00 \cdot 10^8$ and subhalos are destroyed if they lose more than 90% times their infall mass due to tidal effects. For our T1DES simulations, we select "lensing like" subhalos: subhalos with a $20 < r_{2d}/\text{kpc} < 50$ projection of the center of the host. These are subhalos that are likely to be sampled from in a strong lensing study. We also exclude subhalos that have been in the subhalo for less than 1.0Gyr as they likely have not completed a single orbit. Subhalos within x kpc of the host are destroyed.

For this analysis we only first-order substructure; that is subhalos that are field halos until they fall into the primary host. This contrasts with 2nd order and higher subhalos which are accreted into intermediate halos before infalling into the host.

- **Extreme mass histories** — `galacticus` often predicts extreme early mass loss histories for 2nd order and higher subhalos. This is likely an artifact of the way `galacticus` handles the orbits of these subhalos; these extreme mass histories characterized by rapid early mass loss when accreted into the first host and much steadier mass loss when accreted into the final host. This is likely an artifact due to how our `galacticus` handles subhalo orbits: with 2nd order substructure disabled some subhalos rapidly losing mass on first accretion then when the first host is accreted, the subhalo abruptly stops losing mass to its first host and begins losing mass to its new host. Our pipeline often disagrees with `galacticus` on the internal properties of these subhalos. We leave further investigation to future work.
- **Simplifying our analysis** — In our `galacticus` models, 2nd order subhalos have 2 distinct orbits: one in the precursor subhalo and one in the primary host; this would significantly complicate our analysis. For the purpose of this work, we study the simpler case of 1st order substructure and leave these complexities to future work.

Due to these complications, we leave exploring extending the T1DES pipeline to higher order substructure to future work.

B. Gravitational Dynamics

We modify the 1D N-body code `NSphere` to include the influence of a linear tidal force. The tidal dynamics experienced by a subhalo's particle are asymmetric in our subhalo's coordinate frame. To model tidal physics in

our spherically symmetric simulations, we need to choose an effective tidal field; we choose our tidal field to match the target halo's bound mass loss rate. Choosing a tidal field strength can be thought of as choosing a tidal radius. We would expect mass outside the Roche volume of the subhalo to be lost on order the dynamical time of the subhalo. Our choice to match the mass loss history of the subhalo is natural: where the effective tidal force dominates the gravitational subhalo, the subhalo loses mass; this forms a volume () outside of which mass is being lost; we choose a tidal radius enclosing approximately the same mass as this volume.

Under the influence of gravity and a linear tidal force with the assumption of spherical symmetry, the acceleration of i th shell is

$$\ddot{r}_i = -\frac{GM(< r_i)}{r_i^2} + \frac{l_i^2}{r_i^3} + f_t r_i \quad (1)$$

, where r_i is the distance from to center of the subhalo from the i th particle, l_i is the angular momentum per unit mass on each particle, $M(< r_i)$ is the mass enclosed by radius r_i and f_t is the linear tidal field. For each T1DES halo, we calculate a aim to calculate a tidal field to approximate the mass loss history of its corresponding galacticus subhalo. Following [34], outside the tidal radius, we assume the rate of mass loss to be on the dynamical timescale of the halo

$$\frac{dM_{\text{sub}}}{dt} = -\alpha_s \frac{M_{\text{sub}}(> r_t)}{T_{\text{dyn}}} \quad (2)$$

where M_{sub} is the gravitationally bound mass of the subhalo, α_s controls the tidal stripping rate, r_t is the tidal radius, and $T_{\text{dyn}} = 2\pi\sqrt{r_t^3/16GM_{\text{sub}}(< r_t)}$ is the dynamical time of the subhalo. For a T1DES subhalo with bound mass M_T at time t we solve the following equation

$$-\alpha_s \frac{M_T(t, > r_t)}{T_{\text{dyn}}} = \min \left\{ \frac{M_g(t + \Delta t) - M_T(t)}{\Delta t}, 0 \right\} \quad (3)$$

to calculate a tidal radius that results the reduction in mass to the mass galacticus halo, M_g at time $t + \Delta t$; we do not allow the right hand side of the equation to be greater than zero since the subhalo can only lose mass. Due to the self-correcting feedback introduced by the right hand side of Eq. 2, we find our simulations are fairly insensitive to our choice of α_s , we choose $\alpha_s = 4$ to match [34]. The tidal force and gravitational force balance at the tidal radius, r_t ; assuming a linear tidal force $F_t \propto r$ the tidal field corresponding to a tidal radius r_t is

$$f_t = \frac{GM(< r_t)}{r_t^3} \quad (4)$$

. Since our prescription can produce large jumps in the tidal field, we smooth the tidal field on order the dynamical time of the subhalo, T_{dyn} in post processing; we find that running with the smoothed tidal field using




FIG. 2. The evolutionary history of an example CDM T1DES subhalo and its `galacticus` counterpart. We show the bound mass history, M_{bound} as well as the internal properties of the subhalo, R_{max} and V_{max} (see Eq. 7). Each sudden drop in the mass history corresponds to a peri-centric passage of the subhalo, this subhalo completes 10 pericentric passages in its 7Gyr evolution within the host, losing $\sim 90\%$ of its mass since infall. Results from both T1DES stages are shown: results from the T1DES dynamic stage, where the is dynamically calculated as well as the results from the T1DES dynamic stage, where we rerun with the dynamically calculated tidal field smoothed by a gaussian kernel. We note that while our model calculates a tidal field to match the provided `galacticus` mass history, it is not tuned to match the internal R_{max} and V_{max} properties; even so it is in good agreement with `galacticus` for nearly all halos.

a gaussian filter with width $1/10T_{\text{dyn}}$ produces nearly identical halo evolution to the original run.

We show the evolutionary history for both stages of our T1DES model as well as its `galacticus` counterpart in Figure IIB. Due to our choice of tidal field, in CDM, our model is in excellent agreement with the bound mass history of both `galacticus` and 3D idealized N-body simulations. In addition, we also compare our model's prediction for the internal properties R_{max} and V_{max} curves to `galacticus` and predictions from idealized N-body simulations with the caveat that our model sometimes overshoots the mass loss from `galacticus` simulations. Figure IIB shows an important diagnostic for our simulations: we plot the normalized tidal tracks, $R_{\text{max}}/R_{\text{max},0}$ vs $V_{\text{max}}/V_{\text{max},0}$ for each subhalo at apocenter. We include T1DES CDM model, along with results from our `galacticus` model and the empirical Errani and Navarro




FIG. 3. We plot the normalized tidal tracks from our T1DES CDM smoothed stage simulations, our `galacticus` models and from the empirical Errani and Navarro [44]. For each subhalo in our suite, we plot the $R_{\text{max}}/R_{\text{max},0}$ and $V_{\text{max}}/V_{\text{max},0}$ at the subhalo's apocenter where $R_{\text{max},0}$ and $V_{\text{max},0}$ are the initial R_{max} and V_{max} of the subhalo at apocenter. We plot the Normalized tidal tracks from `galacticus`, T1DES and predictions. When properly normalized, CDM tidal tracks are expected to follow a near universal relation. We find general good agreement on the tidal track level between our simulations; with slight disagreement for low R_{max} and V_{max} .

[44] model; These curves are particularly important as they are the scale on which SIDM evolution happens [17]. When properly normalized, CDM tidal tracks are expected to follow a near universal relation. We find general good agreement on the tidal track level between our simulations; with slight disagreement for low R_{max} and V_{max} , however we note that this discrepancy is nearly as large as the difference between the `galacticus` model and Errani and Navarro [44]. Additionally, the `galacticus` tidal tracks results form nearly a perfect curve with very little scatter; while our T1DES model has more significant scatter: likely due to the noise discrete noise inherent to N-body simulations. While our model is not specifically calibrated to match these curves, our model is in reasonable agreement with both the results from `galacticus` and 3D N-body simulations.

C. SIDM Physics

Here, we give an overview of the SIDM physics implemented in our simulations; we use the default NSphere SIDM implementation with the addition of an adaptive timestep. We discuss our choice of cross sections and finally, we discuss the termination conditions we impose on the simulation and how we define core collapse.

SIDM Scattering — The probability of the i th particle scattering in the simulation within some time scale Δt is evaluated by summing over all particles from r_1 to $r_1 + \Delta r_i$

$$P_1(r_1, v_1, c_{\theta_1}) = \frac{\Delta t}{2} \left(\frac{m_p}{4\pi r_1^2 \Delta r} \right) \sum_{r_i=r_1}^{r_1+\Delta r} \sigma_{\text{eff}}(v_{\text{rel},i}) v_{\text{rel},i}, \quad (5)$$

where m_p is the particle mass, v_{rel} is the relative velocity between the two particles and $\sigma_{\text{eff}}(v_{\text{rel}}) = \frac{1}{m} \int_{-1}^1 d\cos\alpha (d\sigma/d\cos\alpha)$ is the total cross section with m being the mass of the elementary particle. We use the default NSphere bin width, $\Delta r = r_{i+j} - r_{i_1}$ with $j_1 = 10$.

After particles scatter, their kinematics must be calculated. When two identical particles with velocities $\mathbf{v}_1$ and $\mathbf{v}_2$ scatter, the Center of Mass (COM) frame has velocity $(\mathbf{v}_1 + \mathbf{v}_2)/2$. COM frame, the particles have outgoing velocities $v_{\text{cm}1}$ and $v_{\text{cm}2}$ both having equal magnitudes to half the incoming relative velocity $v_{\text{rel},2}/2$ and offset by angle α . We assume isotropic scattering; therefore, the cosine of the scattering angle, $\cos\alpha$ is chosen uniformly from -1 to 1 and the polar angle ϕ_f around $v_{\text{rel},2}$ is chosen uniformly on the interval 0 to 2π . After the velocities have been calculated, they are projected into the halo frame and saved for the next timestep.

Timestep — The key timescale for SIDM evolution is the scattering time, $t_{\text{scatter}} = [\rho\sigma_m v]^{-1}$. To accurately model this evolution, the chosen timestep of the simulation, Δt_{step} must be much less than the scattering time ($\Delta t_{\text{step}} < t_{\text{scatter}}$). Therefore, the size of the timestep sets the maximum density (and velocity) that can be resolved by the simulation; by default, NSphere implements a constant timestep, $\Delta t_{\text{step}} = C$. In this method, the fixed timestep size is set by choosing a maximum resolvable density. However, this is inefficient as a very small timestep must be chosen to resolve the high density of core collapse and the halo will only reach this density at the very end of its evolution. Instead, we implement an adaptive timestep; we choose the size of the adaptive timestep based on the probability of two particles scattering in the previous timestep, P_i (see Eq. 5)

$$\Delta t_{\text{step},n+1} = \max \left\{ \epsilon \cdot \frac{\Delta t_{\text{step},n}}{\max\{P_i\}}, \Delta t_{\text{step},\min} \right\} \quad (6)$$

where we ensure that if the particles maintained the same scattering probability in the next timestep, the maximum scattering probability would be ϵ . For our simulations, we find good convergence with $\epsilon = 1.0 \cdot 10^{-4}$. Taking the minimum of all particle scattering probabilities can result in very small time steps, for instance due to a single pair of particles that are highly likely to scatter in the next timestep; we prevent these extremely small timesteps by implementing a timestep floor $\Delta t_{\text{step},\min}$. We set the timestep floor proportional to the maximum density we expect to resolve in the simulation, $\Delta t_{\text{step},\min} = [\bar{\rho}_{\text{max}}(< R_{\text{max}})\sigma_m \max\{V_{\text{max}}\}]$ where $\bar{\rho}_{\text{max}}(< R_{\text{max}})$ is the termination density of our simulations and V_{max} is the circular

velocity predicted for the target halo (in this case, we use the result from **galacticus** model). We conveniently use $\{V_{\text{max}}\}$ as the characteristic velocity because for the cross-sections studied and the termination condition we use, the termination V_{max} in SIDM remains slightly higher than (maximum $\sim 20\%$ more) or less than the $\max\{V_{\text{max}}\}$ in CDM due to tidal stripping.

Cross Section — For this work, we explore a constant cross section per unit mass σ_{eff} and run simulations with 5 cross-sections $\sigma_m/(\text{cm}^2/\text{g}) = 5, 10, 25$ and 50 and assume a SIDM model with elastic scattering. In this work we focus on core collapsed halos; strong lensing probes are most sensitive to core collapsed halos and insensitive to cored halos [45]. Constant SIDM cross sections constraints from clusters [46–48] and dwarf and satellite galaxies [49, 50] constrain constant cross section SIDM models to cross sections below $\sigma_m < 1\text{cm}^2/\text{g}$ which is lower cross section than considered in this work, and below the cross sections we find subhalos collapsing in our simulations. In our simulations, we do not consider SIDM particle interactions with the host; if SIDM interactions with host particles are not suppressed by (as is the case with a truly constant cross section SIDM models) subhalos are not expected to collapse due to evaporation [13]. Therefore, the constant cross sections in this work should be considered as an illustrative case study rather than something that is expected to be probed by strong lensing; mappings exist from velocity dependent cross sections to constant "effective" cross sections, which allow mapping our work onto velocity dependent cross section models, see References [51–53]. Finally, we note many velocity-dependent cross sections considered in lensing analysis such as parametric cross section form considered in [45] asymptotes to a nearly constant cross section in the low velocity regime; these results can be viewed as exploring that regime: near constant cross section at low velocities and highly suppressed cross section on the scale of the host.

Core Collapse and Termination Conditions — Unlike field halos, the characteristic scale of subhalos may greatly decrease over time due to the effects of tidal stripping; to define core collapse we cannot simply choose a density threshold value at a fixed radius. Instead, we use the R_{max} as the characteristic scale, where R_{max} is the radius at which the maximum circular velocity V_{max}

$$V_{\text{max}} = \text{Max} \left\{ \sqrt{\frac{GM(< r)}{r}} \right\} = \sqrt{\frac{GM(< R_{\text{max}})}{R_{\text{max}}}} \quad (7)$$

occurs. From Eq. 7, it follows that the average density within R_{max} , $\bar{\rho}_{\text{max}}(< R_{\text{max}})$ is

$$\bar{\rho}_{\text{max}}(< R_{\text{max}}) = \frac{3}{4\pi G} \left(\frac{V_{\text{max}}}{R_{\text{max}}} \right)^2 \quad (8)$$

. We define a subhalo as core collapsed and terminate with when the density within R_{max} is

$$\bar{\rho}_{\text{max}}(< R_{\text{max}}) > n \frac{3}{4\pi G} \left(\frac{\max\{V_{\text{max}}(t)\}}{\min\{R_{\text{max}}(t)\}} \right)^2 \quad (9)$$

. where $n > 1$ is an arbitrary factor that determines how deep into core collapse we resolve; for this work we chose $n = 10$.

III. RESULTS

Here we discuss results from our suite of **T1DES** simulations. First, we compare results of subhalos with their isolated counterparts. We find that subhalos core collapse rates are strictly accelerated by the presence of a tidal field when compared to their isolated counterparts. Next, we compare our results to previous work including the Yang *et al.* [17] parametric model for core collapse. We find good agreement with the parametric model and a similar order of magnitude core collapse acceleration when compared to Nadler *et al.* [16].

Core Collapse Acceleration — Figure 4 shows the the cross section per unit mass vs the core collapse fraction, the number of total core collapsed subhalos divided by total number of subhalos, f_{collapse} for our **T1DES** simulations compared to the predictions from [17] parametric model. We compare our tides subhalos run with a tidal field (**T1DES** SIDM stage) to their counterparts ran without a tidal field using the same initial conditions (**T1DES** isolated stage). We find that all isolated halos that collapse also collapse when run with a tidal field; the influence of a tidal field in our simulations strictly accelerates core collapse rate. However, we note that our results assume evaporation is negligible; for SIDM models where this is not the case, core collapse may be delayed instead of accelerated. We find roughly similar acceleration rates to Nadler *et al.* [16] who compared the predicted core collapse rates for populations of isolated halos to subhalos in a SIDM model with a velocity dependent cross section; at $\sigma_m = 25\text{cm}^2/\text{g}$ nearly 4 times as many subhalos collapse when compared to their isolated counterparts and at $\sigma_m = 50\text{cm}^2/\text{g}$ (46%) nearly 2 times as many subhalos collapse. We find subhalo collapse for a cross section as low as $\sigma_m = 10\text{cm}^2/\text{g}$; at this cross section around 5% of the subhalos in our models collapse while none of their isolated counterparts do. For some SIDM models it may be possible that a non-negligible fraction of subhalos core collapse before any field halos do, due to accelerated core collapse from a tidal field.

Comparison with previous results — Figure 4 also shows the collapse fraction f_{collapse} , for our **T1DES** simulations compared to the predictions from the Yang *et al.* [17] model; for this model, we show results obtained by integrating over the the tidal tracks from **galacticus** and from the **T1DES** CDM. We find our **T1DES** model is fairly consistent with the predictions from the Yang *et al.* [17] model, though our **T1DES** model may predict slightly higher fractions of core collapsed halos. Interestingly our the Yang *et al.* [17] predictions integrating over the **galacticus** tidal tracks appear to agree slightly better with our **T1DES** predictions than the the

results obtained by integrating over the **galacticus** tidal tracks. We also note that the core collapse definition between our **T1DES** model and the Yang *et al.* [17] model is different; Yang *et al.* [17] defines core collapse as $\tau > 0.75$ while we use the R_{max} based density threshold given by Eq. 9. However, the difference in definitions likely only impacts the number of collapsed halos slightly; the final core collapse happens on short timescale compared to the entire SIDM evolution. We leave a more detailed comparison and investigating these possible discrepancies to future work.

IV. CONCLUSION AND SUMMARY

In this work, we have presented **T1DES**, a novel method of simulating SIDM subhalos using a modified version of the **NSphere** 1D N-body code. Here we provide a summary of this work. In Section II we discuss the methods used in this paper in three subsections. First, in Section IIA we discuss the **galacticus** model used to obtain the mass histories used by **T1DES** and the initial conditions used in our **T1DES** simulations. Then in Section IIB we discuss the dynamics implemented in our fork of **NSphere**; in particular we focus on the tidal physics implemented in our simulations: a linear effective tidal field, calculated in a CDM first pass to match the mass history of a target subhalo. Thirdly, in Section IIC we give an overview of the SIDM physics implemented in **NSphere** and discuss our core collapse conditions for subhalos. In the main section, Section III we discuss the main results from this work: with no evaporation, find that a tidal field strictly accelerates core collapse; and that this acceleration is cross section dependent and significant: up to a factor of nearly 4 time the number of subhalos collapse when compared to their isolated counterparts. Finally, here we discuss future work, discuss some potential caveats and provide concluding statements.

To correctly infer constraints on SIDM models through lensing studies, it will be important to thoroughly understand predicted core collapse rates. Our model gives a distinct advantage here: because it is much more computationally efficient than previous SIDM simulations, by making modifications to our pipeline it may now be practical to test a wider range of parameter and model space within SIDM. In addition to more thoroughly exploring the parameter space of specific SIDM due to the reduction of computational cost in our simulations By making some modifications to our code it is possible to explore a wide range of velocity and angle dependent cross sections. By adding in stochastic scattering chance of a particle interaction with the host simulating evaporation is possible within our model; we leave exploration of this to future work. Additionally, **NSphere** is not limited to dark matter only physics; it may be interesting to explore the effect of a central galaxy on core collapse times.

Our **T1DES** model is in reasonable agreement with




FIG. 4. The cross section per unit mass vs the core collapse fraction for our T1DES simulations compared to as well as predictions from [17]. In red: results from our tides subhalo simulations (red dot with error bars), and the Yang models integrated over the the tidal tracks from `galacticus` (red solid) and from the T1DES CDM (red dotted) in blue. We find our T1DES model is fairly consistent with the predictions from the [17] model, though our T1DES model may predict slightly higher fractions of core collapsed halos. In blue is the results for the isolated case, if the subhalos remained isolated, what fraction would collapse? Our T1DES model is consistent with the predictions of the Yang model here and predicts strictly lower collapse fractions for isolated halos. No halos collapse in the field that also do not collapse as a subhalo.

`galacticus` the tidal tracks of `galacticus` halos, however some outlier halos disagree on order 50% of V_{max} ; this disagreement is particularly pronounced for subhalos that are early in their evolution. This may be due to the initial conditions used by N-Sphere and the non physical stair stepping mass histories predicted by `galacticus`. It may be worth further exploring seeding the T1DES pipeline with N-body mass histories rather than `galacticus` mass histories. We do not include any pre-infall SIDM evolution in our models; depending on the cross section, pre-infall SIDM evolution may be significant enough to form a core that effects the tidal physics of the subhalo. We leave investigating this regime to future work.

In conclusion, by simulating SIDM subhalo evolution in 1D, we are able to be much more computationally efficient in simulating SIDM subhalos when compared to traditional 3D N-body simulations by over an order of magnitude; our simulations are able to be simulated on a laptop and ran in the span of hours (this is greatly particle count and thus mass loss dependent). We shared Previous 1D SIDM subhalo simulations relied on coarse approximations of tidal stripping, such as approximating the tidal stripping as per orbit truncations to the mass profile; we include a much more realistic mass profile that reasonably approximates the mass history over time as well as the internal R_{max} and V_{max} properties of the subhalo. We find that under our assumption of no evaporation, subhalo core collapse is strictly accelerated when compared to isolated field halos. Additionally, we find that our simulations are in broad agreement with the Yang *et al.* [17] parametric model.

ACKNOWLEDGMENTS

We wish to acknowledge the support of the author community in using REVTeX, offering suggestions and encouragement, testing new versions,

Generative AI Disclosure — Generative AI (LLMs) were used in some portions of the research process for this work. Several Claude models were used as a part of this project: Claude Haiku, Sonnet, Opus and Fable[54] were used for coding assistance (including the generation of plots), summarization of background literature, as a general learning tool, transcriptions (taking an image of a diagram drawn by pencil and converting it to latex) and for several discussions about the project. We disclose that coding agents were used for several portions of this project such as writing portions of the tidal physics model, packaging T1DES into an ergonomic command line utility, implement various features into tides such as the "smoothed" and "isolated" modes, various assistance with analysis such as the generation of the figures in this work. Generative AI was also used for summarization of literature and as a general learning tool "E.X What are the ways core collapse in previous SIDM works?". Generative AI was not used in the writing of this manuscript. For some early exploratory work, discussions and brainstorming with AI was tested "E.x What are some ideas for figures?". However, this usage was found to be infetive and for this manuscript nearly all useful AI assistance was directed coding tasks, literature summarization, transcription and as and as a learning tool. We verify the output from AI coding agents in several ways: code generated by agents was reviewed, tests suggested by both agents and humans were set up

and manually reviewed by the authors and finally, for

a variety of tasks was duplicated and key results were checked by human made programs.

-
- [1] Planck Collaboration, Planck2018 results: Vi. cosmological parameters, *Astronomy & Astrophysics* **641**, A6 (2020).
 - [2] S. D. White and M. J. Rees, Core condensation in heavy halos: a two-stage theory for galaxy formation and clustering, *Monthly Notices of the Royal Astronomical Society* **183**, 341 (1978).
 - [3] S. D. White and C. S. Frenk, Galaxy formation through hierarchical clustering, *Astrophysical Journal*, Part 1 (ISSN 0004-637X), vol. 379, Sept. 20, 1991, p. 52-79. Research supported by NASA, NSF, and SERC. **379**, 52 (1991).
 - [4] W. De Blok, F. Walter, E. Brinks, C. Trachternach, S. Oh, and R. C. Kennicutt, High-resolution rotation curves and galaxy mass models from things, *The Astronomical Journal* **136**, 2648 (2008).
 - [5] D. H. Weinberg, J. S. Bullock, F. Governato, R. Kuzio de Naray, and A. H. Peter, Cold dark matter: controversies on small scales, *Proceedings of the National Academy of Sciences* **112**, 12249 (2015).
 - [6] S. Vegetti, L. Koopmans, A. Bolton, T. Treu, and R. Gavazzi, Detection of a dark substructure through gravitational imaging, *Monthly Notices of the Royal Astronomical Society* **408**, 1969 (2010).
 - [7] Q. Minor, S. Gad-Nasr, M. Kaplinghat, and S. Vegetti, An unexpected high concentration for the dark substructure in the gravitational lens sdssj0946+ 1006, *Monthly Notices of the Royal Astronomical Society* **507**, 1662 (2021).
 - [8] M. Tajalli, S. Vegetti, C. M. O’Riordan, S. D. White, C. D. Fassnacht, D. M. Powell, J. P. McKean, and G. Despali, Sharp-ix. the dense, low-mass perturbers in b1938+ 666 and j0946+ 1006: implications for cold and self-interacting dark matter, *Monthly Notices of the Royal Astronomical Society* **543**, 540 (2025).
 - [9] Q. He, A. Robertson, J. W. Nightingale, A. Amvrosiadis, S. Cole, C. S. Frenk, S. C. Lange, S. Li, R. Li, X. Cao, *et al.*, Not so dark, not so dense: an alternative explanation for the lensing subhalo in sdss j0946+ 1006, *The Astrophysical Journal Letters* **991**, L53 (2025).
 - [10] X. Zhang, H.-B. Yu, D. Yang, and E. O. Nadler, The gd-1 stellar stream perturber as a core-collapsed self-interacting dark matter halo, *The Astrophysical Journal Letters* **978**, L23 (2025).
 - [11] S. Tulin and H.-B. Yu, Dark matter self-interactions and small scale structure, *Physics Reports* **730**, 1 (2018).
 - [12] S. Adhikari, A. Banerjee, K. K. Boddy, F.-Y. Cyr-Racine, H. Desmond, C. Dvorkin, B. Jain, F. Kahlhoefer, M. Kaplinghat, A. Nierenberg, *et al.*, Astrophysical tests of dark matter self-interactions, *Reviews of Modern Physics* **97**, 045004 (2025).
 - [13] Z. C. Zeng, A. H. Peter, X. Du, A. Benson, S. Kim, F. Jiang, F.-Y. Cyr-Racine, and M. Vogelsberger, Core-collapse, evaporation, and tidal effects: the life story of a self-interacting dark matter subhalo, *Monthly Notices of the Royal Astronomical Society* **513**, 4845 (2022).
 - [14] N. Shah and S. Adhikari, The abundance of core-collapsed subhaloes in sidm: insights from structure formation in λ cdm, *Monthly Notices of the Royal Astronomical Society* **529**, 4611 (2024).
 - [15] H. Nishikawa, K. K. Boddy, and M. Kaplinghat, Accelerated core collapse in tidally stripped self-interacting dark matter halos, *Physical Review D* **101**, 063009 (2020).
 - [16] E. O. Nadler, D. Kong, D. Yang, and H.-B. Yu, Sidm concerto: compilation and data release of self-interacting dark matter zoom-in simulations, *The Astrophysical Journal* **991**, 69 (2025).
 - [17] D. Yang, E. O. Nadler, H.-B. Yu, and Y.-M. Zhong, A parametric model for self-interacting dark matter halos, *Journal of Cosmology and Astroparticle Physics* **2024** (02), 032.
 - [18] M. Vogelsberger, J. Zavala, and A. Loeb, Subhaloes in self-interacting galactic dark matter haloes, *Monthly Notices of the Royal Astronomical Society* **423**, 3740 (2012).
 - [19] M. Rocha, A. H. Peter, J. S. Bullock, M. Kaplinghat, S. Garrison-Kimmel, J. Onorbe, and L. A. Moustakas, Cosmological simulations with self-interacting dark matter–i. constant-density cores and substructure, *Monthly Notices of the Royal Astronomical Society* **430**, 81 (2013).
 - [20] G. A. Dooley, A. H. Peter, M. Vogelsberger, J. Zavala, and A. Frebel, Enhanced tidal stripping of satellites in the galactic halo from dark matter self-interactions, *Monthly Notices of the Royal Astronomical Society* **461**, 710 (2016).
 - [21] E. O. Nadler, A. Banerjee, S. Adhikari, Y.-Y. Mao, and R. H. Wechsler, Signatures of velocity-dependent dark matter self-interactions in milky way-mass halos, *The Astrophysical Journal* **896**, 112 (2020).
 - [22] H. C. Turner, M. R. Lovell, J. Zavala, and M. Vogelsberger, The onset of gravothermal core collapse in velocity-dependent self-interacting dark matter subhaloes, *Monthly Notices of the Royal Astronomical Society* **505**, 5327 (2021).
 - [23] T. Ebisu, T. Ishiyama, and K. Hayashi, Constraining self-interacting dark matter with dwarf spheroidal galaxies and high-resolution cosmological n-body simulations, *Physical Review D* **105**, 023016 (2022).
 - [24] D. Yang, E. O. Nadler, and H.-B. Yu, Strong dark matter self-interactions diversify halo populations within and surrounding the milky way, *The Astrophysical Journal* **949**, 67 (2023).
 - [25] E. O. Nadler, R. An, D. Yang, H.-B. Yu, A. Benson, and V. Gluscevic, Cosmic. iii. cosmological zoom-in simulations of self-interacting dark matter with suppressed initial conditions, *The Astrophysical Journal* **986**, 129 (2025).
 - [26] F. Kahlhoefer, M. Kaplinghat, T. R. Slatyer, and C.-L. Wu, Diversity in density profiles of self-interacting dark matter satellite halos, *Journal of Cosmology and Astroparticle Physics* **2019** (12), 010.
 - [27] Z. C. Zeng, A. H. Peter, X. Du, S. Yang, A. Benson, F.-Y. Cyr-Racine, F. Jiang, C. Mace, and R. B. Metcalf, Till the core collapses: The evolution and properties

- of self-interacting dark matter subhalos, arXiv preprint arXiv:2310.09910 (2023).
- [28] D. Klemmer, M. S. Fischer, K. K. Boddy, M. Kaplinghat, and L. Sagunski, Survival of the most compact: the life and death of satellite halos in self-interacting dark matter, arXiv preprint arXiv:2603.19362 (2026).
 - [29] A. J. Benson, Galacticus: A semi-analytic model of galaxy formation, *New Astronomy* **17**, 175 (2012).
 - [30] M. Kamionkowski, K. Sigurdson, and O. Slone, Numerical evolution of self-gravitating halos of self-interacting dark matter, *Physical Review Letters* **136**, 201001 (2026).
 - [31] M. Kamionkowski and K. Sigurdson, Gravothermal collapse of self-interacting dark-matter halos with anisotropic velocity distributions, arXiv preprint arXiv:2510.23705 (2025).
 - [32] M. Kamionkowski and K. Sigurdson, Evolution of self-gravitating spherical dark-matter halos with and without new physics, *Physical Review D* **113**, 103026 (2026).
 - [33] Tidal tracks are only required for SIDM mode, as they are used in setting the simulation termination conditions.
 - [34] X. Du, A. Benson, Z. C. Zeng, T. Treu, A. H. Peter, C. Mace, F. Jiang, S. Yang, C. Gannon, D. Gilman, *et al.*, Tidal evolution of cored and cuspy dark matter halos, *Physical Review D* **110**, 023019 (2024).
 - [35] S. Cole, C. G. Lacey, C. M. Baugh, and C. S. Frenk, Hierarchical galaxy formation, *Monthly Notices of the Royal Astronomical Society* **319**, 168 (2000), <https://academic.oup.com/mnras/article-pdf/319/1/168/3734609/319-1-168.pdf>.
 - [36] H. Parkinson, S. Cole, and J. Helly, Generating dark matter halo merger trees, *Monthly Notices of the Royal Astronomical Society* **383**, 557 (2008).
 - [37] W. H. Press and P. Schechter, Formation of galaxies and clusters of galaxies by self-similar gravitational condensation, *Astrophysical Journal*, Vol. 187, pp. 425-438 (1974) **187**, 425 (1974).
 - [38] A. D. Ludlow, S. Bose, R. E. Angulo, L. Wang, W. A. Hellwing, J. F. Navarro, S. Cole, and C. S. Frenk, The mass-concentration-redshift relation of cold and warm dark matter haloes, *Monthly Notices of the Royal Astronomical Society* **460**, 1214 (2016).
 - [39] B. Diemer and M. Joyce, An accurate physical model for halo concentrations, *The Astrophysical Journal* **871**, 168 (2019).
 - [40] S. Chandrasekhar, Dynamical friction. i. general considerations: the coefficient of dynamical friction, *Astrophysical Journal* **97**, 255 (1943).
 - [41] B. F. Griffen, A. P. Ji, G. A. Dooley, F. A. Gómez, M. Vogelsberger, B. W. O'Shea, and A. Frebel, The caterpillar project: a large suite of milky way sized halos, *The Astrophysical Journal* **818**, 10 (2016).
 - [42] O. Y. Gnedin, L. Hernquist, and J. P. Ostriker, Tidal shocking by extended mass distributions, *The Astrophysical Journal* **514**, 109 (1999).
 - [43] A. R. Zentner, A. A. Berlind, J. S. Bullock, A. V. Kravtsov, and R. H. Wechsler, The physics of galaxy clustering. i. a model for subhalo populations, *The Astrophysical Journal* **624**, 505 (2005).
 - [44] R. Errani and J. F. Navarro, The asymptotic tidal remnants of cold dark matter subhaloes, *Monthly Notices of the Royal Astronomical Society* **505**, 18 (2021).
 - [45] D. Gilman, J. Bovy, T. Treu, A. Nierenberg, S. Birrer, A. Benson, and O. Sameie, Strong lensing signatures of self-interacting dark matter in low-mass haloes, *Monthly Notices of the Royal Astronomical Society* **507**, 2432 (2021).
 - [46] D. Eckert, S. Ettori, A. Robertson, R. Massey, E. Pointecouteau, D. Harvey, and I. McCarthy, Constraints on dark matter self-interaction from the internal density profiles of x-cop galaxy clusters, *Astronomy & Astrophysics* **666**, A41 (2022).
 - [47] L. Sagunski, S. Gad-Nasr, B. Colquhoun, A. Robertson, and S. Tulin, Velocity-dependent self-interacting dark matter from groups and clusters of galaxies, *Journal of Cosmology and Astroparticle Physics* **2021** (01), 024.
 - [48] K. E. Andrade, J. Fuson, S. Gad-Nasr, D. Kong, Q. Minor, M. G. Roberts, and M. Kaplinghat, A stringent upper limit on dark matter self-interaction cross-section from cluster strong lensing, *Monthly Notices of the Royal Astronomical Society* **510**, 54 (2022).
 - [49] S. Ando, K. Hayashi, S. Horigome, M. Ibe, and S. Shirai, Stringent constraints on self-interacting dark matter using milky-way satellite galaxies kinematics, arXiv preprint arXiv:2503.13650 (2025).
 - [50] C. A. Correa, Constraining velocity-dependent self-interacting dark matter with the milky way's dwarf spheroidal galaxies, *Monthly Notices of the Royal Astronomical Society* **503**, 920 (2021).
 - [51] D. Yang and H.-B. Yu, Gravothermal evolution of dark matter halos with differential elastic scattering, *Journal of Cosmology and Astroparticle Physics* **2022** (09), 077.
 - [52] S. Yang, X. Du, Z. C. Zeng, A. Benson, F. Jiang, E. O. Nadler, and A. H. Peter, Gravothermal solutions of sidm halos: Mapping from constant to velocity-dependent cross section, *The Astrophysical Journal* **946**, 47 (2023).
 - [53] N. J. Outmezguine, K. K. Boddy, S. Gad-Nasr, M. Kaplinghat, and L. Sagunski, Universal gravothermal evolution of isolated self-interacting dark matter halos for velocity-dependent cross-sections, *Monthly Notices of the Royal Astronomical Society* **523**, 4786 (2023).
 - [54] Many versions of Claude models were used throughout the project, starting at version 4.5 for haiku,sonnet and opus through to version 5.1 of fable.
 - [55] A. Bonaca, D. W. Hogg, A. M. Price-Whelan, and C. Conroy, The spur and the gap in gd-1: Dynamical evidence for a dark substructure in the milky way halo, *The Astrophysical Journal* **880**, 38 (2019).
 - [56] J. Nibauer, A. Bonaca, A. M. Price-Whelan, D. N. Spergel, and J. E. Greene, Measurement of substructure from the kinematics of the gd-1 stellar stream, *The Astrophysical Journal* **1004**, 62 (2026).
 - [57] J. Bond, S. Cole, G. Efstathiou, and N. Kaiser, Excursion set mass functions for hierarchical gaussian fluctuations, *Astrophysical Journal*, Part 1 (ISSN 0004-637X), vol. 379, Oct. 1, 1991, p. 440-460. Research supported by NSERC, NASA, and University of California. **379**, 440 (1991).
 - [58] R. G. Bower, The evolution of groups of galaxies in the press-schechter formalism, *Monthly Notices of the Royal Astronomical Society* **248**, 332 (1991).