



Maxillofacial Trauma Surgery

Purpose: Surgical repair of facial fractures, jaw injuries, or severe soft tissue damage caused by trauma.

First 24–72 Hours:

- Swelling, bruising, and discomfort are common.
- Apply ice packs to the affected area for 15–20 minutes at a time during the first 24 hours.
- Keep head elevated while resting to reduce swelling.

Pain & Sensitivity:

- Take prescribed pain relievers and antibiotics exactly as directed.
- Avoid aspirin unless recommended, as it may increase bleeding risk.

Oral Hygiene:

- Follow your surgeon's instructions for oral care; often includes gentle antimicrobial rinses.
- Avoid brushing near sutured areas until cleared.

Diet:

- Soft or liquid diet as directed, avoiding chewing on the injured side.
- Avoid hot, spicy, or crunchy foods that could irritate healing tissue.

Special Precautions:

- Do not smoke or consume alcohol during recovery.
- Avoid strenuous activities and contact sports until cleared by your surgeon.
- If jaw fixation (wiring) is used, keep wire cutters or scissors nearby in case of emergency.

Follow-Up:

- Multiple follow-up appointments will be necessary to monitor bone and soft tissue healing.
- Contact your surgeon immediately if you experience persistent bleeding, difficulty breathing, fever, or sudden changes in bite alignment.