

Post-Operative Instructions

Cement-Retained Implant Crown

Initial Adjustment

- Mild soreness or pressure around the implant area is normal for the first few days.
- Your bite may feel slightly different at first. This usually improves as your mouth adjusts.
- Increased saliva flow is normal after new crown placement.

Eating & Diet

- Avoid chewing on hard or sticky foods for the first 24–48 hours.
- After adjustment, you may return to a normal diet — but avoid chewing ice, hard candy, or very tough foods that could damage the crown.
- Chew evenly on both sides of your mouth to reduce stress on the implant.

Oral Hygiene

- Brush twice daily with a soft-bristled toothbrush.
- Floss carefully around the implant crown, using implant-safe floss or floss threaders.
- A water flosser may be used gently to keep the area clean.
- Antimicrobial rinses (if prescribed) can reduce bacteria around the crown and implant.

Special Considerations

- Cement-retained crowns are designed to look and feel natural. However, excess cement can sometimes irritate the gum tissue. If you notice persistent soreness, swelling, or bleeding, contact the office.
- Avoid using your teeth as tools (opening packages, biting nails, etc.).

Follow-Up

- Regular dental cleanings and check-ups are essential to monitor the health of the crown and implant.
- Call our office immediately if you notice loosening, pain, swelling, or unusual sensations around the crown.