

Post-Operative Instructions

Screw-Retained Implant Crown

Initial Adjustment

- Some mild tenderness or gum irritation around the implant site is normal for a few days.
- Your bite may feel slightly different at first but will adapt over time.
- You may notice a small access hole on the biting surface of the crown. This is sealed with filling material after placement.

Eating & Diet

- Avoid very hard, crunchy, or sticky foods for the first 24–48 hours.
- After this period, you may return to a normal diet — but avoid habits such as chewing ice or hard candy.
- Cut tougher foods into small pieces to reduce excess force on the crown.

Oral Hygiene

- Brush carefully twice daily, paying extra attention to the gumline around the implant.
- Floss daily using implant-safe floss or threaders to clean under the crown edges.
- Use a water flosser to remove trapped food particles if recommended by your dentist.
- Rinse daily with an antimicrobial mouthwash (alcohol-free preferred).

Special Considerations

- Screw-retained crowns can be removed by your dentist for maintenance without damaging the implant.
- The small filling material covering the screw access may wear over time and require replacement — this is normal.
- Avoid biting very hard objects that may loosen or chip the crown.

Follow-Up

- Schedule and attend regular dental visits for maintenance and professional cleaning.
- Contact our office if you feel any loosening, hear clicking when chewing, or experience pain, swelling, or bleeding near the implant.