



I.V. Sedation

Purpose: Provide safe, comfortable dental treatment using intravenous sedation while maintaining proper

First 24 Hours:

- A responsible adult should remain with you today and overnight.
- No driving, operating machinery, alcohol, recreational drugs, or signing legal documents until tomorrow.
- Expect drowsiness, lightheadedness, or mild forgetfulness—move slowly from lying to standing.
- Sleep with your head elevated; use CPAP/BiPAP if prescribed.

Pain & Swelling:

- Begin prescribed/approved pain medication before numbness fully wears off.
- If permitted, many patients alternate ibuprofen and acetaminophen; follow label limits.
- Apply ice packs to the outside of your face for 20 minutes on/off for the first 24 hours.
- Switch to warm compresses after 48 hours to help with stiffness.
- Gentle jaw stretching after 3 days can help prevent limited opening if your jaw is sore.

Diet:

- Start with clear liquids (water, electrolyte drinks, broth), then advance to soft foods (yogurt, eggs, macaroni).
- Avoid hot foods/drinks until numbness is gone to prevent burns.
- If you had extractions or grafting, avoid straws for at least 1 week to protect the blood clot.

Activity:

- Rest the day of your procedure; light activity the next day as tolerated.
- Avoid strenuous exercise, heavy lifting, or bending over for 48 hours.
- Change positions slowly to reduce dizziness or fainting.

Special Precautions:

- Take only medications listed on your after-visit summary or prescriptions; do not mix with extra sedation.
- Nausea can occur—use small sips of clear fluids; take the prescribed anti-nausea medicine if needed.
- Patients with diabetes should monitor glucose more often today and maintain hydration.
- Breastfeeding: confirm medication compatibility before nursing; you may be advised to pump and discard milk.
- Call if you have trouble breathing, bluish lips, or severe snoring that doesn't improve when repositioned.

Follow-Up:

- Contact the office for: fever over 101°F (38.3°C), heavy bleeding, worsening pain not relieved by medication.
- Use the after-hours number provided on your paperwork. If you experience severe breathing trouble