

# Solving the Incompressible Navier-Stokes Equations using "Stable Fluids"

## 1 Introduction

This document describes the method to solve the incompressible Navier-Stokes equations in a closed box scenario with a forcing function that creates a bloom. The solution strategy follows the "Stable Fluids" approach.

## 2 Governing Equations

The equations governing the flow are:

### 2.1 Momentum Equation

$$\frac{\partial \mathbf{u}}{\partial t} + (\mathbf{u} \cdot \nabla) \mathbf{u} = -\frac{1}{\rho} \nabla p + \nu \nabla^2 \mathbf{u} + \mathbf{f} \quad (1)$$


### 2.2 Incompressibility Condition

$$\nabla \cdot \mathbf{u} = 0 \quad (2)$$

where:

- $\mathbf{u}$ : Velocity (2D vector)
- $p$ : Pressure
- $\mathbf{f}$ : Forcing
- $\nu$ : Kinematic Viscosity
- $\rho$ : Density
- $t$ : Time
- $\nabla$ : Nabla operator (defining nonlinear convection, gradient, and divergence)
- $\nabla^2$ : Laplace Operator


### 3 Closed Box Scenario



- Homogeneous Dirichlet Boundary Conditions are applied everywhere.
- The velocity  $\mathbf{u}$  and pressure  $p$  have zero initial conditions.

### 4 Forcing Function

The forcing function creates an upward force in the lower center of the domain.



## 5 Solution Strategy

The solver follows these steps:

### 5.1 Initialization

Start with zero velocity everywhere:  $\mathbf{u} = [0, 0]$ .

### 5.2 Add Forces

$$\mathbf{w}_1 = \mathbf{u} + \Delta t \mathbf{f} \quad (3)$$

### 5.3 Self-Advection (Convection)

Convect by self-advection (set the value at the current location to be the value at the position backtracked on the streamline). This step is unconditionally stable.

$$\mathbf{w}_2 = \mathbf{w}_1(\mathbf{p}(\mathbf{x}, -\Delta t)) \quad (4)$$

### 5.4 Implicit Diffusion

Diffuse implicitly by solving a linear system matrix-free using Conjugate Gradient. This step is unconditionally stable.

$$(I - \nu \Delta t \nabla^2) \mathbf{w}_3 = \mathbf{w}_2 \quad (5)$$

### 5.5 Pressure Correction

1. Compute a pressure correction by solving a linear system matrix-free using Conjugate Gradient.

$$\nabla^2 p = \nabla \cdot \mathbf{w}_3 \quad (6)$$

2. Correct velocities to be incompressible.

$$\mathbf{w}_4 = \mathbf{w}_3 - \nabla p \quad (7)$$

### 5.6 Advance to Next Time Step

$$\mathbf{u} = \mathbf{w}_4 \quad (8)$$

## 6 Boundary Conditions

Since the fluid is inside a box, completely enclosed by walls. There is no flow velocity at the boundary. The boundary conditions are prescribed indirectly using the discrete differential operators.

## 7 Stability

The solver is unconditionally stable; hence, all parameters can be chosen arbitrarily. However, high timesteps can make the advection step highly incorrect.

## 8 Output

