

Sums. Let n be a natural number and x be a real number.

- $0 + 1 + 2 + \cdots + n = \frac{n(n+1)}{2}$
- $0 + 1^2 + 2^2 + \cdots + n^2 = \frac{n(n+1)(2n+1)}{6}$
- $1 + x + x^2 + \cdots + x^n = \frac{x^{n+1}-1}{x-1}$

Pascal's Rule. Let n be a positive integer and k be a natural number less than n .

- $\binom{n}{k} + \binom{n}{k+1} = \binom{n+1}{k+1}$

Binomial Theorem. Let n be a natural number.

- $(x+y)^n = \binom{n}{0}x^0y^n + \binom{n}{1}x^1y^{n-1} + \cdots + \binom{n}{n}x^ny^0$

Divisibility Properties. Let a, b, c, d be integers.

- If $a \mid b$ and $c \mid d$, then $ac \mid bd$.
- If $a \mid b$ and $b \mid c$, then $a \mid c$.
- If $a \mid b$ and $a \mid c$, then $a \mid (bx + cy)$ for integers x, y .
- If $a \perp b$, and $a \mid c$ and $b \mid c$, then $ab \mid c$.
- If $a \mid bc$ and $a \perp b$, then $a \mid c$.
- If $a \mid b$ and $b \perp c$, then $a \perp c$.

Divisors/Multiples. Let a, b be integers w/ $\gcd(a, b) = d$.

- Any common divisor of a and b also divides d .
- d is the least pos. integral combination of a and b .
- $(a/d) \perp (b/d)$
- If k is a pos. integer, then $\gcd(ka, kb) = kd$.
- If c is another integer, $\gcd(a, b, c) = \gcd(d, c)$.
- $\text{lcm}(a, b) = (ab)/d$ if a and b are positive.
- $\text{lcm}(a, b)$ divides any common multiple of a and b .
- If $a \perp b$, then the pos. divisors of ab are uniquely of the form d_1d_2 where pos. $d_1 \mid a$ and pos. $d_2 \mid b$.

Diophantines. Let a, b, c be integers with $\gcd(a, b) = d$. Consider Diophantine equation $ax + by = c$.

- It admits an integer solution pair iff $d \mid c$.
- If x_0, y_0 is one solution pair, the general solution is:

$$x = x_0 + (b/d)t$$

$$y = y_0 - (a/d)t$$

Primes. Let p be prime and $a, a_1, \dots, a_n$ be integers.

- $p \perp a$ iff $p \nmid a$.
- If $p \mid a_1 \cdots a_n$, then $p \mid a_i$ for at least one a_i .

Square-Free. Positive integer n is square-free iff:

- n is not divisible by any perfect square besides 1.
- or equivalently: n is 1 or a prod. of distinct primes.

Primes Under 100.

- from 1 to 20 are: 2, 3, 5, 7, 11, 13, 17, 19
- from 20 to 40 are: 23, 29, 31, 37
- from 40 to 60 are: 41, 43, 47, 53, 59
- from 60 to 80 are: 61, 67, 71, 73, 79
- from 80 to 100 are: 83, 89, 97

Dirichlet's Theorem. Let $a \perp b$ be given.

- There are infinitely many primes of the form $an + b$ where n is a positive integer.

Fixed Modulus. Let n be a positive integer and a, b, c, d be integers. All $\equiv$ below are modulo n .

- $a \equiv a$
- If $a \equiv b$, then $b \equiv a$.
- If $a \equiv b$ and $b \equiv c$, then $a \equiv c$.
- If $a \equiv b$ and $c \equiv d$, then $a + c \equiv b + d$ and $ac \equiv bd$.
- If $a \equiv b$, then $a + c \equiv b + c$ and $ac \equiv bc$.
- If $a \equiv b$, then $a^k \equiv b^k$ for any positive integer k .
- If $a \perp n$, then there exists integer a^{-1} so $aa^{-1} \equiv 1$.
- If $a \perp n$, then $ax \equiv b$ has a unique solution mod n .
- If $a \equiv b$, then $\gcd(a, n) = \gcd(b, n)$.

Changing Moduli. Let k, m, n be positive integers and a, b be integers.

- $a \equiv b \pmod{n}$ iff $ka \equiv kb \pmod{kn}$
- if $a \equiv b \pmod{n}$ and $d \mid n$, then $a \equiv b \pmod{d}$
- if $m \perp n$, then $a \equiv b \pmod{mn}$ if and only if both:

$$a \equiv b \pmod{m} \text{ and } a \equiv b \pmod{n}$$

Chinese Remainder Theorem. Let $n_1, \dots, n_r$ be relatively prime and $a_1, \dots, a_r$ be integers. Let $n = n_1 \dots n_r$. Then:

$$x \equiv a_1 \pmod{n_1}, \dots, x \equiv a_r \pmod{n_r}$$

is solved by $x = a_1 x_1 N_1 + \dots + a_r x_r N_r$ where:

$$N_k = n/n_k \text{ and } N_k x_k \equiv 1 \pmod{n_k}$$

This solution is unique modulo n .

Fermat's Theorem. Let p be prime and a an integer.

- $a^p \equiv a \pmod{p}$

Pseudoprimes. Let n be a composite integer.

- n is absolute pseudoprime iff always $a^n \equiv a \pmod{n}$
 - absolute pseudoprimes are square-free
-

Wilson's Theorem. Let n be an integer.

- n prime if and only if $(n-1)! \equiv -1 \pmod{n}$
-

Square-Root of -1 . Let p be an odd prime.

- $x^2 \equiv -1 \pmod{p}$ is solvable iff $p \equiv 1 \pmod{4}$, in which case a solution is given by $x = ([p-1]/2)!$.
-

Multiplicative Functions. Let f, g be arithmetic and m, n be positive integers and p be prime.

- multiplicative functions are uniquely determined by their values on powers of primes.
 - if f is multiplicative, and r is a real number, f^r is multiplicative
 - if f and g are multiplicative, then so is fg
 - $1, \text{id}, \tau, \sigma, \phi$ are multiplicative.
 - $\tau(p^n) = n+1, \sigma(p^n) = \frac{p^{n+1}-1}{p-1}, \phi(p^n) = p^n - p^{n-1}$
-

Sum Notation. Let f, g arithmetic and n integer.

- $$\sum_{\substack{d_1|m \\ d_2|n}} f(d_1)g(d_2) = \sum_{d_1|m} f(d_1) \sum_{d_2|n} g(d_2)$$