Older Adult (40-69 Years) Deaths by Symptom **Sierra Leone 2019–2022** (n = 4018)Malaria Infections Non-infections (n=1263)(n=3388)(n=5008)469 984 724 fever (2177, 23%) (37%)(29%)(14%)166 453 560 weightLoss (1179, 12%) (13%)(13%)(11%)102 311 580 breathingProblem (993, 10%) (8%)(9%)(12%)106 458 304 cough (868, 9%)(8%)(14%)(6%)59 197 569 abdominalProblem (825, 9%)(5%)(6%)(11%)218 118 279 vomit (615, 6%)(9%)(6%)(6%)47 209 332 chestPain (588, 6%) (4%)(6%)(7%)65 95 339 unconscious (499, 5%)(5%) (3%)(7%)23 31 348 Number paralysis of deaths (402, 4%) (2%)(1%)(7%)750 500 250 11 23 336 injury (370, 4%)(1%)(7%)(1%) 19 45 183 urinaryProblem (247, 3%)(2%)(4%)(1%) 158 25 64 looseStools (247, 3%)(2%)(5%)(1%) 61 160 6 oedema (227, 2%)(<1%) (2%)(3%)13 91 78 skinProblems (182, 2%)(1%)(3%)(2%)19 29 58 yellowEyes (106, 1%) (2%)(1%)(1%)7 8 36 convulsions (51, 1%) (1%)(<1%) (1%)

14

(<1%)

3

(<1%)

4

(<1%)

4

(<1%)

lumps (50, 1%)

pregnant (33, <1%)

32

(1%)

26

(1%)