Older Adult (40-69 Years) Deaths by Symptom **Sierra Leone 2019–2022** (n = 4018)Malaria Infections Non-infections (n=495)(n=1150)(n=2373)469 982 726 fever (2177, 54%) (95%)(85%)(31%)166 452 561 weightLoss (1179, 29%)(34%)(39%)(24%)102 308 583 breathingProblem (993, 25%) (21%)(27%)(25%)106 454 308 cough (868, 22%) (21%)(39%)(13%)197 59 569 abdominalProblem (825, 21%) (17%)(12%)(24%)118 218 279 vomit (615, 15%) (24%)(19%)(12%)47 207 334 chestPain (588, 15%) (9%)(18%)(14%)65 339 95 unconscious (499, 12%) (13%)(8%)(14%)23 31 348 Number paralysis (402, 10%) of deaths (5%) (3%)(15%)750 500 250 11 23 336 injury (370, 9%)(2%) (2%)(14%)19 45 183 urinaryProblem (247, 6%)(4%)(4%)(8%)25 158 64 looseStools (247, 6%)(5%)(14%) (3%)61 160 6 oedema (227, 6%)(1%)(5%)(7%)13 91 78 skinProblems (182, 5%)(3%)(8%)(3%)19 29 58 yellowEyes (106, 3%) (4%)(3%)(2%)7 8 36 convulsions (51, 1%)(1%)(1%)(2%)13 33 4 lumps (50, 1%)(1%) (1%)(1%)26 4 3 pregnant (33, 1%)(1%)(<1%) (1%)