Young Adult (15–39 Years) Deaths by Symptom Sierra Leone 2019–2022 (n = 2804)Non-infections Malaria Infections (n=487)(n=844)(n=1473)468 677 378 fever (1523, 23%) (96%)(80%)(26%)173 399 261 weightLoss (833, 12%) (36%)(18%)(47%) 76 220 431 abdominalProblem (727, 11%) (16%)(26%)(29%)146 217 200 vomit (563, 8%) (30%)(26%)(14%)109 109 330 cough (548, 8%) (7%)(22%)(39%)205 200 92 breathingProblem (497, 7%) (19%)(24%)(14%)20 396 6 injury (422, 6%)(2%) (1%)(27%)61 143 131 chestPain (335, 5%)(13%)(17%) (9%)57 81 158 Number unconscious of deaths (296, 4%)(12%) (10%)(11%)600 400 200 14 23 179 pregnant (216, 3%) (3%)(3%)(12%)24 104 30 looseStools (158, 2%)(12%)(2%)(5%) 16 69 46 skinProblems (131, 2%)(8%) (3%)(3%)65 5 35 oedema (105, 2%)(1%)(4%)(4%)17 23 64 urinaryProblem (104, 2%)(3%)(3%)(4%)8 22 41 paralysis (71, 1%) (2%)(3%)(3%)34 18 16 convulsions (68, 1%)(4%)(2%)(2%)18 15 28 yellowEyes (61, 1%)(4%)(2%)(2%)10 19 1 lumps (30, <1%)(<1%)(1%)(1%)