Adult (15-69 Years) Deaths by Symptom **Sierra Leone 2019–2022** (n = 7154)Malaria Infections Non-infectionsall_causes (n=752)(n=968) (n=1857) (n=3577) 176 91 384 651 abdominalProblem (12%)(18%)(21%) (18%)144 227 388 759 breathingProblem (19%) (23%)(21%) (21%)74 147 244 465 chestPain (10%)(15%) (13%)(13%)13 36 66 17 convulsions (2%) (2%) (1%) (2%)156 363 183 702 cough (21%)(38%)(10%)(20%)725 722 472 1919 fever (96%) (75%)(25%)(54%)33 143 38 214 looseStools (4%) (15%)(2%) (6%)5 6 19 30 lumps (1%) (1%) (1%) (1%) 29 110 145 6 oedema (1%) (3%) (6%) (4%)12 292 327 23 paralysis (3%) (1%) (16%) (9%)21 47 60 128 skinProblems (3%)(3%)(4%)(5%) 91 75 278 444 unconscious (12%)(8%) (15%) (12%)23 23 108 154 urinaryProblem (3%) (2%) (6%) (4%)201 205 610 204 vomit (27%) (21%) (11%) (17%)252 349 361 962 weightLoss (34%) (37%) (19%) (27%)27 12 40 79 yellowEyes (4%) (2%) (1%) (2%)

Number

of deaths

1500

1000

500