Older Adult (40-69 Years) Deaths by Symptom **Sierra Leone 2019–2022** (n = 4018)Non-infections Malaria Infections (n=2367)(n=495)(n=1156)469 984 724 fever (2177, 23%) (95%)(85%)(31%)166 453 560 weightLoss (1179, 12%) (34%)(39%)(24%)102 311 580 breathingProblem (993, 10%) (21%)(27%)(25%)106 458 304 cough (868, 9%)(21%)(40%)(13%)197 569 59 abdominalProblem (825, 9%)(12%)(17%)(24%)118 218 279 vomit (615, 6%)(24%)(19%)(12%)47 209 332 chestPain (588, 6%) (9%)(18%) (14%)65 339 95 unconscious (499, 5%)(13%)(8%)(14%)23 31 348 Number paralysis of deaths (402, 4%) (5%) (3%)(15%)750 500 250 11 23 336 injury (370, 4%)(2%)(2%)(14%)19 45 183 urinaryProblem (247, 3%)(4%)(4%)(8%)25 158 64 looseStools (247, 3%)(5%) (14%) (3%)61 160 6 oedema (227, 2%)(1%)(5%)(7%)13 91 78 skinProblems (182, 2%)(3%)(8%)(3%)19 29 58 yellowEyes (106, 1%) (4%)(3%)(2%)7 8 36 convulsions (51, 1%)(1%)(1%)(2%)14 32 4 lumps (50, 1%)(1%) (1%)(1%)26 4 3 pregnant (33, <1%)(1%)(<1%) (1%)