Female Child Symptom He

Malaria

(2451, 44%)

195

(8%)

138

(6%)

242

(10%)

5

(0%)

80

(3%)

3

(0%)

6

(0%)

2

(0%)

22

(1%)

50

(2%)

0

(0%)

82

(3%)

1

(0%)

220

(9%)

790

(32%)

16

(1%)

185

(8%)

165

(7%)

198

(8%)

17

(1%)

4

(0%)

30

(1%)

yellowEyes (347, 6%)

weightLoss (380, 7%)

urinaryProblem (28, 1%)

> unconscious (183, 3%)

> > ulcers (51, 1%)

skinRash (70, 1%)

paralysis (17, 0%)

oedema (84, 2%)

neckStiffness (105, 2%)

> lumps (6, 0%)

looseStools (250, 4%)

> injury (55, 1%)

headache (422, 8%)

fever

(1648, 30%)

(34, 1%)

(440, 8%)

cough (390, 7%)

convulsion (352, 6%)

chestPain (48, 1%)

> bleeding (13, 0%)

(125, 2%)

abdominalProblem

difficultySwallowing

difficultyBreathing

vomit (524, 9%)

emale Child S	symptom H	eatm	ap
	Number		
	of deaths ₀	200	400

0 600 Infections (2521, 45%)

121

(5%)

188

(7%)

241

(10%)

17

(1%)

59

(2%)

38

(2%)

48

(2%)

7

(0%)

34

(1%)

45

(2%)

3

(0%)

157

(6%)

1

(0%)

170

(7%)

792

(31%)

12

(0%)

210

(8%)

197

(8%)

108

(4%)

22

(1%)

3

(0%)

48

(2%)

eatr	nap

Non-infections

(600, 11%)

31

(5%)

54

(9%)

41

(7%)

6

(1%)

44

(7%)

10

(2%)

16

(3%)

8

(1%)

28

(5%)

10

(2%)

3

(0%)

11

(2%)

53

(9%)

32

(5%)

66

(11%)

6

(1%)

45

(8%)

28

(5%)

46

(8%)

9

(2%)

6

(1%)

47

(8%)