Young Adult (15–39 Years) Deaths by Symptom Sierra Leone 2019–2022 (n = 2804)Non-infections Malaria Infections (n=1473)(n=487)(n=844)468 677 378 fever (1523, 54%) (96%)(80%)(26%)173 399 261 weightLoss (833, 30%) (36%)(18%)(47%) 76 220 431 abdominalProblem (727, 26%)(16%)(26%)(29%)146 217 200 vomit (563, 20%) (30%)(26%)(14%)109 109 330 cough (548, 20%) (7%)(22%)(39%)205 200 92 breathingProblem (497, 18%) (19%)(24%)(14%)20 396 6 injury (422, 15%) (2%) (1%)(27%)61 143 131 chestPain (335, 12%) (13%)(17%) (9%)158 57 81 Number unconscious (296, 11%) of deaths (12%)(10%)(11%)600 400 200 14 23 179 pregnant (216, 8%)(3%)(3%)(12%)24 104 30 looseStools (158, 6%)(2%) (5%)(12%)16 69 46 skinProblems (131, 5%)(8%) (3%)(3%)65 5 35 oedema (105, 4%)(1%)(4%)(4%)17 23 64 urinaryProblem (104, 4%)(3%)(3%)(4%)8 22 41 paralysis (71, 3%) (2%)(3%)(3%)34 18 16 convulsions (68, 2%)(4%)(2%)(2%)18 15 28 yellowEyes (61, 2%) (4%)(2%)(2%)10 19 1 lumps (30, 1%) (<1%)(1%)(1%)