Older Adult (40-69 Years) Deaths by Symptom **Sierra Leone 2019–2022** (n = 4018)Malaria Infections Non-infections (n=1150) (n=2373)(n=495)29 19 58 yellowEyes (106, 3%)(4%)(3%)(2%)166 452 561 weightLoss (1179, 29%)(34%)(39%)(24%)118 218 279 vomit (615, 15%) (24%)(19%)(12%)19 45 183 urinaryProblem (247, 6%) (4%)(4%)(8%)95 65 339 unconscious (499, 12%) (8%)(13%)(14%)13 91 78 skinProblems (182, 5%)(3%)(8%)(3%)3 26 4 pregnant (33, 1%) (1%)(<1%) (1%)23 31 348 paralysis (402, 10%) (5%) (3%)(15%)160 6 61 Number oedema of deaths (227, 6%)(1%)(5%)(7%)4 13 33 lumps (50, 1%) (1%)(1%) (1%) 25 158 64 looseStools (247, 6%)(5%) (14%) (3%)11 23 336 injury (370, 9%)(2%)(2%)(14%)469 726 982 fever (2177, 54%) (85%)(31%)(95%)106 454 308 cough (868, 22%) (21%)(39%)(13%)7 8 36 convulsions (51, 1%) (1%)(1%)(2%)47 207 334 chestPain (588, 15%) (9%)(18%)(14%)102 308 583 breathingProblem (993, 25%) (21%)(27%)(25%)197 569 59 abdominalProblem (825, 21%) (12%)(17%)(24%)

750

500

250