Older Female Adult (40–69 Years) Deaths by Symptom **Sierra Leone 2019–2022** (n = 1714)Malaria Infections Non-infections (n=549)(n=1374)(n=2043)211 316 422 fever (949, 24%) (38%)(31%)(15%)75 177 230 weightLoss (482, 12%) (14%)(11%)(13%)40 108 246 breathingProblem (394, 10%) (7%)(8%)(12%)41 177 119 cough (337, 8%)(7%)(13%)(6%)24 80 191 abdominalProblem (295, 7%)(6%)(4%)(9%)114 48 96 vomit (258, 7%)(9%)(7%)(6%)15 69 159 chestPain (243, 6%)(3%)(5%)(8%)32 44 155 unconscious (231, 6%)(8%)(6%)(3%)13 13 158 Number paralysis of deaths (184, 5%)(2%)(1%)(8%)400 300 200 100 5 10 102 injury (117, 3%)(1%)(1%)(5%)70 10 24 looseStools (104, 3%)(2%)(5%) (1%) 5 23 67 oedema (95, 2%)(3%)(1%)(2%)7 45 37 skinProblems (89, 2%)(1%) (3%)(2%)10 47 7 urinaryProblem (64, 2%)(1%)(1%)(2%)20 8 14 yellowEyes (42, 1%) (1%)(1%)(1%)26 4 3 pregnant (33, 1%) (1%)(<1%) (1%)3 5 18 convulsions (26, 1%)(1%)(<1%) (1%)1 14 8 lumps (23, 1%) (<1%)(1%)(1%)