

Computer Assisted Verbal Autopsy: Comparing  
Large Language Models to Physicians for  
Assigning Causes to 6939 Deaths in Sierra Leone  
from 2019-2022

Richard Wen<sup>1\*</sup>, Anteneh Tesfaye Assalif<sup>1,2</sup>, Andy Sze-Heng Lee<sup>1</sup>,  
Rajeev Kamadod<sup>1</sup>, Asha Behdinan<sup>1</sup>, Ronald Carshon-Marsh<sup>1</sup>,  
Thomas Kai Sze Ng<sup>1</sup>, Patrick Brown<sup>1</sup>, Prabhat Jha<sup>1</sup>,  
Rashid Ansumana<sup>2</sup>

<sup>1\*</sup>Centre for Global Health Research, St. Michael's Hospital, Unity  
Health Toronto and University of Toronto, 30 Bond St, Toronto, M5B  
1W8, Ontario, Canada.

<sup>2</sup>School of Community Health Sciences, Njala University, Bo, Sierra  
Leone.

\*Corresponding author(s). E-mail(s): [richard.wen@utoronto.ca](mailto:richard.wen@utoronto.ca);  
Contributing authors: [antenehta@gmail.com](mailto:antenehta@gmail.com); [andylee@cs.toronto.edu](mailto:andylee@cs.toronto.edu);  
[rajeevk@kentropy.com](mailto:rajeevk@kentropy.com); [asha.behdinan@mail.utoronto.ca](mailto:asha.behdinan@mail.utoronto.ca);  
[ronald.carshonmarsh@mail.utoronto.ca](mailto:ronald.carshonmarsh@mail.utoronto.ca); [kaisze.ng@unityhealth.to](mailto:kaisze.ng@unityhealth.to);  
[patrick.brown@utoronto.ca](mailto:patrick.brown@utoronto.ca); [prabhat.jha@utoronto.ca](mailto:prabhat.jha@utoronto.ca);  
[rashidansumana@gmail.com](mailto:rashidansumana@gmail.com);

**Abstract**

**Background:** Verbal Autopsies (VAs) collect data on deaths and their causes  
outside of traditional hospital settings to provide representative Causes of Death  
(CODs). Current computer models for COD assignment in VAs perform similar  
to physicians at the population level, but poorly at the individual level, due to  
focuses on questionnaire data and neglecting free text from narratives. Recently, a

large language model called ChatGPT-4 demonstrated human-level performance on professional and academic exams. ChatGPT-4 shows promise in assigning CODs similar to physicians, but has yet been examined for assigning CODs using VA narratives.

**Methods:** 6939 VA records in Sierra Leone from 2019 to 2022 were used to compare four computer models, GPT-3.5, GPT-4, InterVA-5, and InSilicoVA, to physician COD assignment at population and individual levels. Narratives were used for GPT-3.5/4, while questionnaires were used for InterVA-5/InSilicoVA. COD assignments were grouped into general COD categories consisting of 19, 10, and 7 categories for adult, child, and neonatal age groups. Cause Specific Mortality Fraction (CSMF) accuracy and Partial Corrected Concordance (PCCC) were used to compare models to physicians at population and individual levels respectively. CSMF and PCCC were evaluated overall and by COD, age group, and age.

**Results:** GPT-4 had the best performance overall (0.61 PCCC), followed by GPT-3.5 (0.56 PCCC), and InSilicoVA/InterVA-5 (0.44 PCCC). GPT-4 had the best performance for adult and neonatal records (0.64 and 0.58 PCCC), with GPT-3.5 for child records (0.54 PCCC). All models' performances trended upwards from 1 month to 14 years ( $\sim 0.1$ -0.75 PCCC) and downwards from 15-69 years ( $\sim 0.7$ -0.35) of age. GPT-4, GPT-3.5, and InSilicoVA had the highest performances for 17, 9, and 4 of all 30 CODs respectively. At the population level, all models had CSMF accuracies between 0.74-0.79.

**Conclusion:** All models performed similarly at the population level, while GPT-3.5/4 and InSilicoVA performed better for some CODs at the individual level. GPT models made improvements over InSilicoVA and InterVA-5. Our research lays the foundation for future work in computer assisted VA, where physicians utilize alternative COD assignments from computer models to help reduce ill-defined codes and physician disagreement.

**Keywords:** Cause of Death, Physician Coding, Verbal Autopsy, GPT, AI, LLM

## 1 Background

In 2019, 41 million people died prematurely from noncommunicable diseases every year, accounting for 74% of all deaths globally [1]. Most of these deaths are preventable, but require adequate resource allocation, guided by evidence, to implement effective interventions and policies that target populations at risk [2]. Thus, reliable counts and diagnoses of deaths enable decision makers to identify populations at risk to save lives and reduce premature deaths worldwide [3-6]. However, many low-income countries do not have data on deaths or have registered less than half of the deaths in their

country, with an even fewer 8% of these registered deaths having a Cause of Death (COD) recorded [7]. To fill this gap in death registrations, an alternative method known as Verbal Autopsy (VA) is used to collect data on deaths and determine their likely causes at scale [8–10], outside of traditional healthcare facilities where over half of deaths occur at home [11].

VA involves two major components: survey and COD assignment [12–14]. In the survey component, trained lay surveyors interview those familiar with the deceased (e.g. living spouse, children, family, friends) to gather information using standardized questionnaires and open narratives. In the COD assignment component, physicians evaluate information available from the questionnaires and open narratives to assign probable CODs. This component has been criticized to be difficult to reproduce due to reliance on physician assignment [15–19]. As an alternative to physician assignment, computer models, such as InterVA [20] and InSilicoVA [17], have been studied to automatically assign CODs with performances close to physicians at the population level, but poor performances at the individual level [21–25]. These computer models often utilize data from the structured questionnaire, but often omit the free-text open narrative, which misses latent information, such as chronology or health-seeking behaviors, that may potentially help models perform better than using the questionnaire alone [26–28].

Recently, Large Language Models (LLM), leveraging massive datasets and deep learning approaches, have made advances in performing a variety of Natural Language Processing (NLP) tasks using free-text, such as question answering, code generation, and even medical diagnosis [29–32]. In 2022, a widely-available LLM called ChatGPT was released by OpenAI with capabilities of answering natural language text inquiries using training data up to September 2021. ChatGPT-3 was based on several Generative Pre-trained Transformer (GPT) models between 2018 to 2020, namely GPT-1 to


GPT-3, which had notable differences in training data sizes of 5 gigabytes to 45 terabytes from web sources that resulted in 117 million to 175 billion parameter models [33]. In March 2023, ChatGPT-4 was released with human-level performance on various professional and academic exams and benchmarks that outperformed ChatGPT-3 [34]. Given the limited usage of free-text open narratives in computer models for determining CODs, and recent advances in LLMs that leverage natural language text prompts, we conducted a study with Sierra Leone deaths from VA in 2019 to 2022 to compare four models, GPT-3.5, GPT-4, InterVA-5, and InSilicoVA, to physicians for determining CODs.

## 2 Methods

This study details the methods used to compare the COD assignment from four models, GPT-3.5, GPT-4, InterVA-5, and InSilicoVA, to physicians as seen in Figure 1. The initial VA data was filtered for physician agreed records as described in Section 2.1. Section 2.2 describes the input and output of the four models for COD assignment, while section 2.3 details the performance evaluation of the models relative to physicians using population and individual level metrics. See Appendix A for additional details on the methods used in this study.

### 2.1 Verbal Autopsy (VA) Data

Initially, 11,920 records from the HEAL-SL study [35, 36] were collected from dual-coded EVA, where each record was randomly coded by two different physicians that assigned CODs as International Classification of Diseases Revision 10 (ICD-10) codes [37]. For each record, two codes were assigned by two different randomly selected physicians, where codes were evaluated for agreement using Central Medical Evaluation Agreement 10 (CMEA-10) codes. CMEA-10 groups a range of similar ICD-10 codes together, where if they are in agreement if they are within the same group [38]



**Fig. 1** Study methods.

(see Additional File 2). When codes were not in agreement, a record enters the reconciliation phase, where the two physicians were provided reasoning and initial codes from each other to: (1) keep their initial code (2) assign the other physician's code or (3) assign a new code. If codes were not in agreement after the reconciliation phase, a record enters the adjudication phase, where a third senior physician evaluates both physicians' reasoning and codes before and after reconciliation, and assigns a final code based on their evaluation.

Since computer models were compared to physicians in this study, there was more certainty that COD assignments agreed by both physicians were representative of physician assignment than when they disagreed [18, 39, 40]. Thus, we used 6942 physician agreed records of the 11,920 total records. For better comparison, we standardized all codes to CGHR-10 codes (see Additional File 1) that generalized ICD-10 codes into

19, 10, and 7 categories for the adult (12 to 69 years), child (28 days to 11 years), and neonatal (under 28 days) age groups. After conversion, a final total of 6939 physician agreed records (3826 adult, 2636 child, and 477 neonatal) were used for modelling and performance evaluation. See Appendix A.1 for further details on data preprocessing and Tables A1 and A2 for COD and age range distributions of the physician agreed records.

## 2.2 Modelling

Four computer models were used to assign COD for each of the 6939 physician agreed records: GPT-3.5, GPT-4, InterVA-5, and InSilicoVA. InterVA-5 and InSilicoVA are widely used and studied standard statistical models [13, 21, 22, 24, 25, 41, 42] for COD assignment in VAs under the openVA framework [43]. InterVA-5 applies Bayesian probabilistic modelling [44] using a set of standardized symptoms from reports and related conditional probabilities from medical experts to assign CODs based on the highest probability [20, 45]. InSilicoVA improves upon InterVA (e.g. comparable probabilities across individuals, measures of uncertainty, and inclusion of additional data sources) with a hierarchical Bayesian framework and Markov Chain Monte Carlo (MCMC) simulations [46–48] to incorporate multiple sources of uncertainty for assigning CODs based on the highest probability [17]. GPT-3.5 [49] and GPT-4 [34] are LLMs that utilize deep neural networks with transformer architectures [50] and reinforcement learning from human feedback [51–54] to follow instructions from prompts and provide human-level responses, with known differences in GPT-4 possessing multimodal capabilities (e.g. image/voice input/output), more recent training data, and improved responses compared to ChatGPT-3 [33].

For GPT-3.5 and GPT-4, the following user prompt was used to instruct each model to produce COD assignments as ICD-10 codes, where `<age>` and `<sex>` from the questionnaire, and `<narrative>` from the narratives, were replaced with values from the data:

Determine the underlying cause of death and provide the most  
probable ICD–10 code for a verbal autopsy narrative of a <age>  
years old <sex> death in Sierra Leone: <narrative>

For InterVA-5 and InSilicoVA, the standardized questionnaire data from EVA were  
converted into OpenVA format [43], before being used as input for each model to pro-  
duce COD assignments as WHO VA 2016 codes [55]. All model outputs were converted  
to CGHR-10 codes to evaluate performances of models for COD assignment relative  
to physicians. See Appendix A.2 for additional details regarding input parameters,  
output data, and code conversions for each model.

## 2.3 Performance Evaluation

The performance of the four models were evaluated with metrics at the population and  
individual level by comparing their CGHR-10 COD outputs for 6939 records. Cause  
Specific Mortality Fraction (CSMF) accuracy was used to evaluate models on the  
population level (see Appendix A.3.1), while Partial Chance Corrected Concordance  
(PCCC) was used to evaluate models on the individual level (see Appendix A.3.2) [56].  
Both CSMF accuracy and PCCC metrics are between 0 and 1 with 0 indicating low  
performance and 1 indicating perfect performance at the population and individual  
level respectively. As model performance can vary across ages and specific causes  
[41, 42, 57], the CSMF accuracy and PCCC metrics were compared for each model  
overall, by age group (adult, child, neonatal), by CGHR-10 COD codes, and across  
ages. For each of the adult and child age groups, metrics were calculated for five-year  
ages for records with ages at death of one-year or older and five-month ages for 28 days  
or older. For the neonatal age group, the ages of 0-6 days and 7-27 days were used.  
See Appendix A.3 for more details on performance metrics and evaluation strategy  
for comparing each model.

## 3 Results


This section details the performance results of GPT-3.5, GPT-4, InterVA-5, and InSilicoVA models for assigning CGHR-10 CODs after applying the methods in Section 2. GPT-4 performed the best overall at 0.61 PCCC followed by GPT-3.5 at 0.56 PCCC. GPT-4 also had the highest PCCC for most ages and CODs across the adult (12 to 69 years), child (28 days to 11 years), and neonatal (under 28 days) age groups with GPT-3.5, InterVA-5, and InSilicoVA having higher PCCC values for a few ages and CODs. Overall performance results are seen in Section 3.1, and performance by adult, child, and neonatal records are seen in Sections 3.2, 3.3, and 3.4 respectively.

### 3.1 Overall Performance


Of all 6939 records, GPT-4 (0.61 PCCC) had the highest individual performance followed by GPT-3.5 (0.56 PCCC), InSilicoVA (0.44 PCCC), and InterVA-5 (0.44 PCCC) (Figure 2). GPT-3.5 and GPT-4 had improvements ranging from 0.14-0.18 PCCC over InSilicoVA and InterVA-5, while GPT-4 slightly improved over GPT-3.5 by 0.05 PCCC. Population level performances were similar for all models (0.74-0.79 CSMF). Figure 3 shows the PCCC performance across three age groups (adult, child, and neonate). GPT-4 had the best individual performance for adult and neonatal records (0.64 and 0.58 PCCC), while GPT-3.5 had the best performance for child records (0.54 PCCC) with GPT-4 performing slightly worse (0.51 PCCC). InSilicoVA and InterVA-5 performed the worse for adult and child records ( $\leq 0.5$  PCCC), while GPT-3.5 performed the worse for neonatal records (0.42 PCCC). Across ages, all models followed a similar pattern in individual performance (Figure 4). PCCC trended upwards for 1 month to 14 years ( $\sim 0.1$ -0.75), and downwards for ages 15 to 69 years ( $\sim 0.7$ -0.35). The highest and lowest performances were observed for ages 10-29 years ( $\sim$ ) and 1-11 months respectively. Performances varied more across models for ages 0




days to 5 years, while less variation was seen between GPT-3.5 and GPT-4, as well as InSilicoVA and InterVA-5, from 5 to 69 years.



**Fig. 2** Overall model performance.




**Fig. 3** Model performance by age group.



**Fig. 4** Model performance by age range.

### 3.2 Performance for 3826 Adult Records (12 to 69 years)

Figure 5 shows model performance by PCCC across 17 adult CODs excluding suicide due to low sample size ( $n=3$ ,  $<1\%$ ). GPT-4 had the highest individual performance for 10 of 17 CODs (0.35 to 0.99 PCCC), GPT-3.5 for 5 CODs (0.43-0.94 PCCC), and InSilicoVA for 2 CODs (0.71 and 0.84 PCCC). InterVA-5 had the lowest performance for 8 of 17 CODs (0-0.79 PCCC), InSilicoVA for 6 CODs (0.01-0.41 PCCC), and GPT-3.5 for 2 CODs (0.38 and 0.53 PCCC). GPT-3.5/4 models improved over InSilicoVA/InterVA-5 the most for chronic respiratory diseases (0.74-0.94 PCCC difference), and the least for Malaria (0.09-0.17 PCCC difference). All models had  $>0.7$  PCCC for maternal conditions (0.79-0.99 PCCC), while  $<0.5$  PCCC for unspecified infections, malaria, and ill-defined CODs. GPT-4 had performance improvements  $>0.2$  PCCC compared to all other models for cancers (+0.25-0.36 PCCC), stroke (+0.27-0.45 PCCC), and diarrhoeal diseases (+0.37-0.51 PCCC), while GPT-3.5 had similar improvements for liver and alcohol related diseases (+0.27-0.52 PCCC).




**Fig. 5** Model performance for adult records by COD.

### 3.3 Performance for 2636 Child Records (28 Days to 11 Years)

Figure 6 presents individual performances for each of the models by 8 child CODs, excluding congenital anomalies due to low sample size (n=1, <1%). GPT-4 had the

507 highest individual performance for 4 of 8 CODs (0.65-0.94 PCCC), GPT-3.5 for 3  
 508 CODs (0.44-0.88 PCCC), and InSilicoVA for 1 COD (0.78 PCCC). InterVA-5 had  
 509 the lowest performance for 4 of 8 CODs (0.09-0.79 PCCC), InSilicoVA for 3 CODs  
 510 (0-0.35 PCCC), and GPT-3.5 for 1 COD (0.58 PCCC). All models had >0.7 PCCC  
 513 for injuries (0.79-0.94 PCCC), and <0.6 PCCC for malaria (0.35-0.54 PCCC) and  
 514 other infections (0.29-0.44 PCCC). GPT-4 had improvements >0.3 PCCC compared  
 515 to other models for ill-defined CODs (+0.38-0.65 PCCC), and larger improvements  
 516 over other models for injuries (+0.11-0.15 compared to +0.01-0.04 PCCC).  
 517  
 518  
 519




550 **Fig. 6** Model performance for child records by COD.

551

552

### 3.4 Performance for 477 Neonatal Records (Under 28 Days)

Model performance across 5 neonatal CODs, excluding congenital anomalies (n=2, <1%) and other (n=5, 1%) due to small sample sizes is shown in Figure 7. GPT-4 had the highest individual performance for 3 of 5 CODs (0.39-0.71 PCCC), GPT-3.5 for 1 COD (0.57 PCCC), and InSilicoVA for 1 COD (0.86 PCCC). GPT-3.5 had the lowest performance for 3 of 5 CODs (0-0.13 PCCC) and InterVA-5 for 2 CODs (0.01 and 0.48 PCCC). All models had similar performance for stillbirth deaths (0.48-0.57 PCCC), while only GPT-4 had a PCCC >0 PCCC. InSilicoVA had improvements over all other models for neonatal infection deaths (+0.18-0.73 PCCC).



**Fig. 7** Model performance for neonatal records by COD.

## 4 Discussion

This section discusses and summarizes the results from Section 3. Advantages and disadvantages of using GPT-3.5, GPT-4, InterVA-5, and InSilicoVA models for assigning

599 CODs are discussed in Sections 4.1 and 4.2. Limitations of the study are mentioned  
600 in Section 4.3, while opportunities and future work are detailed in Section 4.4.

602

603

604

## 4.1 Advantages

605

606 This section identifies the advantages of models for assigning CODs. Section 4.1.1

607

608 details the application of models for particular CODs and ages. Section 4.1.3 details

609

610 the resource efficiency of computer models for assisting in physician COD assign-

611

612 ment. Section 4.1.4 notes the strength of using natural language text in GPT models  
613 compared to structured questionnaire data for physician COD assignment.

614

615

### 4.1.1 Cause-specific Models

616

617 At the population level, overall performances for all models were similar to physicians  
618 (0.74-0.79 CSMF), indicating potential for adequately estimating COD distributions

619

620 for large populations. Although all models did not perform well for all records at

621

622 the individual level (0.44-0.61 PCCC), several models performed well for certain  
623 CODs (0-0.99 PCCC). For most CODs, GPT-3.5/GPT-4 performed better than

624

625 InSilicoVA/InterVA-5 (top PCCC for 15 of 17, 7 of 8, and 4 of 5 adult, child, and  
626 neonatal CODs respectively), while InSilicoVA performed better for particular CODs

627

628 (road and transport injuries, tuberculosis, pneumonia, and neonatal infections with  
629 0.84, 0.71, 0.78, and 0.86 PCCC respectively). For CODs with high performance (e.g.

630

631 GPT-3.5/4 with 0.91-0.99 PCCC for maternal conditions, chronic respiratory disease,  
632 diabetes melitus, and cancers, InSilicoVA with 0.84 and 0.86 PCCC for road and

633

634 transport injuries, and neonatal infections), the results suggest that GPT-3.5/4 and  
635 InSilicoVA may assign CODs that are very similar to physicians. Thus, it may be ben-

636

637 eficial to evaluate performance at the COD level, and apply a combination of models  
638 that perform well in comparison to physicians for each COD. For example, different

639

640 models perform well for various leading CODs as seen in Table 1 [58, 59].  
641  
642  
643  
644

**Table 1** Leading causes globally in 2021 and most relevant models.

Leading Causes of Death (~53% of 68M deaths) <sup>1</sup>	Deaths (% of 68M) <sup>2</sup>	Best Model(s)	PCCC
Ischaemic heart disease	9M (13%)	GPT-4	0.65 (n=168)
Stroke	7M (10%)	GPT-4	0.77 (n=331)
Cancers	4.3M (4%)	GPT-4	0.91 (n=49)
Lower respiratory infections	2.4M (3%)	GPT-3.5/4	0.78 (n=180) <sup>3</sup>
Diabetes mellitus	1.6M (2%)	GPT-3.5	0.92 (n=27)
Tuberculosis	1.4M (2%)	InSilicoVA	0.71 (n=171)
Hypertensive heart disease	1.4M (2%)	GPT-4	0.78 (n=107) <sup>4</sup>
Cirrhosis of the liver	1.3M (2%)	GPT-3.5	0.78 (n=128) <sup>5</sup>
Diarrhoeal diseases	1.2M (2%)	GPT-4	0.68 (n=205)
Road injury	1,183 (2%)	InSilicoVA	0.84 (n=357)
Preterm birth complications	0.9M (1%)	GPT-4	0.71 (n=73)
Falls	0.7M (1%)	GPT-4	0.89 (n=492) <sup>6</sup>

<sup>1</sup>COVID-19, kidney disease, alzheimer disease, other dementias, and self-harm were excluded as a relevant CGHR-10 code was not present. Trachea, bronchus, lung, colon, rectum, stomach, and breast cancers were generalized into cancers.

<sup>2</sup>Percentage of ~68 Million (M) deaths globally. Numbers are rounded.

<sup>3</sup>Mean of chronic and acute respiratory infections.

<sup>4</sup>Derived from other cardiovascular diseases.

<sup>5</sup>Derived from liver and alcohol related diseases.

<sup>6</sup>Mean of adult and child injuries.

#### 4.1.2 Age-specific Performance Patterns

Across ages, all models followed a similar upward trend from 6 months to 14 years of age, and a downward trend from 15-69 years with GPT models having higher performance than InSilicoVA/InterVA-5 models, while more mixed trends were observed from 0 days to 5 months (recall Figure 4). For adult ages, performance generally decreased as age increased, which suggested that models had difficult assigning CODs for older than younger adults with some improvements after the age of 59. For child and neonatal ages, the performance improved drastically as the age increased after 5 months, suggesting less difficulty in COD assignment when children and neonates are more developed. As the models did not perform particularly well ( $\geq 0.8$  PCCC) for any specific five-year age range, it is not recommended to apply specific models that target cases by age. However, the patterns of increases and decreases of performance in

691 relation to age provide valuable insight for comparison to expected physician diagno-  
692 sis patterns in well-studied medical literature and knowledge. For example, it may be  
693 expected that physicians are more uncertain in diagnosing diseases that are prevalent  
694 in neonatal patients [60, 61], which are present in our findings from Figure 4.  
695

### 696 697 698 **4.1.3 Scalability and Availability** 699

700  
701 The models in this study can assist physicians in assigning CODs in a variety of ways  
702 due to low costs and speed of COD assignment. Similar to differential diagnoses, GPT  
703 and InSilicoVA models offer alternative COD assignments for physicians to consider  
704 [39], which can potentially help lower the number of records with ill-defined causes or  
705 reduce disagreement between physicians. At the time of this study, running GPT-3.5  
706 cost ~\$1.6 USD (\$0.5 per one million tokens), GPT-4 cost ~\$115 USD (\$30 per one  
707 million tokens), and InSilicoVA was cost free on 6939 records [62]. These costs were  
708 lower than physicians (e.g. less than \$3 USD per house in India [15, 16]), while it is  
709 possible to code over 10,000 records in under a day. When physicians are unavailable,  
710 GPT and InSilicoVA models can be a cost-efficient alternative to code large amounts  
711 of records for population estimates of CODs. However, it is recommended to apply  
712 these models only for certain CODs where models perform well, such as in Table 1. In  
713 addition, these models can also help divert physician resources to cases that are more  
714 difficult to code or require more attention. For example, physicians can validate cases  
715 where models performed well (e.g. maternal conditions at 0.79-0.99 PCCC), while  
716 spending more time on cases where models performed poorly (e.g. acute respiratory  
717 infections at 0.25-0.61 PCCC).  
718  
719  
720  
721  
722  
723  
724  
725  
726  
727  
728

### 729 730 **4.1.4 Natural Language Input and Output** 731

732 Training data was not required to assign CODs for all models, which allowed appli-  
733 cation without domain expertise or supplying training datasets. The main advantage  
734 to GPT-3.5/4 was the use of natural language text as input and output. Compared  
735  
736



to InterVA-5 and InSilicoVA, GPT models were able to assign COD codes in ICD-10 standard, as physicians do, and potentially assign CODs in more broad categories depending on the prompts. In comparison, InterVA-5 and InSilicoVA relied on structured input and output data from WHO VA 2016 questionnaires, and assigned CODs in WHO VA 2016 codes only. This required that these codes and forms be maintained with conversions between different form (e.g. WHO VA 2012 to WHO VA 2016) and code standards (e.g. WHO VA 2016 to ICD-10), which reduces interoperability and comparability with other incompatible models. GPT models did not require strict formats for training and testing data, which can capture latent and more ambiguous patterns (e.g. health-seeking behaviours and social issues) outside the scope of WHO VA codes and forms [26, 28]. For example, GPT-3.5/4 had higher performance (+0.35-0.65 PCCC) than InterVA-5 and InSilicoVA for ambiguous ill-defined records across age groups. GPT models also performed better (+0.11-0.61 PCCC) on CODs with a rarer occurrence, such as nutritional deficiencies (n=11) and diabetes mellitus (n=27). Rarer CODs may be more difficult to capture by questionnaire due to lack of sample data, but it may possibly have richer contextual information from articles, web sources, or books that offer knowledge for GPT models to leverage.

## 4.2 Disadvantages

This section discusses the disadvantages of GPT models for COD assignment. Section 4.2.1 identifies issues in reproducing GPT outputs for repeated runs on the same records and lack of up-to-date information, while Section 4.2.2 discusses the resource intensive infrastructure required by GPT and its relation to data privacy.

### 4.2.1 Reproducibility and Timeliness

Recall that the GPT models in this study had the temperature parameter set to 0 for more reproducible and reliable results. A short experiment in Appendix B revealed that GPT-3.5 assigns the same COD for the same record only more than 60% of the

time, based on repeated runs on a sample of 100 records. This suggests that GPT models do not always reliably assign identical CODs for the same case on multiple runs, which may pose issues in reproducibility and reliability. For example, GPT models may achieve correct COD assignments solely due to random chance, but are difficult to test with large numbers (e.g. 10,000) of reruns due to costs (e.g. costs increased 10 fold per record when rerun 10 times). In comparison, InterVA-5 and InSilicoVA are open source and free, allowing a large number of reruns without incurring additional fees. In addition, InterVA-5 and InSilicoVA assign CODs and provide probabilities for each alternative COD, which offers more reproducible and reliable COD assignments despite lower performance overall. Lastly, a major disadvantage in all models was that they were trained on historical data up to particular points in time, which may not utilize the most up-to-date data available (e.g. latest online articles, social media, or books for GPT models). Emergent diseases (e.g. COVID-19) and changes in distributions (e.g. outbreaks) may not be caught by these models depending on how often they are updated.

#### **4.2.2 Infrastructure and Data Privacy**

GPT-3.5 and GPT-4 models required large computing infrastructure to train and run, which was not possible to run on local computers, or setup due to costs and ownership of the models. This poses issues with data privacy as sensitive data (e.g. identifying information) need to be sent to company servers, which can be collected by companies (e.g. OpenAI) and misused [63]. For example, in our study, GPT models use prompts, which contain the narrative data, to assign CODs, and the data in these prompts may be unknowingly collected and misused by companies (e.g. companies) or their users (e.g. malicious prompts) to identify participants or leak sensitive sensitive data [64, 65]. In contrast, InterVA-5 and InSilicoVA can be run on local computers, which allows data to stay with the owner to protect data privacy, without reliance on external services.

### 4.3 Limitations

This section identifies limitations in this research in the context of GPT models. Section 4.3.1 identifies the omission of ICD-10 performance evaluations. Section 4.3.2 mentions the need for parameter tuning and evaluation of consistency and multiple COD assignments.

#### 4.3.1 ICD-10 Evaluation and Low Sample Sizes

For the scope of this study, all models were evaluated for their performance in broad CGHR-10 COD categories as opposed to more specific ICD-10 codes. However, in practical cases, physicians assign more specific ICD-10 codes rather than broader COD categories. InterVA-5 and InSilicoVA assigned broader WHO VA codes, and were unable to assign ICD-10 codes, as the number of cases for specific ICD-10 codes are often low and inadequate for training statistical models. In relation, some broader CGHR-10 CODs were even removed for performance evaluation as <10 cases were captured (e.g. congenital anomalies, suicide). Although GPT models were able to assign ICD-10 codes, lower performance may be expected as even physicians do not agree completely on ICD-10 codes, noted that broader categories (CMEA-10 codes in Additional file 2) were used to assign equivalency or agreement.

#### 4.3.2 Model Tuning, Consistency, and Multiple Outputs

GPT-3.5 and GPT-4 models used default parameters with the exception of setting the temperature to 0 for more consistent results. However, the temperature and other model settings may be adjusted to possibly improve performance for GPT models [66]. This was not examined as sensitivity analyses on model parameters are costly across multiple reruns, noted in Section 4.2.1, which is required when testing various parameter settings. In addition, GPT models may possibly produce inconsistent results even with the temperature set to 0. Thus, it is important to also test the reliability

875 and consistency of GPT outputs to avoid coincidental results due to randomness [67–  
876 69]. InterVA-5 and InSilicoVA were able to provide multiple COD assignments with  
877 probabilities for each COD. GPT models can be prompted to produce more than one  
878 COD assignment, but was not explored in this study as we only evaluated the most  
880 probable COD. This may be useful to evaluate the performance of multiple alternative  
881 COD assignments, which may provide additional diagnoses that have a higher chance  
882 of being similar to physician assignment, and better reflect causes leading to death  
883 [19].  
884  
885  
886  
887

## 889 4.4 Opportunities

891 This section discusses research opportunities to improve GPT models for assigning  
892 CODs. Section 4.4.1 discusses the potential to improve GPT models with prompt  
893 engineering and exploration of misclassified records, while Section 4.4.2 describes the  
894 application of GPT models for improving household surveys for better data quality.  
895 Section 4.4.3 identifies an opportunity to integrate GPT, InterVA-5, and InSilicoVA  
896 models into VA systems for improving physician COD assignment.  
897  
898  
899

### 903 4.4.1 Prompt Engineering and Custom Models

904 Prompt engineering, the design of prompts to guide GPT models for better results [70],  
905 presents an important research opportunity that may improve performance of GPT  
906 models for COD assignment. An example exploration was conducted in Appendix C on  
907 misclassified GPT-4 records for neonatal infections, which found potential issues with  
908 the categorization of CGHR-10 codes, order of information in narratives, and guide-  
909 lines of COD assignments. An analysis of misclassified records with domain experts  
910 (e.g. physicians, specialists) may yield insights on adjusting prompts to assign more  
911 correct CODs, or apply more relevant broad COD categories for evaluation. In addi-  
912 tion, subsequent prompts, data, and examples can be used to include correctional  
913 instructions and refine results, while additional information from the questionnaire and  
914  
915  
916  
917  
918  
919  
920

physician VA manuals can provide contextual information (e.g. retrieval augmented generation [71]) for further performance improvements [72]. Sensitivity analyses may be conducted to assess the effects on performance and consistency of results from modified prompts on a COD basis. GPT models may also be customized to specific domains or contexts, where objectives, behaviours, extra data, privacy, and evaluation tests can be adjusted to produce custom models that perform better in targeted domains or circumstances (e.g. custom models for particular CODs) [73].


#### 4.4.2 Guided and Monitored Household Surveys

Recall that VAs involve surveyors that visit households to gather information about the deceased from their family, next-of-kin, friend, or community. Although standard questionnaires are used during this visit, there is significant information, containing latent patterns, from the narrative that is not always captured by the questionnaire [26, 28]. These narratives often require a human connection between the surveyor and household members, where surveyor characteristics vary in social ability, cultural understanding, emotional capacity, and medical knowledge that affect the quality and bias of narratives [19, 74]. GPT models may help guide surveyors during VA interviews to probe households for better narrative information by generating and suggesting better questions, or providing questions that may have been missed by the surveyors. In addition, as models can assign CODs on-demand, there is potential for models to provide immediate COD estimates during the data collection process to monitor data quality on-demand (e.g. comparing estimated to expected COD distributions for known areas as quality checks).

#### 4.4.3 Computer Assisted Verbal Autopsy

Our study lays the foundation for the integration of GPT, InterVA-5, and InSilicoVA models into VA systems to assist physicians in COD assignment. In dual-coded VA systems (described in Section 2.1), two physicians are randomly assigned to each record

and require second inspections of each other’s assignment (reconciliation) and evaluation by a third more senior physician if their assignments do not agree. As mentioned in Section 4.1.3, suggestion of alternative assignments from GPT and InSilicoVA models potentially reduces the disagreement between physicians, and ill-defined records, while allowing physicians to focus on more difficult records. Thus, model suggestions can be integrated into VA systems by presenting COD suggestions to physicians after their initial COD assignment, which allows them to consider alternative assignments and possibly revise their assignments based on the suggestions. At step 2 in Figure 8, GPT, InterVA-5, and InSilicoVA models can suggest COD assignments to consider, providing the option in step 2b to revise or proceed with their initial assignment. Our future work will be a first step in computer assisted verbal autopsy, assessing the effects of these model suggestions on improve VA data quality (e.g. increase in agreed records, reduction of ill-defined deaths). In preparation, we have integrated GPT-4, InterVA-5, and InSilicoVA model suggestions into our on-going HEAL-SL study after survey round 2 [35] with goals of increasing physician agreement and reducing ill-defined COD assignments.



**Fig. 8** Model suggestions integrated in the physician assignment process.

## 5 Conclusion

This study evaluates the performance of GPT-3.5, GPT-4, InterVA-5, and InSilicoVA models compared to physicians for assigning CODs for 6939 VA records in Sierra Leone (2019-2022). At the population level, all models were similar (0.74-0.79 CSMF accuracy). At the individual level, GPT-4 had the best performance (0.61 PCCC), followed by GPT-3.5 (0.58 PCCC), and InSilicoVA/InterVA-5 (0.44 PCCC). Across CODs, GPT-4 had performed best for 10 of 17 adult, 4 of 8 child, and 3 of 5 neonatal CODs, with GPT-3.5 for 5 adult, 3 child, and one neonatal CODs, and InSilicoVA for 2 adult, one child, and one neonatal CODs. Model performance increased ( $\sim 0.1$ -0.75 PCCC) as children and neonates developed (0 days to 14 years), and decreased ( $\sim 0.7$ -0.35) as adults aged (15 to 69 years). Thus, GPT and InSilicoVA models were comparable to physicians for several CODs, but not across ages. As performance varied across CODs and ages, it is advantageous to combine several models to target CODs that each model performs well for, and to compare age-related performance patterns in relation to physicians. In addition, all models were able to scale to a large number of records and were available on-demand in comparison to physicians, enabling COD estimation and alternative diagnoses in low resource or physician scarce scenarios. As GPT models operate on natural language, they are able to adapt to more loosely defined data structures (e.g. assign in different COD coding standards, provide reasoning, and use contextual information when samples are low), making them behave more similarly to physician assignment. However, GPT models do not provide reliable CODs on repeated assignment, and were limited to past training data, with large computing infrastructure requirements, leading to reproducibility issues in COD assignments, difficulty adapting to new or changing CODs, and data privacy issues. Limitations of this study included difficulty comparing ICD-10 codes directly due to incompatible COD outputs from each model and low sample sizes, difficulty in conducting sensitivity analyses for GPT models due to costs, and omitting evaluation of

multiple COD assignments due to study scope. We identified research opportunities in refining GPT models using prompt engineering and custom models for improving performance, guided household surveys to improve narrative quality, and future work in computer assisted VA, where GPT and other models will be used to assist physician COD assignment by offering multiple alternative assignments, with goals of increasing agreement on COD assignment and reducing ill-defined deaths. GPT-4, InterVA-5, and InSilicoVA has been integrated into future survey rounds of the HEAL-SL study from 2022 onwards, offering alternative COD assignments to assist physicians with second opinions. Future work in evaluating the effectiveness of computer assisted VA to reduce disagreements among physicians and ill-defined deaths will help support the advancement of more accurate and efficient VA systems across the world.

**Supplementary information.** Additional files were used to supplement this paper:

- Additional file 1: Centre for Global Health Research 10 (CGHR-10) codes. Codes grouping ICD-10 code ranges into generalized categories. (.csv)

- Additional file 2: Central Medical Evaluation Agreement 10 (CMEA-10) codes. ICD-10 code ranges considered in physician agreement. (.csv)

**Acknowledgments.** TBD.

## Declarations

## Funding

TBD.

## Competing interests

Not applicable.



<b>Ethics approval</b>	1105
Not applicable.	1106
	1107
	1108
	1109
<b>Consent for publication</b>	1110
	1111
Not applicable.	1112
	1113
	1114
<b>Availability of data and materials</b>	1115
	1116
The datasets supporting the conclusions of this article are included within the article	1117
(and its additional files), at <a href="https://openmortality.org">https://openmortality.org</a> (available upon request) and at	1118
<a href="https://github.com/cghr-toronto/healsl-gpt-paper">https://github.com/cghr-toronto/healsl-gpt-paper</a> . Verbal Autopsy (VA) and narra-	1119
tive data by age group and survey rounds 1 and 2 available at <a href="https://openmortality.org/dataset/heal-sl">https://openmortality.</a>	1120
<a href="https://openmortality.org/dataset/heal-sl">org/dataset/heal-sl</a> . Cause of death code mappings to convert between ICD-10,	1121
WVA-2016, and CGHR-10 codes available at <a href="https://openmortality.org/dataset/icd">https://openmortality.org/dataset/icd</a> .	1122
Model evaluation result files at <a href="https://github.com/cghr-toronto/healsl-gpt-paper/tree/main/data">https://github.com/cghr-toronto/healsl-gpt-paper/</a>	1123
<a href="https://github.com/cghr-toronto/healsl-gpt-paper/tree/main/data">tree/main/data</a> .	1124
	1125
	1126
	1127
	1128
	1129
	1130
	1131
<b>Code availability</b>	1132
	1133
All code for this paper is available at <a href="https://github.com/cghr-toronto/healsl-gpt-paper">https://github.com/cghr-toronto/</a>	1134
<a href="https://github.com/cghr-toronto/healsl-gpt-paper">healsl-gpt-paper</a> .	1135
	1136
	1137
	1138
<b>Authors' contributions</b>	1139
	1140
PJ and PB are the study Principal Investigators. ATA and RK implemented the data	1141
collection procedures. RW, and TKS processed, documented, and prepared the data.	1142
RW, ASL, and RK ran the models. RW wrote the paper and conducted the analysis.	1143
AB and RCM provided medical domain guidance and feedback. All authors reviewed	1144
the results and contributed to the report. All authors read and approved the final	1145
manuscript.	1146
	1147
	1148
	1149
	1150

## 1151 **Appendix A   Details on Methods**

1152

1153 This section provides additional details on the methods described in Section 2. An  
1154

1155 overview of the methods used in this study is seen in Figure A1 as a five-step process.

1156

1157 Section A.1 provides details on the preprocessed data used for modelling. Section A.2

1158 describes the data and parameter inputs and outputs for each model, while Section  
1159

1160 A.3 details the evaluation of model outputs at the individual and population level

1161

1162 across different CODs, age groups, and ages.

1163

1164

1165

1166

1167

1168

1169

1170

1171

1172

1173

1174

1175

1176

1177

1178

1179

1180

1181

1182

1183

1184

1185

1186

1187

1188

1189

1190

1191

1192

1193

1194

1195

1196




Fig. A1 Detailed study methods.


## 1243 **A.1 CGHR-10 Physician Agreed Records**

1244

1245 Initially, 11,920 records were collected from dual-coded EVA in the HEAL-SL study.  
1246  
1247 Physicians were able to assign CODs for 11,820 of the 11,920 records, where 100 of  
1248 these records could not be assigned a COD due to missing or inadequate information  
1249 (e.g. low quality narrative, data loss). The 11,820 physician coded records were further  
1250 filtered for records where both physicians agreed on the assigned codes (records that  
1251 were not reconciled or adjudicated) resulting in 6942 physician agreed records (based  
1252 on comparisons using CMEA-10 codes, see Additional File 2). The 6942 records were  
1253 converted into CGHR-10 codes (see Additional File 1) that generalized ICD-10 codes  
1254 into 19, 10, and 7 categories for the adult (12 to 69 years), child (28 days to 11  
1255 years), and neonatal (under 28 days) age groups. After conversion, a final total of  
1256 6939 physician agreed records (3826 adult, 2636 child, and 477 neonatal) were used  
1257 for modelling and performance evaluation, where three records were removed as their  
1258 ICD-10 codes did not have a matching CGHR-10 code.  
1259

1260 The 6939 physician agreed records were collected using VA from the HEAL-SL  
1261 study between 2019-2022, where records were collected using nation wide samples  
1262 across Sierra Leone provinces seen in Figure A2. More populous areas (e.g. southern  
1263 and north east provinces with ~197,000 and ~135,000 population respectively) had  
1264 more sampling areas versus less populous areas (e.g. north west and eastern provinces  
1265 with ~50,000 and ~69,000 people respectively). The distribution of the study data are  
1266 shown by CGHR-10 causes of death in Table A1. All age groups had relatively evenly  
1267 distributed female and male records (44-55% of 6939 records each). Across CODs,  
1268 there were noticeably more female records for cancers (65%), and maternal condi-  
1269 tions (100%), while more male records for chronic respiratory diseases (61%), other  
1270 noncommunicable diseases (61%), other injuries (77%), road and transport injuries  
1271 (71%), and tuberculosis (68%). Most records were coded by physicians as malaria for  
1272 adults (20%) and children (52%), and stillbirth (36%) and neonatal infections (21%)  
1273  
1274  
1275  
1276  
1277  
1278  
1279  
1280  
1281  
1282  
1283  
1284  
1285  
1286  
1287  
1288

for neonates. Suicide, congenital anomalies, nutritional deficiencies, and other had low sample sizes for each age group (<1% of total records for each age group). Table A2 shows the distribution of the study data by age. Across ages, there were more male records for 50-59 years (60-62%), while all other records had between 49-59% female and male records. Most records were in the 65-69 years age range for adults (15%), 1-5 years for children (62%), and 0-6 days for neonates (83%).



**Fig. A2** Study data sampling areas.

## A.2 Modelling Details

Each model (GPT-3.5, GPT-4, InSilicoVA, and InterVA-5) required pre-processing of the 6939 records into input data, and standardization of output COD codes from models for performance evaluation as not all models produced comparable codes across outputs. Although each model can assign multiple CODs per record, only the first

1335

1336 **Table A1** Study data by cause of death.

Age Group	CGHR-10 Cause of Death (COD)	Female	Male	Total
Adult, 18 CODs (n=3826, 55.1%)	Acute Respiratory Infections	48 (45.7%)	57 (54.3%)	105 (2.7%)
	Cancers	32 (65.3%)	17 (34.7%)	49 (1.3%)
	Chronic Respiratory Diseases	29 (38.7%)	46 (61.3%)	75 (2%)
	Diabetes Mellitus	14 (51.9%)	13 (48.1%)	27 (0.7%)
	Diarrhoeal Diseases	102 (49.8%)	103 (50.2%)	205 (5.4%)
	Ill-Defined	56 (47.9%)	61 (52.1%)	117 (3.1%)
	Ischemic Heart Disease	89 (53%)	79 (47%)	168 (4.4%)
	Liver And Alcohol Related Diseases	58 (45.3%)	70 (54.7%)	128 (3.3%)
	Malaria	372 (46.6%)	427 (53.4%)	799 (20.9%)
	Maternal Conditions	130 (100%)	N/A	130 (3.4%)
	Other Cardiovascular Diseases	59 (55.1%)	48 (44.9%)	107 (2.8%)
	Other Noncommunicable Diseases	160 (38.6%)	254 (61.4%)	414 (10.8%)
	Other Injuries	83 (23.2%)	274 (76.8%)	357 (9.3%)
	Road And Transport Injuries	73 (29.1%)	178 (70.9%)	251 (6.6%)
	Stroke	147 (44.4%)	184 (55.6%)	331 (8.7%)
	Suicide	N/A	3 (100%)	3 (0.1%)
	Tuberculosis	54 (31.6%)	117 (68.4%)	171 (4.5%)
	Unspecified Infections	175 (45%)	214 (55%)	389 (10.2%)
Child, 9 CODs (n=2636, 38%)	Congenital Anomalies	1 (100%)	N/A	1 (0%)
	Diarrhoeal Diseases	79 (45.1%)	96 (54.9%)	175 (6.6%)
	Epilepsy, Leukaemia, And	61 (53.5%)	53 (46.5%)	114 (4.3%)
	Other Noncommunicable Diseases			
	Ill-Defined	34 (48.6%)	36 (51.4%)	70 (2.7%)
	Injuries	51 (37.8%)	84 (62.2%)	135 (5.1%)
	Malaria	680 (49.2%)	702 (50.8%)	1382 (52.4%)
	Nutritional Deficiencies	7 (63.6%)	4 (36.4%)	11 (0.4%)
Child Male (n=1346, 51.1%)	Other Infections	338 (50.7%)	329 (49.3%)	667 (25.3%)
	Pneumonia	39 (48.1%)	42 (51.9%)	81 (3.1%)
Neonate, 7 CODs (n=477, 6.9%)	Birth Asphyxia And Birth Trauma	38 (36.9%)	65 (63.1%)	103 (21.6%)
	Congenital Anomalies	2 (100%)	N/A	2 (0.4%)
	Ill-Defined	11 (47.8%)	12 (52.2%)	23 (4.8%)
	Neonatal Infections	49 (49.5%)	50 (50.5%)	99 (20.8%)
	Other	2 (40%)	3 (60%)	5 (1%)
	Prematurity And Low Birthweight	39 (53.4%)	34 (46.6%)	73 (15.3%)
	Stillbirth	86 (50%)	86 (50%)	172 (36.1%)

1368

1369 generated COD response from GPT-3.5 and GPT-4, and the most probable COD

1370

1371 from InterVA-5 and InSilicoVA were used for evaluation. Section A.2.1 describes the

1372

1373 input data and parameters for each model, while Section A.2.3 details the outputs

1374

1375 from running each model.

1376

1377

1378

1379

1380

**Table A2** Study data by age range.

Age Group	Age Range	Female	Male	Total
Adult (n=3826, 55.1%) Adult Female (n=1681, 43.9%) Adult Male (n=2145, 56.1%)	12-14 Years	51 (37.8%)	84 (62.2%)	135 (3.5%)
	15-19 Years	115 (42.8%)	154 (57.2%)	269 (7%)
	20-24 Years	146 (53.1%)	129 (46.9%)	275 (7.2%)
	25-29 Years	159 (45.2%)	193 (54.8%)	352 (9.2%)
	30-34 Years	174 (50.9%)	168 (49.1%)	342 (8.9%)
	35-39 Years	153 (45.4%)	184 (54.6%)	337 (8.8%)
	40-44 Years	134 (42%)	185 (58%)	319 (8.3%)
	45-49 Years	148 (47%)	167 (53%)	315 (8.2%)
	50-54 Years	134 (39.6%)	204 (60.4%)	338 (8.8%)
	55-59 Years	96 (37.6%)	159 (62.4%)	255 (6.7%)
Child (n=2636, 38%) Child Female (n=1290, 48.9%) Child Male (n=1346, 51.1%)	60-64 Years	128 (40.8%)	186 (59.2%)	314 (8.2%)
	65-69 Years	243 (42.3%)	332 (57.7%)	575 (15%)
	1-5 Months	146 (47.4%)	162 (52.6%)	308 (11.7%)
	6-11 Months	160 (50.8%)	155 (49.2%)	315 (11.9%)
Neonate (n=477, 6.9%) Neonate Female (n=227, 47.6%) Neonate Male (n=250, 52.4%)	1-5 Years	822 (50.3%)	811 (49.7%)	1633 (61.9%)
	6-11 Years	162 (42.6%)	218 (57.4%)	380 (14.4%)
	0-6 Days	184 (46.6%)	211 (53.4%)	395 (82.8%)
	7-27 Days	43 (52.4%)	39 (47.6%)	82 (17.2%)

### A.2.1 Input Data and Preprocessing

For GPT-3.5 and GPT-4, 6939 text prompts were generated for each physician agreed record as input to instruct the models to assign CODs based on the open narratives.

Two types of text prompts were used: user prompts and system prompts. System prompts contained textual instructions to assign the role of a physician ICD-10 coder with expertise in Sierra Leone. The following system prompt was used for each record:

You are a physician with expertise in determining underlying causes of death in Sierra Leone by assigning the most probable ICD-10 code for each death using verbal autopsy narratives. Return only the ICD-10 code without description. E.g. A00. If there are multiple ICD-10 codes, show one code per line.

User prompts contained textual instructions to perform coding of VA records based on the age, sex, and narrative of the deceased. The following template was used to

1427 generate user prompts for each record, where `<age>` and `<sex>` from the questionnaire,  
1428 and `<narrative>` from the narratives, were replaced with values from the data:

```
1430 Determine the underlying cause of death and provide the most  
1431 probable ICD-10 code for a verbal autopsy narrative of a <age>  
1432 years old <sex> death in Sierra Leone: <narrative>
```

1435 For InterVA-5 and InSilicoVA, the standardized questionnaire data from the HEAL-SL  
1436 EVA were first converted into 2016 World Health Organization (WHO) VA question-  
1437 naire revision 1.5.1 Open Data Kit (ODK) format [75, 76] consisting of 526 variables  
1438 [77], followed by further conversion into OpenVA format [43] consisting of 353 vari-  
1439 ables [78] using the `pyCrossVA` version 0.97 Python package [79]. The 6939 records  
1440 were all converted into OpenVA formatted records for InterVA-5 and InSilicoVA.

1445

### 1447 A.2.2 Models and Parameters

1448

1449 The GPT-3.5 and GPT-4 Application Programming Interface (API) was accessed  
1450 using Python version 3.11.4 and used to assign CODs for each record. GPT-3.5 used  
1451 the `gpt-3.5-turbo` model, while GPT-4 used the `gpt-4-0613` model. The parameter  
1452 `temperature` for GPT-3.5 and GPT-4, representing the sampling temperature ranging  
1453 from 0 to 2 (default of 1), was set to 0 to produce more deterministic outputs [66].  
1454 Higher values closer to 2 may produce less deterministic outputs, while lower values  
1455 closer to 0 produce more deterministic outputs.

1456 The `openVA` R package was used to run InterVA-5 and InSilicoVA models to assign  
1457 CODs for each record in R version 4.3.1. The `openVA` package version 1.1.1 used  
1458 dependent packages `InterVA5` version 1.1.3 and `InSilicoVA` version 1.4.0. The `Nsim`  
1459 (number of iterations to run) parameter [80] for InSilicoVA was set to 9500, while  
1460 the `HIV` (level of prevalence of human immunodeficiency virus) and `Malaria` (level  
1461 of prevalence of Malaria) parameters [81] for InterVA-5 were both set to `'h'` (high)  
1462 reflecting HIV and Malaria disease assumptions in Sierra Leone [82, 83]. Note that the  
1472



default value of `Nsim=10000` for InSilicoVA ran until 9500 iterations before it stopped due to errors, thus `Nsim=9500` was used and ran successfully for all iterations.

### A.2.3 Output Data and Code Conversion

Of the 6939 input records, GPT-3.5, GPT-4, InterVA-5, and InSilicoVA were able to assign CODs for 6939 (100%), 6935 (>99%), 6830 (98%), 6830 (98%) records respectively. All 6830 (100%) InterVA-5 and InSilicoVA records with WHO VA 2016 v1.5 output codes [55] were converted into ICD-10 codes respectively. After all model outputs were converted to ICD-10 codes, they were further converted to CGHR-10 codes. The 6939 GPT-3.5 and 6935 GPT-4 output records with ICD-10 codes were converted into 6930 (>99%) and 6931 (>99) records with CGHR-10 codes, where <1% (9 and 8) records did not have matching CGHR-10 codes respectively. The 6830 InterVA-5 and InSilicoVA records with ICD-10 codes were converted into 6802 (>99%) and 6726 (98%) records with CGHR-10 codes respectively, where 28 (<1%) and 104 (1%) of records could not be converted into CGHR-10 codes.

## A.3 Performance Evaluation Details

The performance of GPT-3.5, GPT-4, InSilicoVA, and InterVA-5 models were evaluated with metrics at the population and individual level by comparing their CGHR-10 COD outputs for 6939 records to physician COD assignments. Section A.3.1 describes CSMF accuracy in detail for evaluating models on the population level, Section A.3.2 describes PCCC for evaluating models on the individual level. Records that were assigned a COD by physicians, but not by a model were considered to be an incorrect COD assignment by the model. CSMF accuracy and PCCC were calculated for each model overall and by three age groups (adult, child, and neonatal), then further into age and COD for each age group.

### 1519 **A.3.1 Cause Specific Mortality Fraction (CSMF) Accuracy**

1520

1521 CSMF accuracy measures the performance of models at the population level, compar-  
 1522 ing distributions of CODs between the physicians and the models [56]. To calculate  
 1523 CSMF accuracy, we first calculate  $CSMF_j$  as is the fraction of physician or model  
 1524 records for cause  $j$ , given by dividing the number of records for cause  $j$  with the total  
 1525 number of records as seen in Equation A1. Then, the  $CSMF_{MaximumError}$ , repre-  
 1526 senting the worst possible model, is calculated using Equation A2. Finally, the CSMF  
 1527 accuracy is given by Equation A3, where  $k$  is the number of causes,  $j$  is a cause,  
 1528  $CSMF_j^{true}$  is the true physician CSMF for cause  $j$ , and  $CSMF_j^{pred}$  is the prediction  
 1529 model CSMF for cause  $j$ . CSMF accuracy ranges from 0 to 1, where 1 means that the  
 1530 model completely matched the physician COD distribution and 0 means that it did  
 1531 not match the distribution at all.

1532

1533

$$1541 \quad CSMF_j = Records_j / Records \quad (A1)$$

1542

1543

$$1544 \quad CSMF_{MaximumError} = 2(1 - \min(CSMF_j^{true})) \quad (A2)$$

1545

1546

$$1547 \quad CSMF_{Accuracy} = 1 - \frac{\sum_{j=1}^k |CSMF_j^{true} - CSMF_j^{pred}|}{CSMF_{MaximumError}} \quad (A3)$$

1548

1549

### 1551 **A.3.2 Partial Chance Corrected Concordance (PCCC)**

1552

1553 PCCC measures the performance of models at the individual level, comparing COD  
 1554 assignments between the physicians and models on a record by record basis, correcting  
 1555 for COD assignments made purely by chance [56]. PCCC is given by Equation A5,  
 1556 where  $k$  is the number of top COD assignments from the model to consider,  $N$  is  
 1557 number of causes, and  $C$  is fraction of records where the physician COD assignment is  
 1558 one of the top COD assignments from the model. For this study, we set  $k$  to 1, making  
 1559  $C$  equivalent to the fraction of true positives  $TP$  or records where the physician COD

assignment is equal to the model COD assignment as shown in Equation A4. Higher PCCC values closer to 1 indicate that model COD assignments are similar to physician COD assignments, while values closer to 0 indicate that model COD assignments are not similar to physicians.

$$C = \frac{TP}{Records} \quad (A4)$$

$$PCCC(k) = \frac{C - \frac{k}{N}}{1 - \frac{k}{N}} \quad (A5)$$

## Appendix B Experiment on Repeated Runs of GPT-3.5

A short experiment was conducted to test the consistency of GPT-3.5 outputs repeated on the same record. 100 records, sampled randomly with approximately equal proportions across age groups, CODs, and survey rounds 1 and 2, were used to test repeated runs of GPT-3.5. Each record from the 100 records was rerun 10 times through GPT-3.5, resulting in ten COD outputs per record. The ICD-10 codes were then converted to CGHR-10 codes and tested for consistency, where completely inconsistent results had different ICD-10 or CGHR-10 codes for each of the 10 reruns (1 times+), and completely consistent results had the same ICD-10 or CGHR-10 code for all 10 reruns (10 times), on the same record.

The results are shown in Table B3. For all 100 records, GPT-3.5 assigns the same ICD-10 and CGHR-10 code for the same record 5 times or more out of 10. For 66 and 79 records, GPT-3.5 assigns the same ICD-10 and CGHR-10 code respectively for each record. This number increases to 94 (from 66) and 96 (from 79) when reducing the number of times out of 10 that GPT-3.5 assigns the same ICD-10 and CGHR-10 code respectively. Thus, GPT-3.5 does not always produce the same outputs when repeated on the same record (10 times out of 10), even when the temperature is set

to 0, but does so for more than half the records. For most records (more than 90%), GPT-3.5 will produce the same outputs for the same record 7 times or more out of 10.

**Table B3** Records with same GPT-3.5 outputs based on 10 repeated reruns of 100 records

Times with Same GPT-3.5 Outputs	ICD-10 Records	CGHR-10 Records
1 times+ (inconsistent)	100	100
2 times+	100	100
3 times+	100	100
4 times+	100	100
5 times+	100	100
6 times+	94	96
7 times+	92	94
8 times+	86	91
9 times+	79	86
10 times (consistent)	66	79

## Appendix C Exploration of Neonatal Infections

An exploration of neonatal infections (n=99, 21% of 477 records) was done to understand the low performance of GPT models (0.23 PCCC) for neonatal infections, and high performance of InSilicoVA (0.87 PCCC). In Table C4, about half the records were assigned correctly, and a majority (n=33, 33%) of the other records were misclassified as other, while prematurity and low birthweight, birth asphyxia & birth trauma, and ill-defined make up the rest. On closer inspection of the 49 records with misclassified assignments, the ICD-10 code R50 was assigned in 20 records. R50 falls under unspecified infections in the adult CGHR-10 category, but in the other category for neonates. B50 was assigned in 4 records, falling under malaria, but a similar B54 falls under neonatal infections. P81 was assigned in 3 records, referring to fever of unknown origin, which falls under other, and P07 was assigned in 7 records, falling under prematurity and low birthweight.

In most misclassified records, there is mention of infections, but the misclassifications occur due to the finer details of the ICD-10 code classifications, the categorization

decisions of the CGHR-10 codes, and missing information from the questionnaire. For R50 misclassifications, GPT may have confused descriptions across adult and neonatal age groups. Using the same definition of R50, but in the context of neonates, may result in codes closer to neonatal infections (e.g. B54). For B50 misclassifications, the similar B54 was categorized in CGHR-10 as neonatal infections, but B50 was categorized as other. P81 refers to fever of unknown origin, which may be difficult to differentiate between infection and other causes without information from the questionnaire. P07 refers to prematurity and low birthweight, where GPT initially assigned P07 as the age of the neonate was mentioned first, but later mentions infections as an alternative following the order of information in the narratives. Thus, it may be possible to improve the performance GPT models using better prompts based on the context of VA manuals and CGHR-10 codes, and by also including questionnaire information in the prompts.

**Table C4** GPT-4 CGHR-10 COD assignment for physician coded neonatal infections records.

GPT-4 Assigned Cause of Death (CGHR-10)	Records
Neonatal infections	50 (51%)
Other	33 (33%)
Prematurity and low birthweight	9 (9%)
Birth asphyxia & birth trauma	5 (6%)
Ill-defined	2 (2%)
Total	99 (100%)

## 1703 References

- 1704
- 1705 [1] World Health Organization.: Non Communicable Diseases: Key Facts.
- 1706 <https://www.who.int/news-room/fact-sheets/detail/noncommunicable-diseases>.
- 1707
- 1708
- 1709 [2] Benziger CP, Roth GA, Moran AE. The Global Burden of Disease Study and
- 1710 the Preventable Burden of NCD. *Global Heart*. 2016 Dec;11(4):393–397. <https://doi.org/10.1016/j.gheart.2016.10.024>.
- 1711
- 1712
- 1713
- 1714
- 1715 [3] Lawn JE, Kerber K, Enweronu-Laryea C, Cousens S. 3.6 Million Neonatal
- 1716 Deaths—What Is Progressing and What Is Not? *Seminars in Perinatology*. 2010
- 1717 Dec;34(6):371–386. <https://doi.org/10.1053/j.semperi.2010.09.011>.
- 1718
- 1719
- 1720
- 1721 [4] Lassi ZS, Bhutta ZA. Community-based Intervention Packages for Reducing
- 1722 Maternal and Neonatal Morbidity and Mortality and Improving Neonatal Out-
- 1723 comes. *Cochrane Database of Systematic Reviews*. 2015;(3). [https://doi.org/10.](https://doi.org/10.1002/14651858.CD007754.pub3)
- 1724
- 1725 [1002/14651858.CD007754.pub3](https://doi.org/10.1002/14651858.CD007754.pub3).
- 1726
- 1727
- 1728
- 1729 [5] Liu NH, Daumit GL, Dua T, Aquila R, Charlson F, Cuijpers P, et al. Excess
- 1730 Mortality in Persons with Severe Mental Disorders: A Multilevel Intervention
- 1731 Framework and Priorities for Clinical Practice, Policy and Research Agendas.
- 1732 *World Psychiatry*. 2017;16(1):30–40. <https://doi.org/10.1002/wps.20384>.
- 1733
- 1734
- 1735
- 1736 [6] Ewig S, Torres A. Community-Acquired Pneumonia as an Emergency: Time for
- 1737 an Aggressive Intervention to Lower Mortality. *European Respiratory Journal*.
- 1738 2011 Aug;38(2):253–260. <https://doi.org/10.1183/09031936.00199810>.
- 1739
- 1740
- 1741
- 1742 [7] World Health Organization. SCORE for Health Data Technical Package: Global
- 1743 Report on Health Data Systems and Capacity, 2020; 2021.
- 1744
- 1745
- 1746
- 1747
- 1748

- [8] de Savigny D, Riley I, Chandramohan D, Odhiambo F, Nichols E, Notzon S, et al. Integrating Community-Based Verbal Autopsy into Civil Registration and Vital Statistics (CRVS): System-Level Considerations. *Global Health Action*. 2017 Jan;10(1):1272882. <https://doi.org/10.1080/16549716.2017.1272882>.
- [9] Thomas LM, D’Ambruoso L, Balabanova D. Verbal Autopsy in Health Policy and Systems: A Literature Review. *BMJ Global Health*. 2018 May;3(2):e000639. <https://doi.org/10.1136/bmjgh-2017-000639>.
- [10] Rampatige R, Mikkelsen L, Hernandez B, Riley I, Lopez AD. Systematic Review of Statistics on Causes of Deaths in Hospitals: Strengthening the Evidence for Policy-Makers. *Bulletin of the World Health Organization*. 2014 Sep;92:807–816. <https://doi.org/10.2471/BLT.14.137935>.
- [11] Adair T. Who Dies Where? Estimating the Percentage of Deaths That Occur at Home. *BMJ Global Health*. 2021 Sep;6(9):e006766. <https://doi.org/10.1136/bmjgh-2021-006766>.
- [12] World Health Organization. Verbal Autopsy Standards: 2022 WHO Verbal Autopsy Instrument; 2023.
- [13] Chandramohan D, Fottrell E, Leitao J, Nichols E, Clark SJ, Alsokhn C, et al. Estimating Causes of Death Where There Is No Medical Certification: Evolution and State of the Art of Verbal Autopsy. *Global Health Action*. 2021 Oct;14(sup1):1982486. <https://doi.org/10.1080/16549716.2021.1982486>.
- [14] World Health Organization. Verbal Autopsy Standards: Ascertaining and Attributing Cause of Death. World Health Organization; 2007.
- [15] Gomes M, Begum R, Sati P, Dikshit R, Gupta PC, Kumar R, et al. Nationwide Mortality Studies To Quantify Causes Of Death: Relevant Lessons From India’s

1795 Million Death Study. Health Affairs. 2017 Nov;36(11):1887–1895. [https://doi.](https://doi.org/10.1377/hlthaff.2017.0635)  
1796 [org/10.1377/hlthaff.2017.0635](https://doi.org/10.1377/hlthaff.2017.0635).  
1797  
1798  
1799 [16] Jha P, Gajalakshmi V, Gupta PC, Kumar R, Mony P, Dhingra N, et al. Prospec-  
1800 tive Study of One Million Deaths in India: Rationale, Design, and Validation  
1801 Results. PLOS Medicine. 2005 Dec;3(2):e18. [https://doi.org/10.1371/journal.](https://doi.org/10.1371/journal.pmed.0030018)  
1802 [pmed.0030018](https://doi.org/10.1371/journal.pmed.0030018).  
1803  
1804  
1805  
1806 [17] McCormick TH, Li ZR, Calvert C, Crampin AC, Kahn K, Clark SJ. Prob-  
1807 abilistic Cause-of-death Assignment Using Verbal Autopsies. Journal of the  
1808 American Statistical Association. 2016;111(515):1036–1049. [https://doi.org/10.](https://doi.org/10.1080/01621459.2016.1152191)  
1809 [1080/01621459.2016.1152191](https://doi.org/10.1080/01621459.2016.1152191).  
1810  
1811  
1812  
1813 [18] Morris SK, Bassani DG, Kumar R, Awasthi S, Paul VK, Jha P. Factors Associated  
1814 with Physician Agreement on Verbal Autopsy of over 27000 Childhood Deaths in  
1815 India. PloS one. 2010;5(3):e9583.  
1816  
1817  
1818  
1819 [19] Soleman N, Chandramohan D, Shibuya K. Verbal Autopsy: Current Practices  
1820 and Challenges. Bulletin of the World Health Organization. 2006;84(3):239–245.  
1821  
1822  
1823 [20] Byass P, Hussain-Alkhateeb L, D’Ambruso L, Clark S, Davies J, Fottrell E,  
1824 et al. An Integrated Approach to Processing WHO-2016 Verbal Autopsy Data:  
1825 The InterVA-5 Model. BMC Medicine. 2019 May;17(1):102. [https://doi.org/10.](https://doi.org/10.1186/s12916-019-1333-6)  
1826 [1186/s12916-019-1333-6](https://doi.org/10.1186/s12916-019-1333-6).  
1827  
1828  
1829  
1830 [21] Jha P, Kumar D, Dikshit R, Budukh A, Begum R, Sati P, et al. Automated versus  
1831 Physician Assignment of Cause of Death for Verbal Autopsies: Randomized Trial  
1832 of 9374 Deaths in 117 Villages in India. BMC Medicine. 2019 Jun;17(1):116.  
1833 <https://doi.org/10.1186/s12916-019-1353-2>.  
1834  
1835  
1836  
1837  
1838  
1839  
1840



- [22] Leitao J, Desai N, Aleksandrowicz L, Byass P, Miasnikof P, Tollman S, et al. Comparison of Physician-Certified Verbal Autopsy with Computer-Coded Verbal Autopsy for Cause of Death Assignment in Hospitalized Patients in Low- and Middle-Income Countries: Systematic Review. *BMC Medicine*. 2014 Feb;12(1):22. <https://doi.org/10.1186/1741-7015-12-22>.
- [23] Desai N, Aleksandrowicz L, Miasnikof P, Lu Y, Leitao J, Byass P, et al. Performance of Four Computer-Coded Verbal Autopsy Methods for Cause of Death Assignment Compared with Physician Coding on 24,000 Deaths in Low- and Middle-Income Countries. *BMC Medicine*. 2014 Feb;12(1):20. <https://doi.org/10.1186/1741-7015-12-20>.
- [24] Tunga M, Lungo J, Chambua J, Kateule R. Verbal Autopsy Models in Determining Causes of Death. *Tropical Medicine & International Health*. 2021;26(12):1560–1567. <https://doi.org/10.1111/tmi.13678>.
- [25] Oti SO, Kyobutungi C. Verbal Autopsy Interpretation: A Comparative Analysis of the InterVA Model versus Physician Review in Determining Causes of Death in the Nairobi DSS. *Population Health Metrics*. 2010 Jun;8(1):21. <https://doi.org/10.1186/1478-7954-8-21>.
- [26] Jeblee S, Gomes M, Jha P, Rudzicz F, Hirst G. Automatically Determining Cause of Death from Verbal Autopsy Narratives. *BMC Medical Informatics and Decision Making*. 2019 Jul;19(1):127. <https://doi.org/10.1186/s12911-019-0841-9>.
- [27] Blanco A, Pérez A, Casillas A, Cobos D. Extracting Cause of Death From Verbal Autopsy With Deep Learning Interpretable Methods. *IEEE Journal of Biomedical and Health Informatics*. 2021 Apr;25(4):1315–1325. <https://doi.org/10.1109/JBHI.2020.3005769>.

1887 [28] King C, Zamawe C, Banda M, Bar-Zeev N, Beard J, Bird J, et al. The Quality and  
1888 Diagnostic Value of Open Narratives in Verbal Autopsy: A Mixed-Methods Anal-  
1889 ysis of Partnered Interviews from Malawi. BMC Medical Research Methodology.  
1890 2016 Feb;16(1):13. <https://doi.org/10.1186/s12874-016-0115-5>.  
1891  
1892 [29] Chang Y, Wang X, Wang J, Wu Y, Yang L, Zhu K, et al.: A Survey on Evaluation  
1893 of Large Language Models. arXiv.  
1894  
1895 [30] Lund BD, Wang T. Chatting about ChatGPT: How May AI and GPT Impact  
1896 Academia and Libraries? Library Hi Tech News. 2023 Jan;40(3):26–29. <https://doi.org/10.1108/LHTN-01-2023-0009>.  
1897  
1898 [31] Svyatkovskiy A, Deng SK, Fu S, Sundaresan N. IntelliCode Compose: Code  
1899 Generation Using Transformer. In: Proceedings of the 28th ACM Joint Meeting on  
1900 European Software Engineering Conference and Symposium on the Foundations  
1901 of Software Engineering. ESEC/FSE 2020. New York, NY, USA: Association for  
1902 Computing Machinery; 2020. p. 1433–1443.  
1903  
1904 [32] Haupt CE, Marks M. AI-Generated Medical Advice—GPT and Beyond. JAMA.  
1905 2023 Apr;329(16):1349–1350. <https://doi.org/10.1001/jama.2023.5321>.  
1906  
1907 [33] Wu T, He S, Liu J, Sun S, Liu K, Han QL, et al. A Brief Overview of ChatGPT:  
1908 The History, Status Quo and Potential Future Development. IEEE/CAA Journal  
1909 of Automatica Sinica. 2023;10(5):1122–1136. [https://doi.org/10.1109/JAS.2023.](https://doi.org/10.1109/JAS.2023.123618)  
1910 [123618](https://doi.org/10.1109/JAS.2023.123618).  
1911  
1912 [34] OpenAI, Achiam J, Adler S, Agarwal S, Ahmad L, Akkaya I, et al.: GPT-4  
1913 Technical Report. arXiv.  
1914  
1915 [35] Njala University.: Healthy Sierra Leone. <https://healsl.org/>.  
1916  
1917  
1918  
1919  
1920  
1921  
1922  
1923  
1924  
1925  
1926  
1927  
1928  
1929  
1930  
1931  
1932

- [36] Carshon-Marsh R, Aimone A, Ansumana R, Swaray IB, Assalif A, Musa A, et al. Child, Maternal, and Adult Mortality in Sierra Leone: Nationally Representative Mortality Survey 2018–20. *The Lancet Global Health*. 2022 Jan;10(1):e114–e123. [https://doi.org/10.1016/S2214-109X\(21\)00459-9](https://doi.org/10.1016/S2214-109X(21)00459-9).
- [37] World Health Organization. ICD-10: International Statistical Classification of Diseases and Related Health Problems (10th Revision); 2011.
- [38] Aleksandrowicz L, Malhotra V, Dikshit R, Gupta PC, Kumar R, Sheth J, et al. Performance Criteria for Verbal Autopsy-Based Systems to Estimate National Causes of Death: Development and Application to the Indian Million Death Study. *BMC Medicine*. 2014 Feb;12(1):21. <https://doi.org/10.1186/1741-7015-12-21>.
- [39] Barnett ML, Boddupalli D, Nundy S, Bates DW. Comparative Accuracy of Diagnosis by Collective Intelligence of Multiple Physicians vs Individual Physicians. *JAMA Network Open*. 2019 Mar;2(3):e190096. <https://doi.org/10.1001/jamanetworkopen.2019.0096>.
- [40] Hsiao M, Morris SK, Bassani DG, Montgomery AL, Thakur JS, Jha P. Factors Associated with Physician Agreement on Verbal Autopsy of over 11500 Injury Deaths in India. *PLOS ONE*. 2012 Jan;7(1):e30336. <https://doi.org/10.1371/journal.pone.0030336>.
- [41] Murray CJ, Lozano R, Flaxman AD, Serina P, Phillips D, Stewart A, et al. Using Verbal Autopsy to Measure Causes of Death: The Comparative Performance of Existing Methods. *BMC Medicine*. 2014 Jan;12(1):5. <https://doi.org/10.1186/1741-7015-12-5>.

1979 [42] Benara SK, Sharma S, Juneja A, Nair S, Gulati BK, Singh KJ, et al. Evaluation of  
1980  
1981 Methods for Assigning Causes of Death from Verbal Autopsies in India. *Frontiers*  
1982 in Big Data. 2023 Aug;6:1197471. <https://doi.org/10.3389/fdata.2023.1197471>.  
1983  
1984  
1985 [43] Li ZR, Thomas J, Choi E, McCormick TH, Clark SJ. The openVA Toolkit for  
1986 Verbal Autopsies. *The R Journal*. 2023 Feb;p. 1.  
1987  
1988  
1989 [44] BAYES. An Essay towards Solving a Problem in the Doctrine of Chances.  
1990 *Biometrika*. 1958;45(3-4):296–315.  
1991  
1992  
1993 [45] Byass P, Chandramohan D, Clark SJ, D’Ambruso L, Fottrell E, Graham WJ,  
1994 et al. Strengthening Standardised Interpretation of Verbal Autopsy Data: The  
1995 New InterVA-4 Tool. *Global Health Action*. 2012 Dec;5(1):19281. [https://doi.](https://doi.org/10.3402/gha.v5i0.19281)  
1996  
1997 [org/10.3402/gha.v5i0.19281](https://doi.org/10.3402/gha.v5i0.19281).  
1998  
1999  
2000  
2001 [46] Brooks S. Markov Chain Monte Carlo Method and Its Application. *Journal of*  
2002 the Royal Statistical Society: Series D (The Statistician). 1998 Mar;47(1):69–100.  
2003  
2004 <https://doi.org/10.1111/1467-9884.00117>.  
2005  
2006  
2007 [47] Chib S. Markov Chain Monte Carlo Methods: Computation and Inference.  
2008 *Handbook of econometrics*. 2001;5:3569–3649.  
2009  
2010  
2011 [48] Han C, Carlin BP. Markov Chain Monte Carlo Methods for Computing Bayes  
2012 Factors: A Comparative Review. *Journal of the American Statistical Association*.  
2013 2001 Sep;96(455):1122–1132. <https://doi.org/10.1198/016214501753208780>.  
2014  
2015  
2016 [49] Brown TB, Mann B, Ryder N, Subbiah M, Kaplan J, Dhariwal P, et al.: Language  
2017 Models Are Few-Shot Learners. *arXiv*.  
2018  
2019  
2020 [50] Vaswani A, Shazeer N, Parmar N, Uszkoreit J, Jones L, Gomez AN, et al. Atten-  
2021 tion Is All You Need. In: *Advances in Neural Information Processing Systems*.  
2022  
2023  
2024

vol. 30. Curran Associates, Inc.; 2017. . 2025

[51] Ouyang L, Wu J, Jiang X, Almeida D, Wainwright CL, Mishkin P, et al.: Training 2026  
Language Models to Follow Instructions with Human Feedback. arXiv. 2027  
2028

[52] Christiano PF, Leike J, Brown T, Martic M, Legg S, Amodei D. Deep Rein- 2031  
forcement Learning from Human Preferences. Advances in neural information 2032  
processing systems. 2017;30. 2033  
2034  
2035  
2036

[53] Stiennon N, Ouyang L, Wu J, Ziegler D, Lowe R, Voss C, et al. Learning to 2037  
Summarize with Human Feedback. Advances in Neural Information Processing 2038  
Systems. 2020;33:3008–3021. 2039  
2040  
2041  
2042

[54] Wirth C, Akrou R, Neumann G, Fürnkranz J. A Survey of Preference-Based 2043  
Reinforcement Learning Methods. The Journal of Machine Learning Research. 2044  
2045  
2046  
2047  
2048

[55] World Health Organization.: Verbal Autopsy Standards: The 2016 WHO Ver- 2049  
bal Autopsy Instrument. [https://www.who.int/publications/m/item/verbal-](https://www.who.int/publications/m/item/verbal-autopsy-standards-the-2016-who-verbal-autopsy-instrument) 2050  
[autopsy-standards-the-2016-who-verbal-autopsy-instrument.](https://www.who.int/publications/m/item/verbal-autopsy-standards-the-2016-who-verbal-autopsy-instrument) 2051  
2052  
2053  
2054

[56] Murray CJ, Lozano R, Flaxman AD, Vahdatpour A, Lopez AD. Robust Metrics 2055  
for Assessing the Performance of Different Verbal Autopsy Cause Assignment 2056  
Methods in Validation Studies. Population Health Metrics. 2011 Aug;9(1):28. 2057  
2058  
2059  
<https://doi.org/10.1186/1478-7954-9-28>. 2060  
2061

[57] Setel PW, Whiting DR, Hemed Y, Chandramohan D, Wolfson LJ, Alberti 2062  
KGMM, et al. Validity of Verbal Autopsy Procedures for Determining Cause of 2063  
Death in Tanzania. Tropical Medicine & International Health. 2006;11(5):681– 2064  
696. <https://doi.org/10.1111/j.1365-3156.2006.01603.x>. 2065  
2066  
2067  
2068  
2069  
2070

2071 [58] World Health Organization. World Health Statistics 2024: Monitoring Health for  
2072 the SDGs, Sustainable Development Goals. World Health Organization; 2024.  
2073  
2074

2075 [59] World Health Organization.: Cause-Specific Mortality 2000 and 2021. The Global  
2076 Health Observatory.  
2077  
2078

2079 [60] Rasmussen LA, Cascio MA, Ferrand A, Shevell M, Racine E. The Complex-  
2080 ity of Physicians' Understanding and Management of Prognostic Uncertainty  
2081 in Neonatal Hypoxic-Ischemic Encephalopathy. *Journal of Perinatology*. 2019  
2082 Feb;39(2):278–285. <https://doi.org/10.1038/s41372-018-0296-3>.  
2083  
2084  
2085  
2086

2087 [61] Faison G, Chou FS, Feudtner C, Janvier A. When the Unknown Is Unknowable:  
2088 Confronting Diagnostic Uncertainty. *Pediatrics*. 2023 Sep;152(4):e2023061193.  
2089 <https://doi.org/10.1542/peds.2023-061193>.  
2090  
2091  
2092

2093 [62] OpenAI.: Pricing. <https://openai.com/api/pricing/>.  
2094

2095 [63] Tao G, Cheng S, Zhang Z, Zhu J, Shen G, Zhang X.: Opening A Pandora's Box:  
2096 Things You Should Know in the Era of Custom GPTs. *arXiv*.  
2097  
2098

2099 [64] Khowaja SA, Khuwaja P, Dev K, Wang W, Nkenyereye L. ChatGPT  
2100 Needs SPADE (Sustainability, PrivAcy, Digital Divide, and Ethics) Evalua-  
2101 tion: A Review. *Cognitive Computation*. 2024 May;[https://doi.org/10.1007/](https://doi.org/10.1007/s12559-024-10285-1)  
2102 [s12559-024-10285-1](https://doi.org/10.1007/s12559-024-10285-1).  
2103  
2104  
2105  
2106

2107 [65] Wu X, Duan R, Ni J. Unveiling Security, Privacy, and Ethical Concerns of  
2108 ChatGPT. *Journal of Information and Intelligence*. 2024;2(2):102–115.  
2109  
2110

2111 [66] OpenAI.: OpenAI Platform: API Reference (Tempera-  
2112 ture Parameter). [https://platform.openai.com/docs/api-](https://platform.openai.com/docs/api-reference/completions/create#completions-create-temperature)  
2113 [reference/completions/create#completions-create-temperature](https://platform.openai.com/docs/api-reference/completions/create#completions-create-temperature).  
2114  
2115  
2116

- [67] Johnson D, Goodman R, Patrinely J, Stone C, Zimmerman E, Donald R, et al. Assessing the Accuracy and Reliability of AI-Generated Medical Responses: An Evaluation of the Chat-GPT Model. Research Square. 2023 Feb;p. rs.rs-2566942. <https://doi.org/10.21203/rs.3.rs-2566942/v1>.
- [68] Jang ME, Lukasiewicz T.: Consistency Analysis of ChatGPT. arXiv.
- [69] Krishna S, Bhambra N, Bleakney R, Bhayana R, Atzen S. Evaluation of Reliability, Repeatability, Robustness, and Confidence of GPT-3.5 and GPT-4 on a Radiology Board-Style Examination. Radiology. 2024 May;311(2):e232715. <https://doi.org/10.1148/radiol.232715>.
- [70] Wang J, Shi E, Yu S, Wu Z, Ma C, Dai H, et al.: Prompt Engineering for Healthcare: Methodologies and Applications. arXiv.
- [71] Lewis P, Perez E, Piktus A, Petroni F, Karpukhin V, Goyal N, et al. Retrieval-Augmented Generation for Knowledge-Intensive NLP Tasks. In: Proceedings of the 34th International Conference on Neural Information Processing Systems. NIPS '20. Red Hook, NY, USA: Curran Associates Inc.; 2020. p. 9459–9474.
- [72] Meskó B. Prompt Engineering as an Important Emerging Skill for Medical Professionals: Tutorial. Journal of medical Internet research. 2023;25:e50638.
- [73] Almasre M. Development and Evaluation of a Custom GPT for the Assessment of Students' Designs in a Typography Course. Education Sciences. 2024 Feb;14(2):148. <https://doi.org/10.3390/educsci14020148>.
- [74] Loh P, Fottrell E, Beard J, Bar-Zeev N, Phiri T, Banda M, et al. Added Value of an Open Narrative in Verbal Autopsies: A Mixed-Methods Evaluation from Malawi. BMJ Paediatrics Open. 2021 Feb;5(1):e000961. <https://doi.org/10.1136/bmjpo-2020-000961>.

2163 [75] World Health Organization.: ODK for Verbal Autopsy: A Quick Guide.  
2164 <https://www.who.int/publications/m/item/odk-for-verbal-autopsy-a-quick->  
2165 [guide.](https://www.who.int/publications/m/item/odk-for-verbal-autopsy-a-quick-)  
2166  
2167  
2168  
2169 [76] Nafundi.: ODK - Collect Data Anywhere.  
2170  
2171 [77] DiPasquale A, Maire N, Bratschi M.: Release ODK 2016 WHO VA Instrument  
2172 1.5.1 SwissTPH/WHO-VA. Swiss Tropical and Public Health Institute.  
2173  
2174  
2175 [78] Byass P.: InterVA-5.1 User Guide.  
2176  
2177  
2178 [79] Thomas J, ekarpinskiMITRE, pkmitre, owentrigueros, Choi P, Chu Y.: Pycrossva:  
2179 Prepare Data from WHO and PHRMC Instruments for Verbal Autopsy Algo-  
2180 rithms.  
2181  
2182  
2183  
2184 [80] Li ZR, McCormick T, Clark S.: InSilicoVA: Probabilistic Verbal Autopsy Coding  
2185 with 'InSilicoVA' Algorithm.  
2186  
2187  
2188 [81] Thomas J, Li Z, Byass P, McCormick T, Boyas M, Clark S.: InterVA5: Replicate  
2189 and Analyse 'InterVA5'.  
2190  
2191  
2192 [82] Yendewa GA, Poveda E, Yendewa SA, Sahr F, Quiñones-Mateu ME, Salata RA.  
2193 HIV/AIDS in Sierra Leone: Characterizing the Hidden Epidemic. AIDS reviews.  
2194 2018;20(2).  
2195  
2196  
2197  
2198 [83] Walker PG, White MT, Griffin JT, Reynolds A, Ferguson NM, Ghani AC. Malaria  
2199 Morbidity and Mortality in Ebola-affected Countries Caused by Decreased  
2200 Health-Care Capacity, and the Potential Effect of Mitigation Strategies: A  
2201 Modelling Analysis. The Lancet Infectious Diseases. 2015;15(7):825–832.  
2202  
2203  
2204  
2205  
2206  
2207  
2208