

Computer Assisted Verbal Autopsy: Comparing
Large Language Models to Physicians for
Assigning Causes to 6939 Deaths in Sierra Leone
from 2019-2022

Richard Wen^{1*}, Anteneh Tesfaye Assalif^{1,2}, Andy Sze-Heng Lee¹,
Rajeev Kamadod¹, Asha Behdinan¹, Ronald Carshon-Marsh¹,
Catherine Meh¹, Thomas Kai Sze Ng¹, Patrick Brown¹,
Prabhat Jha¹, Rashid Ansumana²

^{1*}Centre for Global Health Research, St. Michael's Hospital, Unity
Health Toronto and University of Toronto, 30 Bond St, Toronto, M5B
1W8, Ontario, Canada.

²School of Community Health Sciences, Njala University, Bo, Sierra
Leone.

*Corresponding author(s). E-mail(s): richard.wen@utoronto.ca;
Contributing authors: antenehta@gmail.com; andylee@cs.toronto.edu;
rajeevk@kentropy.com; asha.behdinan@mail.utoronto.ca;
ronald.carshonmarsh@mail.utoronto.ca; catherine.meh@unityhealth.to;
kaisze.ng@unityhealth.to; patrick.brown@utoronto.ca;
prabhat.jha@utoronto.ca; rashidansumana@gmail.com;

Abstract

Background: Verbal autopsies (VAs) collect information on deaths occurring
outside traditional healthcare settings to estimate representative Causes of Death
(CODs). Current computer models assign CODs at population-level accuracy
comparable to physicians, but perform poorly at the individual level, largely
due to reliance on structured questionnaire data and neglect of narrative free

text. Recently, the large language model ChatGPT-4 demonstrated human-level performance on professional and academic benchmarks. While ChatGPT-4 shows promise in COD assignment, its application to VA narratives has not yet been evaluated.

Methods: We analyzed 6,939 VA records from Sierra Leone (2019–2022) to compare four models, GPT-3.5, GPT-4, InterVA-5, and InSilicoVA, against physician-assigned CODs at population and individual levels. GPT models used narratives, whereas InterVA-5 and InSilicoVA relied on questionnaires. CODs were grouped into 19, 10, and 7 categories for adult, child, and neonatal deaths. Cause Specific Mortality Fraction (CSMF) accuracy and Partial Chance Corrected Concordance (PCCC) were used to assess population and individual level agreement with physician coding respectively, stratified by age and COD.

Results: GPT-4 outperformed all models overall (PCCC=0.61), followed by GPT-3.5 (0.56) and InSilicoVA/InterVA-5 (0.44). GPT-4 achieved the highest PCCC for adult and neonatal deaths (0.64 and 0.58), with GPT-3.5 for child deaths (0.54). Across ages, model performance increased from 1 month to 14 years (~ 0.10 – 0.75 PCCC) and declined from 15 to 69 years (~ 0.70 – 0.35). GPT-4, GPT-3.5, and InSilicoVA achieved the highest PCCC in 17, 9, and 4 of the 30 CODs, respectively. At the population level, all models achieved comparable CSMF accuracies (0.74–0.79).

Conclusion: All models performed similarly at the population level, but GPT models and InSilicoVA showed greater performance for specific CODs at the individual level. GPT models demonstrated improvements over InterVA-5 and InSilicoVA models. This study provides foundational evidence for integrating computer models to assist physicians with alternative diagnoses, helping reduce ill-defined codes and improve agreement in COD assignment.

Keywords: Cause of Death, Physicians, Computer-Assisted Diagnosis, Artificial Intelligence, Natural Language Processing, Machine Learning, Mortality, Surveillance, Mathematical Models, Global Health

1 Background

Every year, 41 million people died prematurely from noncommunicable diseases, accounting for 74% of all deaths globally [1]. While most of these deaths are preventable, effective intervention requires evidence-based resource allocation that targets high-risk populations [2]. Reliable mortality counts and accurate Cause of Death (COD) data are essential for guiding public health policy and reducing premature mortality [3–6]. However, civil registration and vital statistics systems remain incomplete in many low-income countries. Fewer than half of all deaths are registered, and among

these, only 8% have an assigned COD [7]. To address this gap, Verbal Autopsy (VA) has been deployed as a scalable method for collecting mortality data and assigning likely CODs, particularly for deaths that occur outside of healthcare facilities, which account for more than half of all deaths [8–11].

VA involves two major components: survey and COD assignment [12–14]. In the survey component, trained interviewers use structured questionnaires and open narrative prompts to gather data from relatives or close contacts of the deceased. In the COD assignment component, physicians review these data to determine the most likely COD. However, reliance on physician assignment has been criticized for limited reproducibility and subjectivity [15–19]. To overcome these limitations, automated Computer Coded Verbal Autopsy (CCVA) methods such as InterVA [20] and InSilicoVA [17] have been developed. These models offer scalable and reproducible alternatives and have demonstrated comparable performance to physicians at the population level. However, their performance at the individual level remains limited [21–25], while their reliance on structured questionnaire data often omits open narrative text, which can contain additional contextual and chronological information that may improve diagnostic accuracy [26–28].

Recent advances in large language models (LLMs), trained on vast textual datasets using deep learning methods, have significantly improved natural language processing (NLP) capabilities. These include tasks such as question answering, code generation, and medical reasoning based on free text [29–32]. ChatGPT, developed by OpenAI and released in 2022, is a widely accessible LLM capable of generating human-like responses to natural language queries. Earlier versions (GPT-1 to GPT-3) scaled from 117 million to 175 billion parameters and were trained on data ranging from 5 GB to 45 TB [33]. In 2023, ChatGPT-4 was introduced, achieving human-level performance on a range of academic and professional benchmarks [34]. Given the underutilization of narrative free text in VA analysis and the capabilities of LLMs in processing

such data, we conducted a study using VA records from Sierra Leone (2019–2022) to compare four models, GPT-3.5, GPT-4, InterVA-5, and InSilicoVA, against physician-assigned CODs. This work aims to evaluate the potential of LLMs in enhancing COD assignment from narrative data in low-resource settings.

2 Methods

This study outlines the methodology used to compare cause of death (COD) assignments from four models, GPT-3.5, GPT-4, InterVA-5, and InSilicoVA, with physician-determined CODs, as summarized in Figure 1. The dataset was first filtered to include only records with physician agreement, as described in Section 2.1. Section 2.2 details the input formats and output structures of the four models. Section 2.3 presents the evaluation framework, which compares model outputs to physician-assigned CODs using both population-level and individual-level performance metrics. Additional methodological details are provided in Appendix A.

2.1 Verbal Autopsy (VA) Data

A total of 11,920 verbal autopsy (VA) records were obtained from the HEAL-SL study [35, 36], which employed dual-coded Electronic Verbal Autopsy (EVA). Each record was independently reviewed by two randomly selected physicians, who assigned COD codes based on the International Classification of Diseases, 10th Revision (ICD-10) [37]. Agreement between physician-assigned CODs was evaluated using Central Medical Evaluation Agreement 10 (CMEA-10) codes, which group related ICD-10 codes into broader, clinically similar categories [38] (see Additional File 2). If both codes fell within the same CMEA-10 group, the record was considered in agreement. Disagreements entered a reconciliation phase, where each physician was shown both the assigned codes and the reasoning from the other physician. Physicians could then (1) retain their original code, (2) adopt the other physician’s code, or (3) assign a new




Fig. 1 Study methods.

code. Records that remained unresolved proceeded to adjudication, where a senior physician reviewed all reasoning and assignments and issued a final COD.

To ensure comparability with physician coding, only records with physician agreement were used in this study, as such cases provide higher confidence in the COD assignment [18, 39, 40]. From the original dataset, 6,942 records met this criterion. All ICD-10 codes were then standardized to CGHR-10 categories (see Additional File 1), which group causes into 19, 10, and 7 categories for adults (12–69 years), children (28 days to 11 years), and neonates (under 28 days), respectively. After excluding three records without a valid CGHR-10 category, a total of 6,939 physician-agreed records (3,826 adult, 2,636 child, and 477 neonatal) were used for model comparison and performance evaluation. Further details on data preprocessing are provided in Appendix A.1, with COD and age group distributions summarized in Tables A1 and A2.

2.2 Modelling

Four computational models were used to assign causes of death (CODs) for each of the 6,939 physician-agreed verbal autopsy (VA) records: GPT-3.5, GPT-4, InterVA-5, and InSilicoVA. InterVA-5 and InSilicoVA are widely used statistical models within the OpenVA framework for COD assignment in VAs [13, 21, 22, 24, 25, 41–43]. InterVA-5 applies a Bayesian probabilistic approach, using a standardized set of symptoms and expert-derived conditional probabilities to assign the most likely COD based on maximum probability [20, 44, 45]. InSilicoVA extends this approach by incorporating a hierarchical Bayesian framework and Markov Chain Monte Carlo (MCMC) methods [46–48], allowing for quantification of uncertainty, individual-level probability estimates, and the integration of additional data sources [17]. GPT-3.5 [49] and GPT-4 [34] are large language models (LLMs) based on transformer architectures [50]. These models are trained using reinforcement learning from human feedback [51–54], enabling them to follow natural language instructions and generate human-level responses. GPT-4 introduces improvements over GPT-3.5, including more recent training data, enhanced reasoning capabilities, and multimodal input-output functionality (e.g. text, image, voice) [33].

For GPT-3.5 and GPT-4, the following user prompt was used to instruct each model to produce COD assignments as ICD-10 codes, where `<age>` and `<sex>` from the questionnaire, and `<narrative>` from the narratives, were replaced with values from the data:

```
Determine the underlying cause of death and provide the most
probable ICD-10 code for a verbal autopsy narrative of a <age>
years old <sex> death in Sierra Leone: <narrative>
```

InterVA-5 and InSilicoVA used structured questionnaire data, which were converted into OpenVA-compatible format [43]. Both models produced COD assignments coded using the WHO 2016 VA standard [55]. To ensure comparability across models, all

output CODs were mapped to the CGHR-10 classification system for evaluation relative to physician-assigned CODs. Further details on model input formats, output mappings, and code conversion procedures are provided in Appendix A.2.

2.3 Performance Evaluation

Model performance was assessed at both the population and individual levels by comparing each model’s CGHR-10 COD assignments to those of physicians for all 6,939 records. Cause-Specific Mortality Fraction (CSMF) accuracy was used to evaluate agreement at the population level (see Appendix A.3.1), while Partial Chance-Corrected Concordance (PCCC) was used to assess individual-level agreement (see Appendix A.3.2) [56]. Both metrics range from 0 to 1, where higher values indicate stronger similarity with physician assignment.

Given that model performance can vary by age and different CODs [41, 42, 57], both CSMF accuracy and PCCC were calculated overall and stratified by age group (adult, child, neonatal), CGHR-10 COD, and age at death. For adult and child groups, metrics were computed in five-year age bands for records with age at death of one year or older, and five-month bands for records between 28 days and one year. For the neonatal group, evaluations were conducted separately for age intervals of 0–6 days and 7–27 days. Additional details on the evaluation strategy and metric calculations are provided in Appendix A.3.

3 Results

This section presents the performance of GPT-3.5, GPT-4, InterVA-5, and InSilicoVA in assigning CGHR-10 CODs, based on the methodology described in Section 2. GPT-4 achieved the highest overall individual-level concordance, with a PCCC of 0.61, followed by GPT-3.5 (0.56). GPT-4 also demonstrated the highest PCCC across most age groups and CODs within the adult (12–69 years), child (28 days–11 years), and

neonatal (under 28 days) categories. In contrast, GPT-3.5, InterVA-5, and InSilicoVA showed higher PCCC values for a limited subset of age groups and CODs. Summary results are presented in Section 3.1, with stratified results by age group detailed in Sections 3.2, 3.3, and 3.4.

3.1 Overall Performance

Of all 6939 records, GPT-4 (0.61 PCCC) had the highest individual performance followed by GPT-3.5 (0.56 PCCC), InSilicoVA (0.44 PCCC), and InterVA-5 (0.44 PCCC) (Figure 2). GPT-3.5 and GPT-4 had improvements ranging from 0.14-0.18 PCCC over InSilicoVA and InterVA-5, while GPT-4 slightly improved over GPT-3.5 by 0.05 PCCC. Population level performances were similar for all models (0.74-0.79 CSMF). Figure 3 shows the PCCC performance across three age groups (adult, child, and neonate). GPT-4 had the best individual performance for adult and neonatal records (0.64 and 0.58 PCCC), while GPT-3.5 had the best performance for child records (0.54 PCCC) with GPT-4 performing slightly worse (0.51 PCCC). InSilicoVA and InterVA-5 performed the worse for adult and child records (≤ 0.5 PCCC), while GPT-3.5 performed the worse for neonatal records (0.42 PCCC). Performance varied less for child deaths (0.13 range) than for adult and neonatal deaths (0.24 and 0.22 range). Across ages, all models followed a similar pattern in individual performance (Figure 4), where PCCC trended upwards for 1 month to 14 years (~ 0.1 -0.75), and downwards for ages 15 to 69 years (~ 0.7 -0.35). The highest and lowest performances were observed for ages 12-29 years (~ 0.4 -0.7) and 1-11 months (~ 0.1 -0.35) respectively.

3.2 Performance for 3826 Adult Records (12 to 69 years)

Figure 5 presents model performance across 17 adult CODs, excluding suicide due to a low sample size ($n=3$, $<1\%$). GPT-4 achieved the highest individual level performance for 10 of 17 CODs (0.35–0.99 PCCC), followed by GPT-3.5 for 5 CODs (0.43–0.94




Fig. 2 Overall model performance.




Fig. 3 Model performance by age group.

PCCC), and InSilicoVA for 2 CODs (0.71 and 0.84 PCCC). InterVA-5 showed the lowest performance for 8 CODs (0–0.79 PCCC), InSilicoVA for 6 CODs (0.01–0.41 PCCC), and GPT-3.5 for 2 CODs (0.38 and 0.53 PCCC). The greatest improvements of GPT-3.5/4 over InSilicoVA and InterVA-5 were observed in chronic respiratory




Fig. 4 Model performance by age range.

diseases (+0.74-0.94 PCCC), while the smallest improvements were for malaria (+0.09-0.17 PCCC). All models achieved PCCC values above 0.70 for maternal conditions (0.79-0.99), but remained below 0.50 for unspecified infections (0.35-0.49), malaria (0.26-0.43), and ill-defined CODs (0-0.35). GPT-4 showed performance improvements exceeding 0.20 PCCC over all other models for cancers (+0.25-0.36), stroke (+0.27-0.45), and diarrhoeal diseases (+0.37-0.51). GPT-3.5 demonstrated similar gains for liver and alcohol-related diseases (+0.27-0.52). Performance variability across models was most pronounced for chronic respiratory diseases (range: 0.94), while narrower differences were observed for maternal conditions (0.20), malaria (0.17), ischemic heart disease (0.15), and unspecified infections (0.14).




Fig. 5 Model performance for adult records by COD.

3.3 Performance for 2636 Child Records (28 Days to 11 Years)

Figure 6 shows individual-level performance across 8 child CODs, excluding congenital anomalies due to a low sample size (n=1, <1%). GPT-4 achieved the highest

PCCC for 4 of the 8 CODs (0.65–0.94), followed by GPT-3.5 for 3 CODs (0.44–0.88), and InSilicoVA for 1 COD (0.78). InterVA-5 had the lowest performance for 4 CODs (0.09–0.79), InSilicoVA for 3 CODs (0–0.35), and GPT-3.5 for 1 COD (0.58). All models performed well for injuries, with PCCC values exceeding 0.70 (0.79–0.94), and showed lower performance for malaria (0.35–0.54) and other infections (0.29–0.44). GPT-4 demonstrated an improvement over other models for ill-defined CODs, with improvements greater than 0.30 PCCC (+0.38–0.65), and also showed stronger performance for injuries, with gains of +0.11–0.15 compared to +0.01–0.04 for other models. Performance differences exceeding 0.60 PCCC were observed for epilepsy, leukaemia, other communicable diseases (range: 0.73), ill-defined causes (0.65), and nutritional deficiencies (0.61). In contrast, narrower differences (less than 0.30 PCCC) were seen for malaria (0.20), injuries, and other infections (0.15).

3.4 Performance for 477 Neonatal Records (Under 28 Days)

Figure 7 shows model performance across 5 neonatal CODs, excluding congenital anomalies (n=2, <1%) and other causes (n=5, 1%) due to limited sample sizes. GPT-4 achieved the highest PCCC for 3 of the 5 CODs (0.39–0.71), while GPT-3.5 and InSilicoVA had the highest PCCC for 1 COD each (0.57 and 0.86). GPT-3.5 showed the lowest PCCC for 3 CODs (0–0.13), and InterVA-5 for 2 CODs (0.01 and 0.48). Performance was similar across all models for stillbirths (0.48–0.57 PCCC), though only GPT-4 achieved a PCCC greater than 0 for prematurity-related deaths. InSilicoVA outperformed all other models for neonatal infections, with gains of +0.18–0.73 PCCC. Performance differences greater than 0.6 PCCC were observed for infections (range: 0.73) and prematurity and low birthweight (0.7). Stillbirth showed minimal variation across models (range: 0.09).




Fig. 6 Model performance for child records by COD.

4 Discussion

This section interprets and contextualizes the findings presented in Section 3. The comparative advantages and limitations of GPT-3.5, GPT-4, InterVA-5, and InSilicoVA for COD assignment are discussed in Sections 4.1 and 4.2, respectively. Study limitations are outlined in Section 4.3, and directions for future research are presented in Section 4.4.




Fig. 7 Model performance for neonatal records by COD.

4.1 Advantages

This section outlines the strengths of the evaluated models in assigning CODs. Section 4.1.1 discusses model advantages across specific CODs and age groups. Section 4.1.3 highlights the potential for improving efficiency in physician-assisted COD assignment through computational support. Section 4.1.4 examines the benefits of leveraging natural language narratives in GPT models relative to traditional structured questionnaire data.

4.1.1 Cause-specific Models

At the population level, all models demonstrated comparable performance to physicians (0.74-0.79 CSMF), indicating their potential for estimating COD distributions in large populations. While individual-level performance was lower overall (0.44-0.61 PCCC), several models showed strong performance compared with physicians for specific CODs (up to 0.99 PCCC). GPT-3.5/4 consistently outperformed InSilicoVA and

InterVA-5 across most CODs, achieving the highest PCCC for 15 of 17 adult, 7 of 8 child, and 4 of 5 neonatal CODs. In contrast, InSilicoVA showed better performance for select CODs, including road and transport injuries (0.84 PCCC), tuberculosis (0.71), pneumonia (0.78), and neonatal infections (0.86). For CODs where high performance was observed, such as maternal conditions, chronic respiratory diseases, diabetes mellitus, and cancers for GPT-3.5/4 (0.91–0.99 PCCC), and road/transport injuries and neonatal infections for InSilicoVA (0.84 and 0.86 PCCC), the model outputs were more aligned with physician assignment. These findings support the potential utility of combining models based on their strengths for particular CODs. Evaluating performance at the COD level may allow for more targeted deployment of models, maximizing accuracy across disease categories. Table 1 illustrates how different models align with leading CODs identified in prior Sierra Leone studies [36, 58]. For example, we may deploy models to estimate asthma and chronic respiratory diseases using GPT-3 (0.94 PCCC), while using GPT-4 and InSilicoVA for diarrhoea and tuberculosis respectively (0.79 and 0.71 PCCC).

Table 1 Top ten leading causes of death for Sierra Leone in 2023 and most relevant models.

Top 10 Leading Cause of Death ¹ (~71% of ~76K deaths)	Deaths (% of 76K) ²	Best Model(s)	PCCC ³
Malaria	16,075 (21%)	GPT-3.5/4	0.46 (n=2181)
Infections	11,777 (16%)	GPT-3.5/4/InSilicoVA	0.55 (n=1155)
Ischaemic heart and other vascular	5,747 (8%)	GPT-4	0.65 (n=168)
Diarrhoea	4,285 (6%)	GPT-4	0.79 (n=380)
Stroke	4,262 (6%)	GPT-4	0.77 (n=331)
Pneumonia	3,074 (4%)	GPT-4/InSilicoVA	0.7 (n=186)
Birth asphyxia and birth trauma	2,431 (3%)	GPT-4	0.63 (n=103)
Tuberculosis	2,399 (3%)	InSilicoVA	0.71 (n=171)
Low birth weight/preterm	1,570 (2%)	GPT-4	0.71 (n=103)
Asthma and chronic respiratory	1,551 (2%)	GPT-3	0.94 (n=75)

¹Other infections and severe systemic/localized infections were generalized into infections. Appendix, hernia, intestinal and Peptic ulcer/gastroesophageal causes did not have comparable CGHR-10 codes and were omitted from the top ten.

²Percentage of ~76 Thousand (K) total deaths [58]. Numbers are rounded.

³Adult, child, and neonate mean PCCC and summed n records if available.

691 4.1.2 Age-specific Performance Patterns

692
693 Across age groups, all models exhibited a consistent upward trend in performance
694
695 from 6 months to 14 years, followed by a general decline from ages 15 to 69 years.
696
697 GPT-3.5/4 outperformed InSilicoVA and InterVA-5 throughout this range, while per-
698
699 formance patterns from birth to 5 months were more variable (see Figure 4). In adults,
700
701 performance generally decreased with age, suggesting greater difficulty in assigning
702
703 CODs among older adults, with a modest improvement observed after age 59. Among
704
705 children and neonates, performance increased beyond 5 months, indicating greater
706
707 model reliability as developmental age advanced. Although no model consistently
708
709 achieved performances greater than 0.8 PCCC in any specific five-year age band, these
710
711 age-related trends provide valuable insights. Specifically, they align with expectations
712
713 from clinical literature, where physicians often face greater diagnostic uncertainty in
714
715 neonatal cases [59, 60]. The observed patterns underscore the importance of consid-
716
717 ering developmental stage when interpreting model outputs and comparing them to
718
719 physician-assigned CODs.

716 4.1.3 Scalability and Availability

717
718 The models evaluated in this study offer scalable and cost-effective support for
719
720 physician-assigned CODs, particularly in resource-constrained settings. Similar to
721
722 tools used in differential diagnosis, GPT and InSilicoVA models can provide alterna-
723
724 tive COD suggestions for physician review [39], potentially reducing the proportion of
725
726 ill-defined causes and physician disagreement. At the time of analysis, running GPT-
727
728 3.5 on 6,939 records cost approximately \$1.60 USD (based on \$0.50 per million tokens),
729
730 while GPT-4 cost approximately \$115 USD (at \$30 per million tokens) [61]. InterVA-
731
732 5 and InSilicoVA were freely available as open-source software. These costs compare
733
734 favorably to physician review, which may exceed \$3 USD per household in settings
735
736

like India [15, 16], while the models can also process over 10,000 records within a single day. When physicians are unavailable, these models present a viable alternative for estimating population-level CODs. However, their application should be targeted to CODs where model performance is strong (see Table 1). Additionally, model outputs may be used to prioritize physician review, allocating less physician time to validating high-performing CODs (e.g. maternal conditions with 0.79–0.99 PCCC) and allocating more time to challenging cases (e.g. acute respiratory infections with 0.25–0.61 PCCC).

4.1.4 Natural Language Input and Output

None of the models required training data for COD assignment, enabling their use without domain-specific datasets or expertise. A key advantage of GPT-3.5/4 is their ability to process and generate natural language text as input and output. Unlike InterVA-5 and InSilicoVA, GPT models are able to assign CODs using the ICD-10 standard, mirroring physician practice, and can potentially classify CODs in broader or alternative categories based on prompt design. In contrast, InterVA-5 and InSilicoVA rely exclusively on structured data from WHO VA 2016 questionnaires and assign CODs using WHO VA 2016 codes. This dependency necessitates ongoing maintenance and conversion between questionnaire versions (e.g., WHO VA 2012 to 2016) and coding systems (e.g., WHO VA 2016 to ICD-10), which reduces interoperability and comparability across models. The flexibility of GPT models in handling unstructured data allows them to capture latent and ambiguous information—such as health-seeking behaviors and social context, which are not encompassed by standardized VA codes [26, 28]. For example, GPT-3.5/4 outperformed InterVA-5 and InSilicoVA by +0.35-0.65 PCCC on ill-defined CODs across age groups. They also demonstrated higher performance (+0.11-0.61 PCCC) on rarer CODs, such as nutritional deficiencies

(n=11) and diabetes mellitus (n=27), which may be underrepresented in questionnaire data, but better contextualized through extensive knowledge embedded in GPT training corpora.

4.2 Disadvantages

This subsection addresses the caveats of GPT models in COD assignment. Section 4.2.1 examines challenges related to reproducibility of GPT outputs across repeated runs and their dependence on static training data. Section 4.2.2 explores the substantial computational resources required by GPT models and the associated concerns regarding data privacy and security.

4.2.1 Reproducibility and Timeliness

In this study, GPT models were run with the temperature parameter set to 0 to enhance reproducibility and consistency. However, a brief experiment (Appendix B) showed that GPT-3.5 assigned the same COD for the same record in just over 60% of repeated runs on a sample of 100 records. This variability indicates that GPT models do not consistently produce identical COD assignments for identical inputs, which raises concerns about reproducibility and reliability. For example, GPT models may correctly assign CODs by chance, but extensive testing with large numbers of reruns (e.g., 10,000) is cost-prohibitive, as rerunning increases costs substantially. By contrast, InterVA-5 and InSilicoVA are open-source and free, enabling unlimited reruns without additional expense. Moreover, these models provide COD assignments with probabilities for alternative causes, enhancing reproducibility and transparency despite lower overall performance. Another important limitation common to all models is their reliance on training data that reflect information only up to a fixed point in time. Consequently, they may not incorporate the most current data sources, such as recent scientific literature, social media, or emerging reports. This lag can limit their ability

to detect new or emerging diseases (e.g., COVID-19) and shifts in COD distributions
related to outbreaks or other public health changes unless regularly updated.

4.2.2 Infrastructure and Data Privacy

GPT-3.5/4 require substantial computational infrastructure for training and inference,
making local deployment impractical due to cost and model ownership constraints.
Consequently, sensitive data, such as identifiable personal information, must be trans-
mitted to external servers, raising significant privacy concerns. Data submitted via
prompts, which include narrative content used for COD assignment, may be collected
by service providers (e.g., OpenAI) and potentially misused [62]. There is risk that
sensitive information could be exposed or exploited through malicious actors or poorly
controlled data handling [63, 64]. While jurisdictions, such as the European Union,
enforce strict protections under the General Data Protection Regulation (GDPR),
most low- and middle-income countries are only beginning to formalize regulatory
frameworks for data protection and artificial intelligence governance [65–67]. In con-
trast, InterVA-5 and InSilicoVA can be run entirely on local systems, enabling data
to remain under the control of the data owner. This approach reduces dependency on
external services and better safeguards data privacy.

4.3 Limitations

This section outlines key limitations of the current study related to the use of GPT
models. Section 4.3.1 discusses the use of physician assignment as the reference stan-
dard for comparing models. Section 4.3.2 addresses the need for further parameter
tuning and the evaluation of model consistency and multi-COD assignments, while
Section 4.3.3 notes the importance of more diverse datasets for model evaluation.

875 4.3.1 Physician Reference Standard

876
877 This study evaluated model performance using broad CGHR-10 categories rather than
878 specific ICD-10 codes. In practice, physicians assign more detailed ICD-10 codes, but
879 InterVA-5 and InSilicoVA generate only broader WHO VA codes and cannot assign
880 ICD-10 codes directly, partly due to insufficient sample cases for many specific ICD-10
881 categories to support reliable modeling. For example, even broad CGHR-10 cate-
882 gories had fewer than 10 cases (e.g., congenital anomalies, suicide), and were excluded
883 from evaluation. While GPT models assigned ICD-10 codes, lower performance can
884 be expected, as even physicians show limited agreement on detailed ICD-10 coding,
885 with only 6,939 (58%) of 11,920 records in agreement, necessitating the use of broader
886 categories (e.g., CMEA-10 codes) to assess equivalence. Reliance on physician assign-
887 ment as the reference standard may introduce bias, as physician interpretations may
888 be shaped by local epidemiological knowledge, particularly for more complex cases or
889 ambiguous narratives [22].

900 4.3.2 Model Tuning, Consistency, and Multiple Outputs

901
902 GPT-3.5/4 were used with default parameters except for temperature, which was set
903 to 0 to enhance consistency. However, tuning temperature and other settings could
904 potentially improve performance [68], but was not explored due to the high cost of
905 repeated runs needed for sensitivity analyses, as noted in Section 4.2.1. Despite tem-
906 perature control, GPT outputs may still vary, highlighting the need to assess reliability
907 and consistency to avoid coincidental results [69–71]. Unlike GPT models, InterVA-
908 5 and InSilicoVA provide multiple COD assignments with associated probabilities to
909 measure reliability. In addition, while GPT can be prompted to generate multiple
910 CODs, this study evaluated only the most probable assignment. Considering multiple
911 COD outputs may better capture alternative diagnoses and align more closely with
912 physician assessments [19].

920

4.3.3 Global Validity	921
	922
While this study rigorously compares computer algorithms for COD assignment in	923
Sierra Leone, the extent to which these findings are applied to other geographic or epi-	924
demiological contexts remains limited. Variations in local mortality profiles, linguistic	925
expression, health system infrastructure, and culturally specific interpretations of ill-	926
ness shape the content and structure of VA narratives and questionnaires [72–74]. For	927
example, the ways in which symptoms are described, terminology used, and aspects	928
emphasized by respondents differ across languages and cultural settings. Moreover,	929
Sierra Leone is predominantly driven by infectious diseases, such as malaria and res-	930
piratory infections, a pattern that contrasts with regions where non-communicable	931
diseases typically constitute the leading CODs in North America and Europe, or where	932
violence and road traffic injuries predominate in parts of Latin America and Asia [75–	933
77]. Given ongoing efforts to scale and integrate VA systems for mortality surveillance	934
across diverse low- and middle-income countries, further validation across globally rep-	935
resentative VA datasets is essential to evaluate model robustness, adaptability, and	936
operational utility in practice [78–80].	937
	938
	939
4.4 Opportunities	940
	941
This section explores opportunities to enhance GPT models for assigning CODs.	942
Section 4.4.1 highlights improvements through prompt engineering and analysis of mis-	943
classified cases. Section 4.4.2 discusses leveraging GPT to improve household survey	944
data quality. Section 4.4.3 considers integrating GPT with InterVA-5 and InSilicoVA	945
to support and enhance physician COD assignment within VA systems.	946
	947
	948
	949
	950
	951
	952
	953
	954
	955
	956
	957
	958
	959
	960
	961
	962
	963
	964
	965
	966

967 4.4.1 Prompt Engineering and Custom Models

968
969 Prompt engineering, the design of input prompts to guide GPT models toward
970 improved outputs [81], offers a key opportunity to enhance COD assignment per-
971 formance. An exploratory analysis in Appendix C of misclassified GPT-4 records
972 for neonatal infections identified potential issues related to CGHR-10 code catego-
973 rization, narrative information order, and COD assignment guidelines. Collaborating
974 with domain experts (e.g., physicians, specialists) to review misclassified cases could
975 inform prompt refinements that increase correct COD assignments or better align
976 with broader COD categories. Furthermore, iterative prompt adjustments incorporat-
977 ing additional questionnaire data and physician manuals (e.g., via retrieval-augmented
978 generation [82]) may improve model accuracy [83]. Sensitivity analyses can evaluate
979 how prompt modifications affect performance and output consistency on a cause-
980 specific basis. Additionally, GPT models can be customized to specific domains or
981 contexts, adjusting objectives, behavior, data inputs, privacy considerations, and eval-
982 uation criteria to create specialized models optimized for particular CODs or settings
983 [84].
984

995 4.4.2 Guided and Monitored Household Surveys

996
997 Verbal autopsies involve surveyors visiting households to collect information about
998 the deceased from family, friends, or community members. While standardized ques-
999 tionnaires are used, important latent information within free-text narratives often
1000 goes uncaptured [26, 28]. Narrative quality depends heavily on the surveyor’s social
1001 skills, cultural understanding, emotional capacity, and medical knowledge, all of which
1002 influence data completeness and potential bias [19, 85]. GPT models may support
1003 surveyors by suggesting improved or overlooked questions during interviews to elicit
1004 richer narratives. Moreover, as these models can assign CODs in real-time, they offer
1005 the opportunity to monitor data quality during collection. For example, by comparing
1006
1007
1008
1009
1010
1011
1012

estimated COD distributions with expected patterns for specific regions as a form of immediate quality control, where surveyors may be required to undergo review when estimated and expected COD distributions diverge significantly.

4.4.3 Computer Assisted Verbal Autopsy (CAVA)

This study establishes a basis for integrating GPT, InterVA-5, and InSilicoVA models into VA systems to support physicians in assigning CODs. In dual-coded VA systems (Section 2.1), two physicians independently assign CODs for each record and review each other’s assignments (reconciliation), while a senior physician adjudicates if disagreements persist. As noted in Section 4.1.3, presenting alternative COD suggestions from GPT and InSilicoVA models may reduce physician disagreement and the frequency of ill-defined records, allowing physicians to focus on more complex cases. Model-generated COD suggestions can be offered to physicians after their initial assignment, enabling reconsideration or confirmation of CODs (step 2 and option 2b in Figure 8). Future work will evaluate the impact of these suggestions on improving VA data quality, including increasing physician agreement and reducing ill-defined deaths. GPT-4, InterVA-5, and InSilicoVA suggestions have been incorporated into the ongoing HEAL-SL study [35], aiming to improve physician agreement and lower ill-defined COD assignments.




Fig. 8 Model suggestions integrated in the physician assignment process.

1059 5 Conclusion

1060
1061 This study evaluated the performance of GPT-3.5, GPT-4, InterVA-5, and InSili-
1062 coVA models against physicians in assigning CODs for 6,939 VA records from Sierra
1063 Leone (2019–2022). At the population level, all models achieved similar CSMF accu-
1064 racy (0.74–0.79). At the individual level, GPT-4 had the highest performance (0.61
1065 PCCC), followed by GPT-3.5 (0.58), and InSilicoVA/InterVA-5 (0.44). By COD,
1066 GPT-4 performed best for 10 of 17 adult, 4 of 8 child, and 3 of 5 neonatal causes,
1067 while GPT-3.5 led in 5 adult, 3 child, and 1 neonatal CODs, and InSilicoVA led in
1068 2 adult, 1 child, and 1 neonatal cause. Performance increased (~ 0.1 – 0.75 PCCC) as
1069 children and neonates matured (0 days to 14 years) and decreased (~ 0.7 – 0.35) with
1070 adult aging (15 to 69 years). These findings suggest that combining models tailored
1071 to specific CODs and age groups may optimize performance relative to physicians.
1072 All models demonstrated scalability and on-demand availability, enabling COD esti-
1073 mation and alternative diagnoses in low-resource or physician-scarce settings. GPT
1074 models’ natural language processing capability allowed flexible data input and out-
1075 put, aligning closer to physician reasoning, but issues remain with reproducibility,
1076 reliance on historical training data, computational demands, and data privacy. Study
1077 limitations included challenges comparing ICD-10 codes across models, limited sen-
1078 sitivity analyses due to costs, and exclusion of multiple COD assignment evaluation.
1079 Future research opportunities include prompt engineering and custom GPT models to
1080 improve accuracy, guided household surveys to enhance narrative quality, and CAVA
1081 systems integrating GPT and other models to support physicians by suggesting alter-
1082 native COD assignments. GPT-4, InterVA-5, and InSilicoVA have been incorporated
1083 into ongoing HEAL-SL study since 2022 to provide second-opinion support for physi-
1084 cian COD assignment. Evaluating the impact of computer-assisted VA on physician
1085 agreement and reduction of ill-defined deaths will be critical to advancing accurate,
1086 efficient VA systems worldwide.
1087
1088
1089
1090
1091
1092
1093
1094
1095
1096
1097
1098
1099
1100
1101
1102
1103
1104

Supplementary information.	Additional files were used to supplement this paper:	1105
		1106
•	Additional file 1: Centre for Global Health Research 10 (CGHR-10) codes. Codes	1107
	grouping ICD-10 code ranges into generalized categories. (.csv)	1108
		1109
•	Additional file 2: Central Medical Evaluation Agreement 10 (CMEA-10) codes. ICD-	1110
	10 code ranges considered in physician agreement. (.csv)	1111
		1112
		1113
Acknowledgments.	TBD.	1114
		1115
		1116
Declarations		1117
		1118
		1119
Funding		1120
		1121
TBD.		1122
		1123
		1124
Competing interests		1125
		1126
Not applicable.		1127
		1128
		1129
Ethics approval		1130
		1131
Not applicable.		1132
		1133
		1134
Consent for publication		1135
		1136
Not applicable.		1137
		1138
		1139
Availability of data and materials		1140
		1141
		1142
The datasets supporting the conclusions of this article are included within the article		1143
(and its additional files), at https://openmortality.org (available upon request). Verbal		1144
Autopsy (VA) and narrative data by age group and survey rounds 1 and 2 available at		1145
https://openmortality.org/dataset/heal-sl . Cause of death code mappings to convert		1146
		1147
		1148
		1149
		1150

1151 between ICD-10, WVA-2016, and CGHR-10 codes available at <https://openmortality.org/dataset/icd>.

1154

1155

1156 **Code availability**

1157

1158 All code for this paper is available at <https://github.com/cghr-toronto/healsl-gpt-paper>.

1160

1161

1162 **Authors' contributions**

1163

1164 PJ and PB are the study Principal Investigators. ATA and RK implemented the data
1165 collection procedures. RW, and TKS processed, documented, and prepared the data.
1166 RW, ASL, and RK ran the models. RW wrote the paper and conducted the analysis.
1167 AB and RCM provided medical domain guidance and feedback. All authors reviewed
1171 the results and contributed to the report. All authors read and approved the final
1172 manuscript.

1174

1175

1176 **Appendix A Details on Methods**

1177

1178 This section provides additional details on the methods described in Section 2. An
1179 overview of the methods used in this study is seen in Figure A1 as a five-step process.
1180 Section A.1 provides details on the preprocessed data used for modelling. Section A.2
1183 describes the data and parameter inputs and outputs for each model, while Section
1185 A.3 details the evaluation of model outputs at the individual and population level
1186 across different CODs, age groups, and ages.

1188

1189

1190

1191

1192

1193

1194

1195

1196




Fig. A1 Detailed study methods.

1243 **A.1 CGHR-10 Physician Agreed Records**

1244

1245 Initially, 11,920 records were collected from dual-coded EVA in the HEAL-SL study.
1246
1247 Physicians were able to assign CODs for 11,820 of the 11,920 records, where 100 of
1248 these records could not be assigned a COD due to missing or inadequate information
1249 (e.g. low quality narrative, data loss). The 11,820 physician coded records were further
1250 filtered for records where both physicians agreed on the assigned codes (records that
1251 were not reconciled or adjudicated) resulting in 6942 physician agreed records (based
1252 on comparisons using CMEA-10 codes, see Additional File 2). The 6942 records were
1253 converted into CGHR-10 codes (see Additional File 1) that generalized ICD-10 codes
1254 into 19, 10, and 7 categories for the adult (12 to 69 years), child (28 days to 11
1255 years), and neonatal (under 28 days) age groups. After conversion, a final total of
1256 6939 physician agreed records (3826 adult, 2636 child, and 477 neonatal) were used
1257 for modelling and performance evaluation, where three records were removed as their
1258 ICD-10 codes did not have a matching CGHR-10 code.
1259

1260 The 6939 physician agreed records were collected using VA from the HEAL-SL
1261 study between 2019-2022, where records were collected using nation wide samples
1262 across Sierra Leone provinces seen in Figure A2. More populous areas (e.g. southern
1263 and north east provinces with ~197,000 and ~135,000 population respectively) had
1264 more sampling areas versus less populous areas (e.g. north west and eastern provinces
1265 with ~50,000 and ~69,000 people respectively). The distribution of the study data are
1266 shown by CGHR-10 causes of death in Table A1. All age groups had relatively evenly
1267 distributed female and male records (44-55% of 6939 records each). Across CODs,
1268 there were noticeably more female records for cancers (65%), and maternal condi-
1269 tions (100%), while more male records for chronic respiratory diseases (61%), other
1270 noncommunicable diseases (61%), other injuries (77%), road and transport injuries
1271 (71%), and tuberculosis (68%). Most records were coded by physicians as malaria for
1272 adults (20%) and children (52%), and stillbirth (36%) and neonatal infections (21%)
1273
1274
1275
1276
1277
1278
1279
1280
1281
1282
1283
1284
1285
1286
1287
1288

for neonates. Suicide, congenital anomalies, nutritional deficiencies, and other had low sample sizes for each age group (<1% of total records for each age group). Table A2 shows the distribution of the study data by age. Across ages, there were more male records for 50-59 years (60-62%), while all other records had between 49-59% female and male records. Most records were in the 65-69 years age range for adults (15%), 1-5 years for children (62%), and 0-6 days for neonates (83%).




Fig. A2 Study data sampling areas.

A.2 Modelling Details

Each model (GPT-3.5, GPT-4, InSilicoVA, and InterVA-5) required pre-processing of the 6939 records into input data, and standardization of output COD codes from models for performance evaluation as not all models produced comparable codes across outputs. Although each model can assign multiple CODs per record, only the first

1335

1336 **Table A1** Study data by cause of death.

Age Group	CGHR-10 Cause of Death (COD)	Female	Male	Total
Adult, 18 CODs (n=3826, 55.1%)	Acute Respiratory Infections	48 (45.7%)	57 (54.3%)	105 (2.7%)
	Cancers	32 (65.3%)	17 (34.7%)	49 (1.3%)
	Chronic Respiratory Diseases	29 (38.7%)	46 (61.3%)	75 (2%)
	Diabetes Mellitus	14 (51.9%)	13 (48.1%)	27 (0.7%)
	Diarrhoeal Diseases	102 (49.8%)	103 (50.2%)	205 (5.4%)
	Ill-Defined	56 (47.9%)	61 (52.1%)	117 (3.1%)
	Ischemic Heart Disease	89 (53%)	79 (47%)	168 (4.4%)
	Liver And Alcohol Related Diseases	58 (45.3%)	70 (54.7%)	128 (3.3%)
	Malaria	372 (46.6%)	427 (53.4%)	799 (20.9%)
	Maternal Conditions	130 (100%)	N/A	130 (3.4%)
	Other Cardiovascular Diseases	59 (55.1%)	48 (44.9%)	107 (2.8%)
	Other Noncommunicable Diseases	160 (38.6%)	254 (61.4%)	414 (10.8%)
	Other Injuries	83 (23.2%)	274 (76.8%)	357 (9.3%)
	Road And Transport Injuries	73 (29.1%)	178 (70.9%)	251 (6.6%)
	Stroke	147 (44.4%)	184 (55.6%)	331 (8.7%)
	Suicide	N/A	3 (100%)	3 (0.1%)
	Tuberculosis	54 (31.6%)	117 (68.4%)	171 (4.5%)
	Unspecified Infections	175 (45%)	214 (55%)	389 (10.2%)
Child, 9 CODs (n=2636, 38%)	Congenital Anomalies	1 (100%)	N/A	1 (0%)
	Diarrhoeal Diseases	79 (45.1%)	96 (54.9%)	175 (6.6%)
	Epilepsy, Leukaemia, And	61 (53.5%)	53 (46.5%)	114 (4.3%)
	Other Noncommunicable Diseases			
	Ill-Defined	34 (48.6%)	36 (51.4%)	70 (2.7%)
	Injuries	51 (37.8%)	84 (62.2%)	135 (5.1%)
	Malaria	680 (49.2%)	702 (50.8%)	1382 (52.4%)
	Nutritional Deficiencies	7 (63.6%)	4 (36.4%)	11 (0.4%)
Child Male (n=1346, 51.1%)	Other Infections	338 (50.7%)	329 (49.3%)	667 (25.3%)
	Pneumonia	39 (48.1%)	42 (51.9%)	81 (3.1%)
Neonate, 7 CODs (n=477, 6.9%)	Birth Asphyxia And Birth Trauma	38 (36.9%)	65 (63.1%)	103 (21.6%)
	Congenital Anomalies	2 (100%)	N/A	2 (0.4%)
	Ill-Defined	11 (47.8%)	12 (52.2%)	23 (4.8%)
	Neonatal Infections	49 (49.5%)	50 (50.5%)	99 (20.8%)
	Other	2 (40%)	3 (60%)	5 (1%)
	Prematurity And Low Birthweight	39 (53.4%)	34 (46.6%)	73 (15.3%)
	Stillbirth	86 (50%)	86 (50%)	172 (36.1%)

1368

1369 generated COD response from GPT-3.5 and GPT-4, and the most probable COD

1370

1371 from InterVA-5 and InSilicoVA were used for evaluation. Section A.2.1 describes the

1372

1373 input data and parameters for each model, while Section A.2.3 details the outputs

1374

1375 from running each model.

1376

1377

1378

1379

1380

Table A2 Study data by age range.

Age Group	Age Range	Female	Male	Total
Adult (n=3826, 55.1%) Adult Female (n=1681, 43.9%) Adult Male (n=2145, 56.1%)	12-14 Years	51 (37.8%)	84 (62.2%)	135 (3.5%)
	15-19 Years	115 (42.8%)	154 (57.2%)	269 (7%)
	20-24 Years	146 (53.1%)	129 (46.9%)	275 (7.2%)
	25-29 Years	159 (45.2%)	193 (54.8%)	352 (9.2%)
	30-34 Years	174 (50.9%)	168 (49.1%)	342 (8.9%)
	35-39 Years	153 (45.4%)	184 (54.6%)	337 (8.8%)
	40-44 Years	134 (42%)	185 (58%)	319 (8.3%)
	45-49 Years	148 (47%)	167 (53%)	315 (8.2%)
	50-54 Years	134 (39.6%)	204 (60.4%)	338 (8.8%)
	55-59 Years	96 (37.6%)	159 (62.4%)	255 (6.7%)
Child (n=2636, 38%) Child Female (n=1290, 48.9%) Child Male (n=1346, 51.1%)	60-64 Years	128 (40.8%)	186 (59.2%)	314 (8.2%)
	65-69 Years	243 (42.3%)	332 (57.7%)	575 (15%)
	1-5 Months	146 (47.4%)	162 (52.6%)	308 (11.7%)
	6-11 Months	160 (50.8%)	155 (49.2%)	315 (11.9%)
Neonate (n=477, 6.9%) Neonate Female (n=227, 47.6%) Neonate Male (n=250, 52.4%)	1-5 Years	822 (50.3%)	811 (49.7%)	1633 (61.9%)
	6-11 Years	162 (42.6%)	218 (57.4%)	380 (14.4%)
	0-6 Days	184 (46.6%)	211 (53.4%)	395 (82.8%)
	7-27 Days	43 (52.4%)	39 (47.6%)	82 (17.2%)

A.2.1 Input Data and Preprocessing

For GPT-3.5 and GPT-4, 6939 text prompts were generated for each physician agreed record as input to instruct the models to assign CODs based on the open narratives.

Two types of text prompts were used: user prompts and system prompts. System prompts contained textual instructions to assign the role of a physician ICD-10 coder with expertise in Sierra Leone. The following system prompt was used for each record:

You are a physician with expertise in determining underlying causes of death in Sierra Leone by assigning the most probable ICD-10 code for each death using verbal autopsy narratives. Return only the ICD-10 code without description. E.g. A00. If there are multiple ICD-10 codes, show one code per line.

User prompts contained textual instructions to perform coding of VA records based on the age, sex, and narrative of the deceased. The following template was used to

1427 generate user prompts for each record, where `<age>` and `<sex>` from the questionnaire,
1428 and `<narrative>` from the narratives, were replaced with values from the data:

```
1430 Determine the underlying cause of death and provide the most  
1431 probable ICD-10 code for a verbal autopsy narrative of a <age>  
1432 years old <sex> death in Sierra Leone: <narrative>
```

1435 For InterVA-5 and InSilicoVA, the standardized questionnaire data from the HEAL-SL
1436 EVA were first converted into 2016 World Health Organization (WHO) VA question-
1437 naire revision 1.5.1 Open Data Kit (ODK) format [86, 87] consisting of 526 variables
1438 [88], followed by further conversion into OpenVA format [43] consisting of 353 vari-
1439 ables [89] using the `pyCrossVA` version 0.97 Python package [90]. The 6939 records
1440 were all converted into OpenVA formatted records for InterVA-5 and InSilicoVA.

1445

1447 A.2.2 Models and Parameters

1448

1449 The GPT-3.5 and GPT-4 Application Programming Interface (API) was accessed
1450 using Python version 3.11.4 and used to assign CODs for each record. GPT-3.5 used
1451 the `gpt-3.5-turbo` model, while GPT-4 used the `gpt-4-0613` model. The parameter
1452 `temperature` for GPT-3.5 and GPT-4, representing the sampling temperature ranging
1453 from 0 to 2 (default of 1), was set to 0 to produce more deterministic outputs [68].
1454 Higher values closer to 2 may produce less deterministic outputs, while lower values
1455 closer to 0 produce more deterministic outputs.

1456

1461 The `openVA` R package was used to run InterVA-5 and InSilicoVA models to assign
1462 CODs for each record in R version 4.3.1. The `openVA` package version 1.1.1 used
1463 dependent packages `InterVA5` version 1.1.3 and `InSilicoVA` version 1.4.0. The `Nsim`
1464 (number of iterations to run) parameter [91] for InSilicoVA was set to 9500, while
1465 the `HIV` (level of prevalence of human immunodeficiency virus) and `Malaria` (level
1466 of prevalence of Malaria) parameters [92] for InterVA-5 were both set to `'h'` (high)
1467 reflecting HIV and Malaria disease assumptions in Sierra Leone [93, 94]. Note that the
1471

default value of `Nsim=10000` for InSilicoVA ran until 9500 iterations before it stopped due to errors, thus `Nsim=9500` was used and ran successfully for all iterations.

A.2.3 Output Data and Code Conversion

Of the 6939 input records, GPT-3.5, GPT-4, InterVA-5, and InSilicoVA were able to assign CODs for 6939 (100%), 6935 (>99%), 6830 (98%), 6830 (98%) records respectively. All 6830 (100%) InterVA-5 and InSilicoVA records with WHO VA 2016 v1.5 output codes [55] were converted into ICD-10 codes respectively. After all model outputs were converted to ICD-10 codes, they were further converted to CGHR-10 codes. The 6939 GPT-3.5 and 6935 GPT-4 output records with ICD-10 codes were converted into 6930 (>99%) and 6931 (>99) records with CGHR-10 codes, where <1% (9 and 8) records did not have matching CGHR-10 codes respectively. The 6830 InterVA-5 and InSilicoVA records with ICD-10 codes were converted into 6802 (>99%) and 6726 (98%) records with CGHR-10 codes respectively, where 28 (<1%) and 104 (1%) of records could not be converted into CGHR-10 codes.

A.3 Performance Evaluation Details

The performance of GPT-3.5, GPT-4, InSilicoVA, and InterVA-5 models were evaluated with metrics at the population and individual level by comparing their CGHR-10 COD outputs for 6939 records to physician COD assignments. Section A.3.1 describes CSMF accuracy in detail for evaluating models on the population level, Section A.3.2 describes PCCC for evaluating models on the individual level. Records that were assigned a COD by physicians, but not by a model were considered to be an incorrect COD assignment by the model. CSMF accuracy and PCCC were calculated for each model overall and by three age groups (adult, child, and neonatal), then further into age and COD for each age group.

1519 **A.3.1 Cause Specific Mortality Fraction (CSMF) Accuracy**

1520

1521 CSMF accuracy measures the performance of models at the population level, compar-
 1522 ing distributions of CODs between the physicians and the models [56]. To calculate
 1523 CSMF accuracy, $CSMF_j$ was calculated as is the fraction of physician or model records
 1524 for cause j , given by dividing the number of records for cause j with the total number
 1525 of records as seen in Equation A1. Then, the $CSMF_{MaximumError}$, representing
 1526 the worst possible model, is calculated using Equation A2. Finally, the CSMF accuracy
 1527 is given by Equation A3, where k is the number of causes, j is a cause, $CSMF_j^{true}$ is
 1528 the true physician CSMF for cause j , and $CSMF_j^{pred}$ is the prediction model CSMF
 1529 for cause j . CSMF accuracy ranges from 0 to 1, where 1 means that the model com-
 1530 pletely matched the physician COD distribution and 0 means that it did not match
 1531 the distribution at all.

1532

1533

$$1541 \quad CSMF_j = Records_j / Records \quad (A1)$$

1542

1543

$$1544 \quad CSMF_{MaximumError} = 2(1 - \min(CSMF_j^{true})) \quad (A2)$$

1545

1546

$$1547 \quad CSMF_{Accuracy} = 1 - \frac{\sum_{j=1}^k |CSMF_j^{true} - CSMF_j^{pred}|}{CSMF_{MaximumError}} \quad (A3)$$

1548

1549

1551 **A.3.2 Partial Chance Corrected Concordance (PCCC)**

1552

1553 PCCC measures the performance of models at the individual level, comparing COD
 1554 assignments between the physicians and models on a record by record basis, correcting
 1555 for COD assignments made purely by chance [56]. PCCC is given by Equation A5,
 1556 where k is the number of top COD assignments from the model to consider, N is
 1557 number of causes, and C is fraction of records where the physician COD assignment is
 1558 one of the top COD assignments from the model. For this study, k was set to 1, making
 1559 C equivalent to the fraction of true positives TP or records where the physician COD
 1560

assignment is equal to the model COD assignment as shown in Equation A4. Higher PCCC values closer to 1 indicate that model COD assignments are similar to physician COD assignments, while values closer to 0 indicate that model COD assignments are not similar to physicians.

$$C = \frac{TP}{Records} \quad (A4)$$

$$PCCC(k) = \frac{C - \frac{k}{N}}{1 - \frac{k}{N}} \quad (A5)$$

Appendix B Experiment on Repeated Runs of GPT-3.5

A short experiment was conducted to test the consistency of GPT-3.5 outputs repeated on the same record. 100 records, sampled randomly with approximately equal proportions across age groups, CODs, and survey rounds 1 and 2, were used to test repeated runs of GPT-3.5. Each record from the 100 records was rerun 10 times through GPT-3.5, resulting in ten COD outputs per record. The ICD-10 codes were then converted to CGHR-10 codes and tested for consistency, where completely inconsistent results had different ICD-10 or CGHR-10 codes for each of the 10 reruns (1 times+), and completely consistent results had the same ICD-10 or CGHR-10 code for all 10 reruns (10 times), on the same record.

The results are shown in Table B3. For all 100 records, GPT-3.5 assigns the same ICD-10 and CGHR-10 code for the same record 5 times or more out of 10. For 66 and 79 records, GPT-3.5 assigns the same ICD-10 and CGHR-10 code respectively for each record. This number increases to 94 (from 66) and 96 (from 79) when reducing the number of times out of 10 that GPT-3.5 assigns the same ICD-10 and CGHR-10 code respectively. Thus, GPT-3.5 does not always produce the same outputs when repeated on the same record (10 times out of 10), even when the temperature is set

to 0, but does so for more than half the records. For most records (more than 90%), GPT-3.5 will produce the same outputs for the same record 7 times or more out of 10.

Table B3 Records with same GPT-3.5 outputs based on 10 repeated reruns of 100 records

Times with Same GPT-3.5 Outputs	ICD-10 Records	CGHR-10 Records
1 times+ (inconsistent)	100	100
2 times+	100	100
3 times+	100	100
4 times+	100	100
5 times+	100	100
6 times+	94	96
7 times+	92	94
8 times+	86	91
9 times+	79	86
10 times (consistent)	66	79

Appendix C Exploration of Neonatal Infections

An exploration of neonatal infections (n=99, 21% of 477 records) was done to understand the low performance of GPT models (0.23 PCCC) for neonatal infections, and high performance of InSilicoVA (0.87 PCCC). In Table C4, about half the records were assigned correctly, and a majority (n=33, 33%) of the other records were misclassified as other, while prematurity and low birthweight, birth asphyxia & birth trauma, and ill-defined make up the rest. On closer inspection of the 49 records with misclassified assignments, the ICD-10 code R50 was assigned in 20 records. R50 falls under unspecified infections in the adult CGHR-10 category, but in the other category for neonates. B50 was assigned in 4 records, falling under malaria, but a similar B54 falls under neonatal infections. P81 was assigned in 3 records, referring to fever of unknown origin, which falls under other, and P07 was assigned in 7 records, falling under prematurity and low birthweight.

In most misclassified records, there is mention of infections, but the misclassifications occur due to the finer details of the ICD-10 code classifications, the categorization

decisions of the CGHR-10 codes, and missing information from the questionnaire. For R50 misclassifications, GPT may have confused descriptions across adult and neonatal age groups. Using the same definition of R50, but in the context of neonates, may result in codes closer to neonatal infections (e.g. B54). For B50 misclassifications, the similar B54 was categorized in CGHR-10 as neonatal infections, but B50 was categorized as other. P81 refers to fever of unknown origin, which may be difficult to differentiate between infection and other causes without information from the questionnaire. P07 refers to prematurity and low birthweight, where GPT initially assigned P07 as the age of the neonate was mentioned first, but later mentions infections as an alternative following the order of information in the narratives. Thus, it may be possible to improve the performance GPT models using better prompts based on the context of VA manuals and CGHR-10 codes, and by also including questionnaire information in the prompts.

Table C4 GPT-4 CGHR-10 COD assignment for physician coded neonatal infections records.

GPT-4 Assigned Cause of Death (CGHR-10)	Records
Neonatal infections	50 (51%)
Other	33 (33%)
Prematurity and low birthweight	9 (9%)
Birth asphyxia & birth trauma	5 (6%)
Ill-defined	2 (2%)
Total	99 (100%)

1703 References

- 1704
- 1705 [1] World Health Organization.: Non Communicable Diseases: Key Facts.
- 1706 <https://www.who.int/news-room/fact-sheets/detail/noncommunicable-diseases>.
- 1707
- 1708
- 1709 [2] Benziger CP, Roth GA, Moran AE. The Global Burden of Disease Study and
- 1710 the Preventable Burden of NCD. *Global Heart*. 2016 Dec;11(4):393–397. <https://doi.org/10.1016/j.gheart.2016.10.024>.
- 1711
- 1712
- 1713
- 1714
- 1715 [3] Lawn JE, Kerber K, Enweronu-Laryea C, Cousens S. 3.6 Million Neonatal
- 1716 Deaths—What Is Progressing and What Is Not? *Seminars in Perinatology*. 2010
- 1717 Dec;34(6):371–386. <https://doi.org/10.1053/j.semperi.2010.09.011>.
- 1718
- 1719
- 1720
- 1721 [4] Lassi ZS, Bhutta ZA. Community-based Intervention Packages for Reducing
- 1722 Maternal and Neonatal Morbidity and Mortality and Improving Neonatal Out-
- 1723 comes. *Cochrane Database of Systematic Reviews*. 2015;(3). [https://doi.org/10.](https://doi.org/10.1002/14651858.CD007754.pub3)
- 1724 [1002/14651858.CD007754.pub3](https://doi.org/10.1002/14651858.CD007754.pub3).
- 1725
- 1726
- 1727
- 1728
- 1729 [5] Liu NH, Daumit GL, Dua T, Aquila R, Charlson F, Cuijpers P, et al. Excess
- 1730 Mortality in Persons with Severe Mental Disorders: A Multilevel Intervention
- 1731 Framework and Priorities for Clinical Practice, Policy and Research Agendas.
- 1732 *World Psychiatry*. 2017;16(1):30–40. <https://doi.org/10.1002/wps.20384>.
- 1733
- 1734
- 1735
- 1736 [6] Ewig S, Torres A. Community-Acquired Pneumonia as an Emergency: Time for
- 1737 an Aggressive Intervention to Lower Mortality. *European Respiratory Journal*.
- 1738 2011 Aug;38(2):253–260. <https://doi.org/10.1183/09031936.00199810>.
- 1739
- 1740
- 1741
- 1742 [7] World Health Organization. SCORE for Health Data Technical Package: Global
- 1743 Report on Health Data Systems and Capacity, 2020; 2021.
- 1744
- 1745
- 1746
- 1747
- 1748

- [8] de Savigny D, Riley I, Chandramohan D, Odhiambo F, Nichols E, Notzon S, et al. Integrating Community-Based Verbal Autopsy into Civil Registration and Vital Statistics (CRVS): System-Level Considerations. *Global Health Action*. 2017 Jan;10(1):1272882. <https://doi.org/10.1080/16549716.2017.1272882>.
- [9] Thomas LM, D’Ambruoso L, Balabanova D. Verbal Autopsy in Health Policy and Systems: A Literature Review. *BMJ Global Health*. 2018 May;3(2):e000639. <https://doi.org/10.1136/bmjgh-2017-000639>.
- [10] Rampatige R, Mikkelsen L, Hernandez B, Riley I, Lopez AD. Systematic Review of Statistics on Causes of Deaths in Hospitals: Strengthening the Evidence for Policy-Makers. *Bulletin of the World Health Organization*. 2014 Sep;92:807–816. <https://doi.org/10.2471/BLT.14.137935>.
- [11] Adair T. Who Dies Where? Estimating the Percentage of Deaths That Occur at Home. *BMJ Global Health*. 2021 Sep;6(9):e006766. <https://doi.org/10.1136/bmjgh-2021-006766>.
- [12] World Health Organization. Verbal Autopsy Standards: 2022 WHO Verbal Autopsy Instrument; 2023.
- [13] Chandramohan D, Fottrell E, Leitao J, Nichols E, Clark SJ, Alsokhn C, et al. Estimating Causes of Death Where There Is No Medical Certification: Evolution and State of the Art of Verbal Autopsy. *Global Health Action*. 2021 Oct;14(sup1):1982486. <https://doi.org/10.1080/16549716.2021.1982486>.
- [14] World Health Organization. Verbal Autopsy Standards: Ascertaining and Attributing Cause of Death. World Health Organization; 2007.
- [15] Gomes M, Begum R, Sati P, Dikshit R, Gupta PC, Kumar R, et al. Nationwide Mortality Studies To Quantify Causes Of Death: Relevant Lessons From India’s

1795 Million Death Study. Health Affairs. 2017 Nov;36(11):1887–1895. [https://doi.](https://doi.org/10.1377/hlthaff.2017.0635)
1796 [org/10.1377/hlthaff.2017.0635](https://doi.org/10.1377/hlthaff.2017.0635).
1797
1798
1799 [16] Jha P, Gajalakshmi V, Gupta PC, Kumar R, Mony P, Dhingra N, et al. Prospec-
1800 tive Study of One Million Deaths in India: Rationale, Design, and Validation
1801 Results. PLOS Medicine. 2005 Dec;3(2):e18. [https://doi.org/10.1371/journal.](https://doi.org/10.1371/journal.pmed.0030018)
1802 [pmed.0030018](https://doi.org/10.1371/journal.pmed.0030018).
1803
1804
1805
1806 [17] McCormick TH, Li ZR, Calvert C, Crampin AC, Kahn K, Clark SJ. Prob-
1807 abilistic Cause-of-death Assignment Using Verbal Autopsies. Journal of the
1808 American Statistical Association. 2016;111(515):1036–1049. [https://doi.org/10.](https://doi.org/10.1080/01621459.2016.1152191)
1809 [1080/01621459.2016.1152191](https://doi.org/10.1080/01621459.2016.1152191).
1810
1811
1812
1813
1814 [18] Morris SK, Bassani DG, Kumar R, Awasthi S, Paul VK, Jha P. Factors Associated
1815 with Physician Agreement on Verbal Autopsy of over 27000 Childhood Deaths in
1816 India. PloS one. 2010;5(3):e9583.
1817
1818
1819
1820 [19] Soleman N, Chandramohan D, Shibuya K. Verbal Autopsy: Current Practices
1821 and Challenges. Bulletin of the World Health Organization. 2006;84(3):239–245.
1822
1823
1824 [20] Byass P, Hussain-Alkhateeb L, D’Ambruso L, Clark S, Davies J, Fottrell E,
1825 et al. An Integrated Approach to Processing WHO-2016 Verbal Autopsy Data:
1826 The InterVA-5 Model. BMC Medicine. 2019 May;17(1):102. [https://doi.org/10.](https://doi.org/10.1186/s12916-019-1333-6)
1827 [1186/s12916-019-1333-6](https://doi.org/10.1186/s12916-019-1333-6).
1828
1829
1830
1831 [21] Jha P, Kumar D, Dikshit R, Budukh A, Begum R, Sati P, et al. Automated versus
1832 Physician Assignment of Cause of Death for Verbal Autopsies: Randomized Trial
1833 of 9374 Deaths in 117 Villages in India. BMC Medicine. 2019 Jun;17(1):116.
1834
1835
1836 <https://doi.org/10.1186/s12916-019-1353-2>.
1837
1838
1839
1840

- [22] Leitao J, Desai N, Aleksandrowicz L, Byass P, Miasnikof P, Tollman S, et al. Comparison of Physician-Certified Verbal Autopsy with Computer-Coded Verbal Autopsy for Cause of Death Assignment in Hospitalized Patients in Low- and Middle-Income Countries: Systematic Review. BMC Medicine. 2014 Feb;12(1):22. <https://doi.org/10.1186/1741-7015-12-22>.
- [23] Desai N, Aleksandrowicz L, Miasnikof P, Lu Y, Leitao J, Byass P, et al. Performance of Four Computer-Coded Verbal Autopsy Methods for Cause of Death Assignment Compared with Physician Coding on 24,000 Deaths in Low- and Middle-Income Countries. BMC Medicine. 2014 Feb;12(1):20. <https://doi.org/10.1186/1741-7015-12-20>.
- [24] Tunga M, Lungo J, Chambua J, Kateule R. Verbal Autopsy Models in Determining Causes of Death. Tropical Medicine & International Health. 2021;26(12):1560–1567. <https://doi.org/10.1111/tmi.13678>.
- [25] Oti SO, Kyobutungi C. Verbal Autopsy Interpretation: A Comparative Analysis of the InterVA Model versus Physician Review in Determining Causes of Death in the Nairobi DSS. Population Health Metrics. 2010 Jun;8(1):21. <https://doi.org/10.1186/1478-7954-8-21>.
- [26] Jeblee S, Gomes M, Jha P, Rudzicz F, Hirst G. Automatically Determining Cause of Death from Verbal Autopsy Narratives. BMC Medical Informatics and Decision Making. 2019 Jul;19(1):127. <https://doi.org/10.1186/s12911-019-0841-9>.
- [27] Blanco A, Pérez A, Casillas A, Cobos D. Extracting Cause of Death From Verbal Autopsy With Deep Learning Interpretable Methods. IEEE Journal of Biomedical and Health Informatics. 2021 Apr;25(4):1315–1325. <https://doi.org/10.1109/JBHI.2020.3005769>.

1887 [28] King C, Zamawe C, Banda M, Bar-Zeev N, Beard J, Bird J, et al. The Quality and
1888 Diagnostic Value of Open Narratives in Verbal Autopsy: A Mixed-Methods Anal-
1889 ysis of Partnered Interviews from Malawi. BMC Medical Research Methodology.
1890 2016 Feb;16(1):13. <https://doi.org/10.1186/s12874-016-0115-5>.
1891
1892 [29] Chang Y, Wang X, Wang J, Wu Y, Yang L, Zhu K, et al.: A Survey on Evaluation
1893 of Large Language Models. arXiv.
1894
1895 [30] Lund BD, Wang T. Chatting about ChatGPT: How May AI and GPT Impact
1896 Academia and Libraries? Library Hi Tech News. 2023 Jan;40(3):26–29. <https://doi.org/10.1108/LHTN-01-2023-0009>.
1897
1898 [31] Svyatkovskiy A, Deng SK, Fu S, Sundaresan N. IntelliCode Compose: Code
1899 Generation Using Transformer. In: Proceedings of the 28th ACM Joint Meeting on
1900 European Software Engineering Conference and Symposium on the Foundations
1901 of Software Engineering. ESEC/FSE 2020. New York, NY, USA: Association for
1902 Computing Machinery; 2020. p. 1433–1443.
1903
1904 [32] Haupt CE, Marks M. AI-Generated Medical Advice—GPT and Beyond. JAMA.
1905 2023 Apr;329(16):1349–1350. <https://doi.org/10.1001/jama.2023.5321>.
1906
1907 [33] Wu T, He S, Liu J, Sun S, Liu K, Han QL, et al. A Brief Overview of ChatGPT:
1908 The History, Status Quo and Potential Future Development. IEEE/CAA Journal
1909 of Automatica Sinica. 2023;10(5):1122–1136. [https://doi.org/10.1109/JAS.2023.](https://doi.org/10.1109/JAS.2023.123618)
1910 123618.
1911
1912 [34] OpenAI, Achiam J, Adler S, Agarwal S, Ahmad L, Akkaya I, et al.: GPT-4
1913 Technical Report. arXiv.
1914
1915 [35] Njala University.: Healthy Sierra Leone. <https://healsl.org/>.
1916
1917
1918
1919
1920
1921
1922
1923
1924
1925
1926
1927
1928
1929
1930
1931
1932

[36] Carshon-Marsh R, Aimone A, Ansumana R, Swaray IB, Assalif A, Musa A, et al. Child, Maternal, and Adult Mortality in Sierra Leone: Nationally Representative Mortality Survey 2018–20. *The Lancet Global Health*. 2022 Jan;10(1):e114–e123. [https://doi.org/10.1016/S2214-109X\(21\)00459-9](https://doi.org/10.1016/S2214-109X(21)00459-9).

[37] World Health Organization. ICD-10: International Statistical Classification of Diseases and Related Health Problems (10th Revision); 2011.

[38] Aleksandrowicz L, Malhotra V, Dikshit R, Gupta PC, Kumar R, Sheth J, et al. Performance Criteria for Verbal Autopsy-Based Systems to Estimate National Causes of Death: Development and Application to the Indian Million Death Study. *BMC Medicine*. 2014 Feb;12(1):21. <https://doi.org/10.1186/1741-7015-12-21>.

[39] Barnett ML, Boddupalli D, Nundy S, Bates DW. Comparative Accuracy of Diagnosis by Collective Intelligence of Multiple Physicians vs Individual Physicians. *JAMA Network Open*. 2019 Mar;2(3):e190096. <https://doi.org/10.1001/jamanetworkopen.2019.0096>.

[40] Hsiao M, Morris SK, Bassani DG, Montgomery AL, Thakur JS, Jha P. Factors Associated with Physician Agreement on Verbal Autopsy of over 11500 Injury Deaths in India. *PLOS ONE*. 2012 Jan;7(1):e30336. <https://doi.org/10.1371/journal.pone.0030336>.

[41] Murray CJ, Lozano R, Flaxman AD, Serina P, Phillips D, Stewart A, et al. Using Verbal Autopsy to Measure Causes of Death: The Comparative Performance of Existing Methods. *BMC Medicine*. 2014 Jan;12(1):5. <https://doi.org/10.1186/1741-7015-12-5>.

1979 [42] Benara SK, Sharma S, Juneja A, Nair S, Gulati BK, Singh KJ, et al. Evaluation of
1980
1981 Methods for Assigning Causes of Death from Verbal Autopsies in India. *Frontiers*
1982 in Big Data. 2023 Aug;6:1197471. <https://doi.org/10.3389/fdata.2023.1197471>.
1983
1984
1985 [43] Li ZR, Thomas J, Choi E, McCormick TH, Clark SJ. The openVA Toolkit for
1986 Verbal Autopsies. *The R Journal*. 2023 Feb;p. 1.
1987
1988
1989 [44] Byass P, Chandramohan D, Clark SJ, D’Ambruoso L, Fottrell E, Graham WJ,
1990 et al. Strengthening Standardised Interpretation of Verbal Autopsy Data: The
1991 New InterVA-4 Tool. *Global Health Action*. 2012 Dec;5(1):19281. [https://doi.](https://doi.org/10.3402/gha.v5i0.19281)
1992
1993 [org/10.3402/gha.v5i0.19281](https://doi.org/10.3402/gha.v5i0.19281).
1994
1995
1996 [45] BAYES. An Essay towards Solving a Problem in the Doctrine of Chances.
1997 *Biometrika*. 1958;45(3-4):296–315.
1998
1999
2000 [46] Brooks S. Markov Chain Monte Carlo Method and Its Application. *Journal of*
2001 the Royal Statistical Society: Series D (The Statistician). 1998 Mar;47(1):69–100.
2002
2003 <https://doi.org/10.1111/1467-9884.00117>.
2004
2005
2006 [47] Chib S. Markov Chain Monte Carlo Methods: Computation and Inference.
2007 *Handbook of econometrics*. 2001;5:3569–3649.
2008
2009
2010 [48] Han C, Carlin BP. Markov Chain Monte Carlo Methods for Computing Bayes
2011 Factors: A Comparative Review. *Journal of the American Statistical Association*.
2012 2001 Sep;96(455):1122–1132. <https://doi.org/10.1198/016214501753208780>.
2013
2014
2015
2016 [49] Brown TB, Mann B, Ryder N, Subbiah M, Kaplan J, Dhariwal P, et al.: Language
2017 Models Are Few-Shot Learners. *arXiv*.
2018
2019
2020 [50] Vaswani A, Shazeer N, Parmar N, Uszkoreit J, Jones L, Gomez AN, et al. Atten-
2021 tion Is All You Need. In: *Advances in Neural Information Processing Systems*.
2022
2023
2024

vol. 30. Curran Associates, Inc.; 2017. . 2025

[51] Ouyang L, Wu J, Jiang X, Almeida D, Wainwright CL, Mishkin P, et al.: Training 2026
Language Models to Follow Instructions with Human Feedback. arXiv. 2027
2028

[52] Christiano PF, Leike J, Brown T, Martic M, Legg S, Amodei D. Deep Rein- 2031
forcement Learning from Human Preferences. Advances in neural information 2032
processing systems. 2017;30. 2033
2034
2035
2036

[53] Stiennon N, Ouyang L, Wu J, Ziegler D, Lowe R, Voss C, et al. Learning to 2037
Summarize with Human Feedback. Advances in Neural Information Processing 2038
Systems. 2020;33:3008–3021. 2039
2040
2041
2042

[54] Wirth C, Akrou R, Neumann G, Fürnkranz J. A Survey of Preference-Based 2043
Reinforcement Learning Methods. The Journal of Machine Learning Research. 2044
2045
2046
2047
2048

[55] World Health Organization.: Verbal Autopsy Standards: The 2016 WHO Ver- 2049
bal Autopsy Instrument. [https://www.who.int/publications/m/item/verbal-](https://www.who.int/publications/m/item/verbal-autopsy-standards-the-2016-who-verbal-autopsy-instrument) 2050
[autopsy-standards-the-2016-who-verbal-autopsy-instrument.](https://www.who.int/publications/m/item/verbal-autopsy-standards-the-2016-who-verbal-autopsy-instrument) 2051
2052
2053
2054

[56] Murray CJ, Lozano R, Flaxman AD, Vahdatpour A, Lopez AD. Robust Metrics 2055
for Assessing the Performance of Different Verbal Autopsy Cause Assignment 2056
Methods in Validation Studies. Population Health Metrics. 2011 Aug;9(1):28. 2057
2058
2059
<https://doi.org/10.1186/1478-7954-9-28>. 2060
2061

[57] Setel PW, Whiting DR, Hemed Y, Chandramohan D, Wolfson LJ, Alberti 2062
KGMM, et al. Validity of Verbal Autopsy Procedures for Determining Cause of 2063
Death in Tanzania. Tropical Medicine & International Health. 2006;11(5):681– 2064
696. <https://doi.org/10.1111/j.1365-3156.2006.01603.x>. 2065
2066
2067
2068
2069
2070

2071 [58] Ansumana R, Mohamed V, Carshon-Marsh R, Jambai A, Smart F, Sartie K,
2072 et al. Report on Causes of Death in Sierra Leone 2018 – 2023; 2023.
2073
2074

2075 [59] Rasmussen LA, Cascio MA, Ferrand A, Shevell M, Racine E. The Complex-
2076 ity of Physicians’ Understanding and Management of Prognostic Uncertainty
2077 in Neonatal Hypoxic-Ischemic Encephalopathy. *Journal of Perinatology*. 2019
2078 Feb;39(2):278–285. <https://doi.org/10.1038/s41372-018-0296-3>.
2079
2080
2081

2082 [60] Faison G, Chou FS, Feudtner C, Janvier A. When the Unknown Is Unknowable:
2083 Confronting Diagnostic Uncertainty. *Pediatrics*. 2023 Sep;152(4):e2023061193.
2084
2085 <https://doi.org/10.1542/peds.2023-061193>.
2086
2087

2088 [61] OpenAI.: Pricing. <https://openai.com/api/pricing/>.
2089
2090

2091 [62] Tao G, Cheng S, Zhang Z, Zhu J, Shen G, Zhang X.: Opening A Pandora’s Box:
2092 Things You Should Know in the Era of Custom GPTs. *arXiv*.
2093
2094

2095 [63] Khowaja SA, Khuwaja P, Dev K, Wang W, Nkenyereye L. ChatGPT
2096 Needs SPADE (Sustainability, PrivAcY, Digital Divide, and Ethics) Evalua-
2097 tion: A Review. *Cognitive Computation*. 2024 May;[https://doi.org/10.1007/](https://doi.org/10.1007/s12559-024-10285-1)
2098
2099 [s12559-024-10285-1](https://doi.org/10.1007/s12559-024-10285-1).
2100
2101

2102 [64] Wu X, Duan R, Ni J. Unveiling Security, Privacy, and Ethical Concerns of
2103 ChatGPT. *Journal of Information and Intelligence*. 2024;2(2):102–115.
2104
2105

2106 [65] Intersoft Consulting.: General Data Protection Regulation (GDPR) – Legal Text.
2107
2108 <https://gdpr-info.eu/>.
2109
2110

2111 [66] Beck EJ, G Wayne, and De Lay PR. Protecting the Confidentiality and Security
2112 of Personal Health Information in Low- and Middle-Income Countries in the Era
2113 of SDGs and Big Data. *Global Health Action*. 2016 Dec;9(1):32089. [https://doi.](https://doi.org/10.1080/17445019.2016.1210891)
2114
2115
2116

org/10.3402/gha.v9.32089. 2117
2118

[67] Kwarkye TG. “We Know What We Are Doing”: The Politics and Trends in 2119
Artificial Intelligence Policies in Africa. Canadian Journal of African Studies / 2120
Revue canadienne des études africaines. 2025;p. 1–19. [https://doi.org/10.1080/](https://doi.org/10.1080/00083968.2025.2456619) 2121
[00083968.2025.2456619](https://doi.org/10.1080/00083968.2025.2456619). 2122
2123
2124
2125
2126

[68] OpenAI.: OpenAI Platform: API Reference (Tempera- 2127
ture Parameter). [https://platform.openai.com/docs/api-](https://platform.openai.com/docs/api-reference/completions/create#completions-create-temperature) 2128
[reference/completions/create#completions-create-temperature](https://platform.openai.com/docs/api-reference/completions/create#completions-create-temperature). 2129
2130
2131
2132

[69] Johnson D, Goodman R, Patrinely J, Stone C, Zimmerman E, Donald R, et al. 2133
Assessing the Accuracy and Reliability of AI-Generated Medical Responses: An 2134
Evaluation of the Chat-GPT Model. Research Square. 2023 Feb;p. rs.3.rs– 2135
2566942. <https://doi.org/10.21203/rs.3.rs-2566942/v1>. 2136
2137
2138
2139

[70] Jang ME, Lukasiewicz T.: Consistency Analysis of ChatGPT. arXiv. 2140
2141
2142

[71] Krishna S, Bhambra N, Bleakney R, Bhayana R, Atzen S. Evaluation of Reli- 2143
ability, Repeatability, Robustness, and Confidence of GPT-3.5 and GPT-4 on 2144
a Radiology Board–Style Examination. Radiology. 2024 May;311(2):e232715. 2145
<https://doi.org/10.1148/radiol.232715>. 2146
2147
2148
2149

[72] Setel PW, Macfarlane SB, Szreter S, Mikkelsen L, Jha P, Stout S, et al. A 2150
Scandal of Invisibility: Making Everyone Count by Counting Everyone. The 2151
Lancet. 2007;370(9598):1569–1577. 2152
2153
2154
2155

[73] Fottrell E, Byass P. Verbal Autopsy: Methods in Transition. Epidemiologic 2156
reviews. 2010;32(1):38–55. 2157
2158
2159
2160
2161
2162

2163 [74] Byass P. The Imperfect World of Global Health Estimates. PLoS Medicine. 2010
2164 Nov;7(11):e1001006. <https://doi.org/10.1371/journal.pmed.1001006>.
2165
2166

2167 [75] Murray CJ, Lopez AD. Mortality by Cause for Eight Regions of the World:
2168 Global Burden of Disease Study. The lancet. 1997;349(9061):1269–1276.
2169
2170

2171 [76] Mathers CD, Boerma T, Ma Fat D. Global and Regional Causes of Death. British
2172 medical bulletin. 2009;92(1):7–32.
2173
2174

2175 [77] Vos T, Lim SS, Abbafati C, Abbas KM, Abbasi M, Abbasifard M, et al. Global
2176 Burden of 369 Diseases and Injuries in 204 Countries and Territories, 1990–2019:
2177 A Systematic Analysis for the Global Burden of Disease Study 2019. The lancet.
2178 2020;396(10258):1204–1222.
2179
2180
2181
2182

2183 [78] Shawon MTH, Ashrafi SAA, Azad AK, Firth SM, Chowdhury H, Mswia RG,
2184 et al. Routine Mortality Surveillance to Identify the Cause of Death Pattern
2185 for Out-of-Hospital Adult (Aged 12+ Years) Deaths in Bangladesh: Introduction
2186 of Automated Verbal Autopsy. BMC Public Health. 2021 Dec;21(1):491. <https://doi.org/10.1186/s12889-021-10468-7>.
2187
2188
2189
2190
2191

2192 [79] Maqungo M, Nannan N, Nojilana B, Nichols E, Morof D, Cheyip M, et al. Can
2193 Verbal Autopsies Be Used on a National Scale? Key Findings and Lessons from
2194 South Africa’s National Cause-of-Death Validation Study. Global Health Action.
2195 2024 Dec;17(1):2399413. <https://doi.org/10.1080/16549716.2024.2399413>.
2196
2197
2198
2199

2200 [80] Onyango D, Awuonda B. Using Verbal Autopsy to Enhance Mortality Surveil-
2201 lance. The Lancet Global Health. 2024;12(8):e1217–e1218.
2202
2203

2204 [81] Wang J, Shi E, Yu S, Wu Z, Ma C, Dai H, et al.: Prompt Engineering for
2205 Healthcare: Methodologies and Applications. arXiv.
2206
2207
2208

[82] Lewis P, Perez E, Piktus A, Petroni F, Karpukhin V, Goyal N, et al. Retrieval-Augmented Generation for Knowledge-Intensive NLP Tasks. In: Proceedings of the 34th International Conference on Neural Information Processing Systems. NIPS '20. Red Hook, NY, USA: Curran Associates Inc.; 2020. p. 9459–9474.	2209 2210 2211 2212 2213 2214 2215
[83] Meskó B. Prompt Engineering as an Important Emerging Skill for Medical Professionals: Tutorial. Journal of medical Internet research. 2023;25:e50638.	2216 2217 2218 2219 2220
[84] Almasre M. Development and Evaluation of a Custom GPT for the Assessment of Students' Designs in a Typography Course. Education Sciences. 2024 Feb;14(2):148. https://doi.org/10.3390/educsci14020148 .	2221 2222 2223 2224 2225
[85] Loh P, Fottrell E, Beard J, Bar-Zeev N, Phiri T, Banda M, et al. Added Value of an Open Narrative in Verbal Autopsies: A Mixed-Methods Evaluation from Malawi. BMJ Paediatrics Open. 2021 Feb;5(1):e000961. https://doi.org/10.1136/bmjpo-2020-000961 .	2226 2227 2228 2229 2230 2231 2232 2233
[86] World Health Organization.: ODK for Verbal Autopsy: A Quick Guide. https://www.who.int/publications/m/item/odk-for-verbal-autopsy-a-quick-guide .	2234 2235 2236 2237 2238 2239
[87] Nafundi.: ODK - Collect Data Anywhere.	2240 2241
[88] DiPasquale A, Maire N, Bratschi M.: Release ODK 2016 WHO VA Instrument 1.5.1 SwissTPH/WHO-VA. Swiss Tropical and Public Health Institute.	2242 2243 2244 2245
[89] Byass P.: InterVA-5.1 User Guide.	2246 2247 2248
[90] Thomas J, ekarpinskiMITRE, pkmitre, owentrigueros, Choi P, Chu Y.: Pycrossva: Prepare Data from WHO and PHRMC Instruments for Verbal Autopsy Algorithms.	2249 2250 2251 2252 2253 2254

2255 [91] Li ZR, McCormick T, Clark S.: InSilicoVA: Probabilistic Verbal Autopsy Coding
2256 with 'InSilicoVA' Algorithm.
2257
2258
2259 [92] Thomas J, Li Z, Byass P, McCormick T, Boyas M, Clark S.: InterVA5: Replicate
2260 and Analyse 'InterVA5'.
2261
2262
2263 [93] Yendewa GA, Poveda E, Yendewa SA, Sahr F, Quiñones-Mateu ME, Salata RA.
2264 HIV/AIDS in Sierra Leone: Characterizing the Hidden Epidemic. AIDS reviews.
2265 2018;20(2).
2266
2267
2268
2269 [94] Walker PG, White MT, Griffin JT, Reynolds A, Ferguson NM, Ghani AC. Malaria
2270 Morbidity and Mortality in Ebola-affected Countries Caused by Decreased
2271 Health-Care Capacity, and the Potential Effect of Mitigation Strategies: A
2272 Modelling Analysis. The Lancet Infectious Diseases. 2015;15(7):825–832.
2273
2274
2275
2276
2277
2278
2279
2280
2281
2282
2283
2284
2285
2286
2287
2288
2289
2290
2291
2292
2293
2294
2295
2296
2297
2298
2299
2300