

Book II of Euclid's Elements

Formal Reconstruction — Technical Documentation

Jurek Różański

Military University of Technology, Warsaw, Poland

CGJTEAMLAB

August 29, 2026

Document Status

This volume is the technical documentation on formal reconstruction of Book II of Euclid's *Elements*. It records the mathematical architecture of the reconstruction, the Lean formalization, the dependency structure of individual propositions, the reusable rectangle and scissors-congruence infrastructure, and the relation between the formal development and the classical sources.

The documentation accompanies the formal development. The Lean source is authoritative for theorem statements, hypotheses, dependencies, and proof status.

Contents

Document Status	ii
List of Figures	ix
1 Proposition II.1	1
1.1 Introduction	1
1.2 Euclid’s “Rectangle Contained By”	2
1.2.1 The original geometric meaning	2
1.2.2 Why this already looks algebraic	2
1.3 The Synthetic Rectangle Layer	2
1.3.1 Rectangles	2
1.3.2 The formal meaning of “contained by”	3
1.3.3 No segment multiplication and no numerical area	3
1.4 Exact Scissors Equality and Equal Content	4
1.4.1 Exact scissors equality	4
1.4.2 The more general equal-content relation	4
1.5 The Classical and Formal Configurations	4
1.5.1 The binary split	4
1.5.2 The three-part split	5
1.5.3 Scope of the present formal statement	6
1.6 From a Concrete Diagram to Euclid’s Language	6
1.7 Representative Independence	6
1.7.1 Why representatives matter formally	6
1.8 Commutativity by Rectangle Rotation	7
1.9 Proof Architecture of the Reconstruction	8
1.9.1 Step 1: exact binary dissection	8
1.9.2 Step 2: iterate the split	8
1.9.3 Step 3: attach the “contained by” data	9
1.9.4 Step 4: remove the choice of representatives	9
1.9.5 Step 5: expose the symmetric side order	9
1.10 Classical Proof and Formal Proof	9
1.10.1 Euclid’s proof pattern	9
1.10.2 The formal DAG	10
1.11 Geometry, Content, and Representation	10
1.11.1 Geometry	10
1.11.2 Content	10
1.11.3 Representation	10
1.12 Hidden Configuration Data	11
1.13 Axiomatic Status	11
1.14 Relationship with Book I and the Following Propositions	12
1.15 Source Audit	12
1.15.1 Euclid	12

1.15.2	Hartshorne	12
1.15.3	Hilbert	12
1.15.4	The phrase “geometric algebra”	13
1.16	Dependency Summary	13
1.17	Conclusion	14
2	Proposition II.2	16
2.1	Introduction	16
2.2	Euclid’s Statement and Classical Configuration	16
2.2.1	The statement	16
2.2.2	The classical diagram	17
2.3	Classical Proof	17
2.4	The Geometric Identity	18
2.5	Reusable Book II Infrastructure	18
2.5.1	A square as a rectangle contained by the same side twice	18
2.5.2	The binary form of II.1	19
2.5.3	Exchanging the containing-side order	19
2.6	Proof Architecture of the Reconstruction	20
2.6.1	Step 1: recognize the square	20
2.6.2	Step 2: rotate the two arbitrary part representatives	20
2.6.3	Step 3: apply II.1 with fixed segment BC	20
2.6.4	Step 4: return to Euclid’s ordering	21
2.7	Lean Formalization	21
2.8	Relationship with Proposition II.1	22
2.9	Relationship with Book I Infrastructure	23
2.10	Formalization Notes and Hidden Geometric Data	23
2.11	Geometry, Content, Representation, and Later Arithmetic	24
2.11.1	Geometry	24
2.11.2	Content / scissors	24
2.11.3	Representation	24
2.11.4	Later arithmetic / numerical area	24
2.12	Axiomatic and Semantic Status	24
2.13	Hartshorne’s Semantic Control	25
2.14	Hilbert’s Layering	25
2.15	Dependency Summary	25
2.16	Conclusion	26
3	Proposition II.3	27
3.1	Introduction	27
3.2	Euclid’s Statement and Classical Configuration	27
3.2.1	The statement	27
3.2.2	The classical diagram	28
3.3	Classical Proof	28
3.4	The Geometric Identity	30
3.5	Reusable Book II Infrastructure	30
3.5.1	The binary form of II.1	30
3.5.2	A square as $\text{Rect}(MC, MC)$	31
3.5.3	What II.3 does not use	31
3.6	Proof Architecture of the Reconstruction	31
3.6.1	Step 1: recognize the square	31
3.6.2	Step 2: instantiate II.1 with the fixed segment MC	32
3.7	Lean Formalization	32

3.8	Relationship with Earlier Book II Propositions	34
3.8.1	Proposition II.1	34
3.8.2	Proposition II.2	34
3.9	Relationship with Book I Infrastructure	34
3.10	Formalization Notes and Hidden Geometric Data	35
3.11	Geometry, Content, Representation, and Later Arithmetic	35
3.11.1	Geometry	35
3.11.2	Content / scissors	35
3.11.3	Representation	35
3.11.4	Later arithmetic / numerical area	36
3.12	Axiomatic and Semantic Status	36
3.13	Hartshorne's Semantic Control	36
3.14	Hilbert's Layering	37
3.15	Dependency Summary	37
3.16	Conclusion	38
4	Proposition II.4	39
4.1	Introduction	39
4.2	Euclid's Statement and Classical Configuration	40
4.2.1	The statement	40
4.2.2	The classical diagram	40
4.3	Classical Proof	40
4.4	The Geometric Identity in the Current Formalization	42
4.5	Reusable Book II Infrastructure	42
4.5.1	Proposition II.2: split the square on the whole	42
4.5.2	Proposition II.3: expand a whole-by-part rectangle	43
4.5.3	Rectangle rotation and endpoint reversal	43
4.5.4	Exact scissors addition	44
4.6	Proof Architecture of the Reconstruction	44
4.6.1	Step 1: reverse the cut orientation for the left branch	44
4.6.2	Step 2: apply II.2 to the square on the whole	44
4.6.3	Step 3: prepare the reversed left II.3 call	44
4.6.4	Step 4: expand the left rectangle by II.3	45
4.6.5	Step 5: expand the right rectangle by II.3	46
4.6.6	Step 6: add the branches and attach them to the square	46
4.7	Lean Formalization	48
4.8	Relationship with Earlier Book II Propositions	51
4.8.1	Proposition II.1	51
4.8.2	Proposition II.2	51
4.8.3	Proposition II.3	51
4.9	Relationship with Book I Infrastructure	51
4.10	Formalization Notes and Hidden Geometric Data	52
4.11	Geometry, Content, Representation, and Later Arithmetic	52
4.11.1	Geometry	52
4.11.2	Content / scissors	52
4.11.3	Representation	53
4.11.4	Later arithmetic / numerical area	53
4.12	Axiomatic and Semantic Status	53
4.13	Hartshorne's Semantic Control	54
4.14	Hilbert's Layering	54
4.15	Dependency Summary	55
4.16	Conclusion	55

5	Proposition II.5	57
5.1	Introduction	57
5.2	Euclid's Statement and Classical Configuration	57
5.2.1	The statement	57
5.2.2	The classical diagram	58
5.3	Classical Proof	58
5.4	The Geometric Identity in the Current Formalization	59
5.5	Reusable Book II Infrastructure	59
5.5.1	The binary rectangle split from II.1	59
5.5.2	Proposition II.3	60
5.5.3	Rectangle representation and transport	60
5.6	Proof Architecture of the Reconstruction	60
5.6.1	Step 1: recover the full order and midpoint congruence	60
5.6.2	Step 2: apply II.3 on the left half	60
5.6.3	Step 3: transport the midpoint side	61
5.6.4	Step 4: split the unequal part DB	61
5.6.5	Step 5: the scissors bridge	61
5.6.6	Step 6: assemble the right side	61
5.6.7	Step 7: recognize the half-square	61
5.7	Lean Formalization	62
5.7.1	The public theorem signature	62
5.7.2	The final three-stage assembly	64
5.8	Relationship with Earlier Book II Propositions	66
5.8.1	Proposition II.1	66
5.8.2	Proposition II.3	66
5.8.3	Proposition II.4	66
5.9	Relationship with Book I Infrastructure	66
5.10	Formalization Notes and Hidden Geometric Data	67
5.10.1	Order orientation	67
5.10.2	Midpoint as geometry, not arithmetic	67
5.10.3	Representative independence	67
5.10.4	No gnomon primitive	67
5.10.5	No cancellation axiom	67
5.11	Axiomatic and Semantic Status	67
5.12	Dependency Summary	68
5.13	Conclusion	69
6	Proposition II.6	70
6.1	Introduction	70
6.2	Euclid's Statement and Classical Configuration	70
6.2.1	The statement	70
6.2.2	The classical diagram	71
6.3	Classical Proof	72
6.4	The Geometric Identity in the Current Formalization	72
6.5	Reusable Book II Infrastructure Used	73
6.5.1	Proposition II.3: extend the unequal-parts rectangle	73
6.5.2	Proposition II.1: split the old segment at its midpoint	73
6.5.3	Midpoint transport	73
6.5.4	Proposition II.4 on the produced segment	74
6.6	Proof Architecture of the Reconstruction	74
6.6.1	Step 1: recover the full order	74
6.6.2	Step 2: decompose the rectangle on AD, BD	74

6.6.3	Step 3: split AB at C	74
6.6.4	Step 4: compose the two decompositions	74
6.6.5	Step 5: use the midpoint congruence	75
6.6.6	Step 6: normalize the left decomposition	75
6.6.7	Step 7: expand the square on CD by II.4	75
6.6.8	Step 8: the abstract scissors bridge	75
6.6.9	Step 9: assemble II.6	75
6.7	Lean Formalization	76
6.7.1	The public theorem signature	76
6.7.2	The final bridge call	79
6.8	Relationship with Earlier Book II Propositions	79
6.8.1	Proposition II.1	79
6.8.2	Proposition II.3	79
6.8.3	Proposition II.4	79
6.8.4	Proposition II.5	80
6.9	Relationship with Book I Infrastructure	80
6.10	Formalization Notes and Hidden Geometric Data	80
6.10.1	The extension order is proved	80
6.10.2	Midpoint data have two independent roles	80
6.10.3	Representative reuse is deliberate	80
6.10.4	The square terms remain geometric representatives	81
6.10.5	No gnomon primitive	81
6.10.6	No cancellation or subtraction of content	81
6.11	Axiomatic and Semantic Status	81
6.12	Dependency Summary	82
6.13	Conclusion	82
A	Greenberg: Foundational and Semantic Control	84
.1	Euclid I.1–I.20 in Greenberg’s Hilbert-Style Development	84
.1.1	Purpose of the comparison	84
.1.2	Comparison table	84
.1.3	Result of the query	86
.1.4	Consequences for the formal reconstruction	86
.1.5	Current checkpoint	86
A	Hartshorne: Equal Content and the Algebra of Areas	87
A.1	Purpose and scope	87
A.2	Hartshorne’s organization of Book II	87
A.3	Comparison table	88
A.4	Semantic observations retained from Hartshorne	89
A.5	II.1–II.6: from distribution to the two midpoint identities	89
A.6	Result of the Hartshorne audit	90
A.7	Current checkpoint	90
B	Hilbert: Content and Segment Arithmetic	91
B.1	Purpose and scope	91
B.2	Hilbert’s organization of the relevant theory	91
B.3	Comparison table	91
B.4	Structural observations retained from Hilbert	93
B.5	II.1–II.6: exact dissection below segment arithmetic	93
B.6	Result of the Grundlagen audit	94
B.7	Current checkpoint	94

C	Borsuk–Szmielw: Axiomatic Layers and Representation	95
C.1	Why this source matters	95
C.2	Layer notation	96
C.3	Euclid in the Borsuk–Szmielw development	96
C.4	The strongest architectural correspondences	101
C.4.1	Transport is primitive-strength infrastructure	101
C.4.2	Pairs of Euclidean propositions become biconditionals	101
C.4.3	Midpoints and bisectors include positional information	101
C.5	The neutral parallel theory is already large	102
C.6	Theorem 12 of Chapter V as an interface theorem	102
C.7	What the book contributes at I.35	103
C.8	Consequences for the formal library	103
C.8.1	Hilbert interface candidates	104
C.8.2	Book Zero candidates	104
C.8.3	Euclidean interface candidates	105
C.9	Conclusion	105
	Bibliography	106

List of Figures

1.1	Binary and three-part rectangle dissections	5
1.2	Commutativity from cyclic rectangle rotation	8
2.1	Square dissection and rectangle rotation	17
3.1	Binary rectangle split with the square on MC	29
4.1	Square decomposition into two squares and two equal rectangles	41
5.1	The gnomon mechanism in II.5	58
6.1	The gnomon mechanism in II.6	71

Chapter 1

Proposition II.1

1.1 Introduction

Proposition II.1 opens Book II with a change of mathematical language. Book I has developed a synthetic theory of congruence, parallels, parallelograms, squares, and equal content. Book II begins to combine these ingredients in a systematic calculus of rectangles. A modern reader immediately recognizes in II.1 the pattern of a distributive law. That recognition is useful, but it must be stated carefully: Euclid does not introduce variables for numerical lengths, does not define multiplication of lengths, and does not prove an identity in a number field.

The primitive object in the proposition is a geometric figure: a rectangle “contained by” two straight lines. When one of the two lines is divided into parts, the large rectangle is divided into smaller rectangles having the same other side. The equality asserted by the proposition is therefore first of all a statement about decomposition of figures.

This is a natural point at which to speak of the beginning of the *geometrical algebra* characteristic of Book II. The phrase is an interpretive description of a stable pattern of geometric operations, not a claim that Euclid was secretly manipulating modern symbolic formulas. In the present formal reconstruction this distinction is especially sharp: the code contains rectangles, segment congruence, betweenness, parallelograms, right angles, and scissors relations, but no multiplication operation on segment lengths and no numerical area function.

The formal development separates three levels:

1. a concrete rectangle and its geometric cuts;
2. the Euclidean phrase “rectangle contained by” two given segments;
3. independence of the particular rectangle chosen to represent that phrase.

The final theorem is stronger than a bare equal-content statement. Its conclusion is the exact relation

```
HilbertScissorsEq
```

for the whole rectangle and the formal sum of its parts.

1.2 Euclid’s “Rectangle Contained By”

1.2.1 The original geometric meaning

Book II begins with the terminology that a rectangular parallelogram is “contained by” the two straight lines which contain one of its right angles. Thus the expression does not initially denote a product of two numbers. It names a rectangle by two adjacent sides.

Suppose a rectangle has adjacent sides congruent to two given segments PQ and RS . Euclid’s phrase

the rectangle contained by PQ, RS

should be read geometrically: choose a right-angled parallelogram whose two adjacent sides have those magnitudes. The rectangle itself remains a plane figure.

For exposition only, we will occasionally write

$$\text{Rect}(PQ, RS).$$

This is documentary shorthand. It is not a Lean definition, it is not a number-valued function, and the comma is not multiplication.

1.2.2 Why this already looks algebraic

If BC is divided into consecutive parts BM , MN , and NC , II.1 says in this shorthand

$$\text{Rect}(BC, PQ) = \text{Rect}(BM, PQ) + \text{Rect}(MN, PQ) + \text{Rect}(NC, PQ).$$

A modern algebraist naturally sees the shape

$$(b_1 + b_2 + b_3)a = b_1a + b_2a + b_3a.$$

The formal proof, however, establishes the geometric statement on the first line, not the numerical identity on the second. The second formula is only a modern mnemonic for the combinatorial pattern of the dissection.

This is consistent with the modern synthetic reading of Euclid’s area theory. Hartshorne treats Book II as a collection of equal-content statements about figures and describes the resulting manipulation as “geometrical algebra.” The point is precisely that the manipulation precedes any assignment of numerical areas to the figures.

1.3 The Synthetic Rectangle Layer

1.3.1 Rectangles

The reconstruction introduces a small reusable rectangle API in `HilbertRectangle.lean`. A rectangle is a parallelogram with one specified right angle:

```
def IsRectangle
  (A B C D : Geo.Point) : Prop :=
  IsParallelogram Geo A B C D /\
  HilbertRightAngle Geo A B C
```

The parallelogram structure supplies the opposite-side relations, while the right-angle component distinguishes rectangles from arbitrary parallelograms. The remaining right angles can then be propagated synthetically.

1.3.2 The formal meaning of “contained by”

The Euclidean phrase is represented by the predicate

```
def IsRectangleContainedBy
  (A B C D P Q R S : Geo.Point) : Prop :=
  IsRectangle Geo A B C D /\
  Geo.Congruent A B P Q /\
  Geo.Congruent B C R S
```

Thus

```
IsRectangleContainedBy Geo A B C D P Q R S
```

means exactly that $ABCD$ is a concrete rectangle, $AB \cong PQ$, and $BC \cong RS$. Nothing in this definition assigns a number to PQ or RS . The two given segments specify the two adjacent geometric magnitudes of a representative rectangle.

The layer also proves existence of such a representative and, more importantly for the semantics of Book II, uniqueness of its scissors content. The theorem

```
rectangle_contained_by_unique
```

says that any two rectangles contained by the same ordered pair of segments satisfy the exact scissors relation. Consequently Euclid’s definite description “the rectangle contained by PQ, RS ” becomes well-defined at the scissors level even though many different concrete quadrilaterals can represent it.

1.3.3 No segment multiplication and no numerical area

There are two tempting modernizations that the present reconstruction explicitly avoids.

First, we do not define a product of segment lengths $|PQ| |RS|$. The pair (PQ, RS) remains a pair of geometric segments, and the rectangle is related to that pair by congruence of adjacent sides.

Second, we do not map a rectangle to a real number called its area. The content of the proposition is expressed through relations between finite formal sums of triangles. This permits addition, dissection, transport, and comparison of figures at the synthetic level.

This is also conceptually distinct from Hilbert’s later arithmetic of segments, where arithmetic operations are introduced as an additional construction. No such field or segment-arithmetic layer is imported into II.1.

1.4 Exact Scissors Equality and Equal Content

The project uses two related but different content relations. They should not be conflated.

1.4.1 Exact scissors equality

`HilbertScissorsEq` records an exact finite cut-and-paste relation between formal scissors terms. In II.1 it is produced by explicit splits of the rectangle into smaller rectangles and by normalization of the triangulated parallelogram terms. Schematically,

$$P \simeq_{\text{sc}} Q + R$$

means that the two sides can be matched by the exact scissors machinery already developed in the project.

The rectangle transport theorem is already available in this stronger form:

```
rectangle_transport_scissorsEq
```

for rectangles with congruent corresponding adjacent sides.

1.4.2 The more general equal-content relation

The broader project notion corresponding to equal content is

```
HilbertScissorsEquicomplementable
```

It allows suitable complementary terms to be adjoined before exact scissors equality is established. Thus it is designed to capture a more general synthetic equal-content relation.

The theorem

```
rectangle_transport
```

is obtained by weakening the exact transport theorem from `HilbertScissorsEq` to equicomplementability.

For II.1 no such weakening is necessary. The final theorem proves `HilbertScissorsEq` itself. Therefore II.1 certainly yields an equal-content consequence, but its formal statement retains the stronger fact that the displayed equality comes from an exact dissection.

1.5 The Classical and Formal Configurations

1.5.1 The binary split

The reusable core is a one-cut theorem. Let $B - M - C$ and let the corresponding point L divide the opposite side DE . The rectangle $BCED$ is cut into the rectangles $BMLD$ and $MCEL$:

$$BCED \simeq_{\text{sc}} BMLD + MCEL.$$

In Lean the theorem is

```
rectangle_split
```

and it returns both the rectangle structure of the two pieces and the exact scissors equality.

The formal configuration contains one additional point X . It satisfies $D - X - C$ and $M - X - L$. This is not decorative data: the inherited exact split lemma from the I.47 scissors infrastructure uses the intersection of the cut line with a diagonal. The classical rectangle picture suppresses this witness; the formal theorem makes it explicit.

1.5.2 The three-part split

The three-part form applies the binary split twice. First cut at $M - L$; then cut the remaining rectangle at $N - K$. The result is

$$\begin{aligned} BCED &\simeq_{\text{sc}} BMLD \\ &\quad + MNKL \\ &\quad + NCEK. \end{aligned}$$

The second split has its own diagonal-intersection witness Y , with $L - Y - C$ and $N - Y - K$.

Figure 1.1 shows both levels and includes the hidden witnesses X and Y required by the current Lean interface.




Figure 1.1: The exact rectangle splits underlying II.1. Panel (a) is the binary kernel: $B - M - C$ and the cut ML divide $BCED$ into $BMLD$ and $MCEL$; the dashed diagonal DC meets the cut at the formal witness X . Panel (b) applies the same construction twice: BC is divided into BM , MN , and NC , producing the three rectangles $BMLD$, $MNKL$, and $NCEK$. The dashed diagonals display the witnesses X and Y used by the two exact split calls.

The actual three-part theorem is

```
rectangle_split_three
```

Its proof is structurally transparent: obtain the first binary equality, split the remaining right-hand rectangle, add the unchanged left piece to that second equality, and compose by transitivity of `HilbertScissorsEq`.

1.5.3 Scope of the present formal statement

Euclid states II.1 for one line cut into an arbitrary finite number of parts. The current Lean file does not yet package an n -ary list-valued theorem. It proves the binary split and the explicit three-part form used in the standard II.1 configuration. The binary theorem is the reusable kernel from which a general finite version could later be obtained by iteration.

This is a deliberate statement-fidelity point. The documentation records Euclid’s general formulation, but the final formal statement displayed below is the representative-independent three-part theorem actually present in `Proposition2_1.lean`.

1.6 From a Concrete Diagram to Euclid’s Language

The geometric split alone produces concrete rectangles. Euclid II.1 is more abstract: it speaks of the rectangles *contained by* the whole segment and the several parts together with one fixed segment.

The theorem

```
rectangle_split_three_contained_by
```

bridges these levels. If $CE \cong PQ$, it packages the four concrete rectangles as

$$\begin{aligned} BCED & : \text{Rect}(BC, PQ), \\ BMLD & : \text{Rect}(BM, PQ), \\ MNKL & : \text{Rect}(MN, PQ), \\ NCEK & : \text{Rect}(NC, PQ). \end{aligned}$$

The opposite-side congruences of the parallelograms are what transport the fixed side PQ through the two cuts. The conclusion still contains the same exact scissors equality between the concrete quadrilateral terms.

At this stage the geometry corresponding to Euclid’s wording is visible, but the theorem still uses one particular drawn rectangle and its particular cut points. The final step removes this representational dependence.

1.7 Representative Independence

1.7.1 Why representatives matter formally

A phrase such as

$$\text{Rect}(BC, PQ)$$

does not identify four vertices. Lean, however, must manipulate an actual term such as

```
hilbertParallelogramTerm Geo U0 U1 U2 U3
```

with named vertices. The final theorem therefore accepts arbitrary representatives for the whole rectangle and for each of the three part rectangles.

The concrete dissection first proves

$$BCED \simeq_{\text{sc}} BMLD + MNKL + NCEK.$$

Then four calls to

```
rectangle_contained_by_unique
```

transport the whole and the three pieces to arbitrary representatives U , V , W , and Z . Compatibility of `HilbertScissorsEq` with formal addition transports the nested sum. Transitivity completes the chain

arbitrary whole representative
 $\Downarrow$
 concrete whole rectangle
 $\Downarrow$
 concrete three-part dissection
 $\Downarrow$
 arbitrary representatives of the three parts.

This is the formal content of representative independence. It is not merely a convenience for Lean. It explains why the Euclidean expression “the rectangle contained by two segments” can function as a stable geometric magnitude without first quotienting segments into numerical lengths or assigning numerical areas to rectangles.

1.8 Commutativity by Rectangle Rotation

The second algebra-like feature is symmetry in the two containing segments. For a numerical product one would write $ab = ba$. The synthetic proof does not appeal to a commutative multiplication law. Instead it rotates the presentation of a rectangle.

If $ABCD$ is contained by PQ, RS , then the cyclically reoriented rectangle $BCDA$ has adjacent sides BC and CD . The first is already congruent to RS ; the second is congruent to AB , hence to PQ . Thus the same geometric rectangle, with its vertices read from the next corner, is contained by RS, PQ .

The exact theorem available in the rectangle layer is

```
rectangle_contained_by_swap
```

It returns both the swapped `IsRectangleContainedBy` fact and a `HilbertScissorsEq` identifying the two quadrilateral presentations. The current rectangle API also packages the representative-independent consequence as

```
rectangle_contained_by_comm
```

so the commutativity needed by later Book II propositions is a theorem of rectangle geometry rather than an assumed law of segment multiplication.

Figure 1.2 shows the geometric mechanism. The operation is a cyclic change of the distinguished corner of the same rectangle; this exchanges the two adjacent side roles.




Figure 1.2: The geometric source of commutativity. On the left, $ABCD$ represents the rectangle contained by PQ, RS . Reading the same rectangle from the next vertex gives $BCDA$: now the first adjacent side represents RS and the second represents PQ . Exact scissors invariance under this cyclic presentation change yields the synthetic counterpart of exchanging the two factors.

This lemma is already operational in the next proposition. II.2 naturally writes rectangles in the order “whole side, part”, whereas the II.1 cut lemma produces them in the order “part, fixed side”. The implementation of II.2 uses the swap theorem to pass between these two presentations before applying the binary form of II.1.

1.9 Proof Architecture of the Reconstruction

The Lean proof is best read as five mathematical stages.

1.9.1 Step 1: exact binary dissection

```
rectangle_split_scissorsEq
rectangle_split_left
rectangle_split_right
rectangle_split
```

The inherited I.47 split theorem supplies the exact scissors decomposition. The two local rectangle helpers prove that each piece remains a rectangle by propagating right angles through the relevant parallelograms and along the same rays determined by betweenness.

1.9.2 Step 2: iterate the split

```
rectangle_split_three
```

The right-hand remainder of the first split is itself a rectangle, so the same binary theorem can be applied again. Addition and transitivity of exact scissors equality compose the two decompositions.

1.9.3 Step 3: attach the “contained by” data

```
rectangle_split_three_contained_by
```

Opposite-side congruence transports the fixed segment PQ through all three pieces. Each concrete piece is thereby promoted from `IsRectangle` to `IsRectangleContainedBy`.

1.9.4 Step 4: remove the choice of representatives

```
rectangle_contained_by_unique
euclid_proposition_2_1_three
```

The uniqueness theorem replaces the concrete whole and each concrete part by an arbitrary rectangle contained by the same pair of segments. This is the semantic step that makes the final statement faithful to Euclid’s definite phrase “the rectangle contained by.”

1.9.5 Step 5: expose the symmetric side order

```
rectangle_contained_by_swap
rectangle_contained_by_comm
```

Cyclic rotation of the rectangle exchanges the two containing segments. This is not needed to establish the displayed three-part split itself, but it is an important part of the Book II rectangle interface and is immediately useful in II.2 and later propositions.

1.10 Classical Proof and Formal Proof

1.10.1 Euclid’s proof pattern

In the classical three-part configuration, take one undivided segment A and a segment BC divided at D and E . Erect a perpendicular at one endpoint of BC , lay off on it a segment equal to A , complete the outer rectangle, and through the division points draw parallels to the perpendicular side. The outer rectangle is visibly partitioned into three rectangles. Each smaller rectangle is contained by A and one of the three consecutive parts of BC . Common Notion 2 then combines the parts into the whole equality.


The classical DAG is therefore extremely small:

```

      divide one side
        ↓↓
    draw the corresponding parallels
        ↓↓
    obtain three subrectangles
        ↓↓
whole rectangle = sum of the three subrectangles.
```

1.10.2 The formal DAG

The current reconstruction expands two things that the classical diagram treats as immediate: exact scissors bookkeeping and independence of the chosen rectangle representatives.



The proof is therefore not a mechanical transcription of Euclid. It preserves the geometric core, but makes the semantic and representational layers explicit.

1.11 Geometry, Content, and Representation

1.11.1 Geometry

The geometric data are substantial even though the proposition is visually simple. The current theorem receives the outer rectangle, strict betweenness relations on the divided side and its opposite, two diagonal-intersection witnesses, and four parallelogram facts for the two successive cuts. Right angles of the two resulting rectangles are not read from the picture; they are proved by right-angle transport and parallelogram geometry.

No coordinates are introduced. The cut points are controlled by **Between**, the side lengths by segment congruence, and the rectangular structure by parallels and synthetic right angles.

1.11.2 Content

Content enters in its strongest local form. The core statements are exact **HilbertScissorsEq** decompositions. The three-part theorem is composed from two exact binary splits by addition and transitivity. Representative transport is also exact scissors equality.

The more general equicomplementability relation is part of the surrounding Book I content theory, but II.1 does not need to weaken its conclusion to that level.

1.11.3 Representation

Representation is central to the formal meaning of the proposition. A rectangle contained by PQ, RS is not one canonical four-tuple of vertices. The predicate **IsRectangleContainedBy** describes a representative; **rectangle_contained_by_unique** proves that different representatives have the same exact scissors content.

There is a second representation issue inside the split itself. The inherited scissors theorem and the Book II statement use different cyclic orders for some parallelogram terms. The proof normalizes these by

```
parallelogram_term_rotateOne
```

before composing the scissors equalities. These cyclic changes are bookkeeping of a geometric representation, not new area facts.

1.12 Hidden Configuration Data

The classical drawing makes several facts appear automatic. Lean records them explicitly:

- $B - M - C$ for the first cut and $M - N - C$ for the second;
- $D - L - E$ and $L - K - E$ on the opposite side;
- $D - X - C$ together with $M - X - L$ for the first diagonal/cut intersection;
- $L - Y - C$ together with $N - Y - K$ for the second diagonal/cut intersection;
- explicit parallelogram witnesses for the left and right pieces of each split;
- congruence $CE \cong PQ$, which identifies the fixed side used in the “contained by” descriptions;
- four arbitrary rectangle representatives in the final theorem.

The points X and Y do not appear in Euclid’s statement and need not be emphasized in an elementary picture of II.1. They are nevertheless real data of the current exact-dissection API, inherited from the triangulated proof of the rectangle split.

1.13 Axiomatic Status

The public II.1 theorems assume

```
[HilbertIncidence Geo]
[HilbertEuclideanPlane Geo]
```

so II.1 is Euclidean in the current project API. Group IV is available through this typeclass and is consumed transitively by the parallelogram and parallel infrastructure on which the rectangle layer and inherited split theorem depend. The proposition itself does not introduce a new parallel axiom.

No continuity typeclass, Archimedean axiom, numerical area function, or segment field is introduced in `Proposition2_1.lean`. No proposition-local `axiom` or `sorry` occurs in the audited source. The modular builds for `HilbertRectangle` and `Proposition2_1` are recorded as clean at the present project checkpoint.

The content status is also worth separating from the later converse arguments of Book I. II.1 proceeds from geometry to an exact dissection. It does not infer metric geometry from an equal-content hypothesis, so it does not need a proper-part or De Zolt-type strictness argument.

1.14 Relationship with Book I and the Following Propositions

The exact split theorem reuses the scissors infrastructure established during I.47. In particular, the underlying theorem historically named

```
i47_square_split
```

actually requires only parallelogram structure, and II.1 repackages it as a general rectangle split.

The rectangle layer also uses Book I results about parallelograms, opposite sides, congruent right angles, SAS/SSS transport, and the scissors representation of parallelograms. Thus Book II does not start a new foundational geometry; it starts a new way of organizing geometric magnitudes on top of the completed Book I infrastructure.

II.2 immediately imports II.1. Its formal proof illustrates why the symmetric rectangle API matters: Euclid’s order of the two containing sides is opposite to the order produced by the binary cut theorem, so the representatives are swapped by rectangle rotation before II.1 is applied. II.3 continues the same rectangle calculus. In this sense II.1 is not only the first theorem of Book II but the API-level distributive kernel for the propositions that follow.

1.15 Source Audit

1.15.1 Euclid

Book II Definition 1 supplies the semantic key: the two lines “containing” a rectangle are the adjacent sides that contain a right angle. Proposition II.1 then divides one of two given lines into several parts and identifies the whole rectangle with the sum of the rectangles formed from the undivided line and those parts.

The proof is geometric throughout. The division points on one side are carried across the outer rectangle by parallels. The resulting subrectangles literally partition the original rectangle. Nothing in the proposition requires a numerical assignment to a segment or a plane figure.

1.15.2 Hartshorne

Hartshorne’s modern synthetic discussion is a useful semantic control. He interprets Euclid’s equalities of figures through equal content rather than numerical area, and notes that all the Book II propositions are valid at that level. He also uses “geometrical algebra” for the systematic manipulation of such geometric contents.

Our reconstruction is deliberately slightly sharper for II.1: the final result is not only equicomplementability but exact `HilbertScissorsEq`. This is possible because II.1 is itself a direct dissection theorem.

1.15.3 Hilbert

Hilbert is relevant here mainly as a warning against collapsing distinct layers. His later arithmetic of segments introduces arithmetic operations only after substantial synthetic development. The current Book II reconstruction does not import that arithmetic and does not use Hilbert’s

later numerical surface-area theory. It remains in the incidence/order/congruence plus scissors language already available after Book I.

1.15.4 The phrase “geometric algebra”

The phrase is therefore appropriate only with a qualification. II.1 exhibits an algebraic *pattern*: partition of one side corresponds to additive decomposition of a rectangle; exchanging the two containing sides corresponds to cyclic rotation; a square is the special case in which the two containing segments are the same. These operations anticipate the formal laws of a product-like calculus.


But the objects being manipulated by Euclid are still geometric figures and relations of equality between them. The documentation should therefore resist rewriting the theorem as if Euclid had stated

$$a(b + c + d) = ab + ac + ad$$


in symbolic algebra. That equation is a modern shadow of the configuration, not the proposition itself.

1.16 Dependency Summary

The actual formal dependency chain is



The symmetric side-order branch is



New geometric axiom in II.1:	no
New proposition-local axiom:	no
New rectangle abstraction:	yes, in <code>HilbertRectangle</code>
New reusable split helper:	yes
New representative-independence theorem:	yes
Exact scissors dissection used:	yes
General equicomplementability required by II.1:	no
Numerical area semantics:	no
Segment multiplication / field theory:	no
Group IV available/used transitively:	yes
Continuity assumption in the file:	no
Proposition-local <code>sorry</code> /admit:	no
Modular build:	clean

1.17 Conclusion

Proposition II.1 is the first clean instance in Book II of a product-like law being realized entirely by geometry. A segment is divided; a rectangle is cut along corresponding parallels; exact scissors equality identifies the whole with the sum of the pieces. The rectangle layer then removes the accidental choice of a concrete drawing, so Euclid’s phrase “the rectangle contained by” becomes a representative-independent synthetic magnitude.

The formalization also isolates the two structural laws that make the language look algebraic. Additivity comes from actual dissection. Commutativity comes from cyclic rotation of the rectangle. A square is the special case in which the two containing segments agree. None of these steps requires a product of segment lengths or a numerical area map.

Thus the “geometric algebra” of II.1 is best understood not as hidden modern symbolism but as a calculus of geometric magnitudes whose algebraic laws emerge from constructions, congruence, and finite decomposition.

The representative-independent three-part theorem at the end of the current formalization is:

```

theorem euclid_proposition_2_1_three
  [HilbertIncidence Geo]
  [HilbertEuclideanPlane Geo]

  -- Concrete dissection diagram.
  (B C D E L M K N X Y P Q : Geo.Point)
  (hRect : IsRectangle Geo B C E D)
  (hCE_PQ : Geo.Congruent C E P Q)

  -- First cut.
  (hBMC : Geo.Between B M C)
  (hDLE : Geo.Between D L E)
  (hDXC : Geo.Between D X C)
  (hMXL : Geo.Between M X L)
  (hLeftPar :
    IsParallelogram Geo L D B M)
  (hRightPar :
    IsParallelogram Geo C E L M)

  -- Second cut.
  (hMNC : Geo.Between M N C)

```



```

(hLKE : Geo.Between L K E)
(hLYC : Geo.Between L Y C)
(hNYK : Geo.Between N Y K)
(hMiddlePar :
  IsParallelogram Geo K L M N)
(hFinalPar :
  IsParallelogram Geo C E K N)

-- Arbitrary representatives of the four rectangles.
(U0 U1 U2 U3 : Geo.Point)
(V0 V1 V2 V3 : Geo.Point)
(W0 W1 W2 W3 : Geo.Point)
(Z0 Z1 Z2 Z3 : Geo.Point)

(hWhole :
  IsRectangleContainedBy Geo
    U0 U1 U2 U3 B C P Q)

(hPart1 :
  IsRectangleContainedBy Geo
    V0 V1 V2 V3 B M P Q)

(hPart2 :
  IsRectangleContainedBy Geo
    W0 W1 W2 W3 M N P Q)

(hPart3 :
  IsRectangleContainedBy Geo
    Z0 Z1 Z2 Z3 N C P Q) :
HilbertScissorsEq Geo
  (hilbertParallelogramTerm Geo U0 U1 U2 U3)
  (hilbertParallelogramTerm Geo V0 V1 V2 V3 +
    (hilbertParallelogramTerm Geo W0 W1 W2 W3 +
      hilbertParallelogramTerm Geo Z0 Z1 Z2 Z3)) := by

```

Chapter 2

Proposition II.2

2.1 Introduction

Proposition II.2 is the first place where the rectangle calculus established in II.1 is reused rather than developed further. The classical proposition is visually elementary: a square on a straight line is cut by a parallel through an arbitrary point of that line, and the two resulting rectangles fill the square. The formal reconstruction preserves exactly that geometric content, but its proof architecture is deliberately different from Euclid’s.

Euclid proves II.2 directly from a square construction and a parallel through the point of division. The current Lean theorem instead imports `Proposition2_1` and applies the representative-independent binary form of II.1. One additional operation is needed: Euclid names each part rectangle with the whole side first, whereas the II.1 cut theorem returns the divided part first. The proof therefore rotates each rectangle representative by one vertex, applies II.1, and then transports the result back to Euclid’s original order.

This makes II.2 a particularly clean test of the Book II API. Two algebra-like laws are already available, but neither is primitive algebra:

- additivity is realized by an exact rectangle dissection;
- exchange of the two containing segments is realized by cyclic rotation of a rectangle.

No multiplication of segment lengths and no numerical area function occurs in the theorem.

2.2 Euclid’s Statement and Classical Configuration

2.2.1 The statement

In Heath’s wording, Proposition II.2 states:

If a straight line be cut at random, the rectangle contained by the whole and both of the segments is equal to the square on the whole.

Here “both of the segments” means the two rectangles obtained by pairing the whole line with each of the two parts. If AB is cut at C , the geometric statement is therefore

$$\text{Rect}(AB, AC) + \text{Rect}(AB, CB) = \text{Sq}(AB).$$

The notation is documentary shorthand only. It does not introduce a number-valued rectangle function.

2.2.2 The classical diagram

Euclid takes a straight line AB cut at C , constructs the square $ADEB$ on AB by I.46, and through C draws CF parallel to the two opposite sides AD and BE by I.31 [1]. The square is thereby partitioned into two rectangles. One is contained by the whole AB and the segment AC ; the other is contained by the whole AB and the segment CB .

The current Lean configuration uses different letters:

classical role	Lean role
AB	BC
C	M
$ADEB$	$BCED$
CF	ML

Thus $B - M - C$, the outer figure $BCED$ is a square, and the cut ML divides it into the left and right rectangular pieces.

Figure 2.1 shows both the configured cut and the separate representation change used by the formal proof.




Figure 2.1: Two mechanisms used in the formal II.2 proof. Panel (a) is the instantiated square $BCED$ with $B - M - C$ and the cut ML ; the dashed diagonal DC meets the cut at the hidden witness X , so the formal data include both $D - X - C$ and $M - X - L$. Panel (b) shows the cyclic presentation change $V_0V_1V_2V_3 \mapsto V_1V_2V_3V_0$, which exchanges the two adjacent side roles while preserving exact scissors content.

2.3 Classical Proof

The classical proof has only four mathematical steps.

First, construct the square on the whole segment. Second, draw through the point of section a line parallel to the appropriate pair of sides of the square. Third, observe that this line partitions the square into two rectangles. Finally, identify those rectangles by Book II Definition 1: their adjacent sides are the whole segment together with the two parts of the divided segment.

In the standard lettering this gives

$$\text{Rect}(AB, AC) + \text{Rect}(AB, CB) = \text{Sq}(AB).$$

Euclid does not invoke II.1 in this proof. The local classical dependency graph is instead

$$\begin{array}{c} \text{I.46: construct the square on } AB \\ \Downarrow \\ \text{I.31: draw the parallel through } C \\ \Downarrow \\ \text{II.Def.1: identify the two rectangles} \\ \Downarrow \\ \text{the two pieces constitute the square.} \end{array}$$

This fact is important for documentation. The formal theorem proves the same geometric identity, but it is not a line-by-line formalization of Euclid’s historical derivation.

2.4 The Geometric Identity

In the current Lean lettering the square side is BC and the cut point is M . The two Euclidean rectangles are therefore

$$\text{Rect}(BC, BM), \quad \text{Rect}(BC, MC),$$

and the theorem asserts the exact scissors relation

$$\text{Rect}(BC, BM) + \text{Rect}(BC, MC) \simeq_{\text{sc}} \text{Sq}(BC).$$

A modern algebraic shadow is obtained by writing $x = y + z$:

$$x^2 = xy + xz.$$

This is useful as a mnemonic for the pattern, but it is not the object proved by Lean and it is not a claim that Euclid has defined multiplication on segment lengths. The formal terms remain rectangle representatives and their exact scissors relation.

2.5 Reusable Book II Infrastructure

II.2 imports

```
import CGJteamLab.Proposition2_1
```

and introduces no new rectangle abstraction. Its proof consumes four pieces of the existing Book II infrastructure.

2.5.1 A square as a rectangle contained by the same side twice

From

```
hSquare : IsSquare Geo B C E D
```

the theorem

```
square_is_rectangle_contained_by_side
```

produces

```
IsRectangleContainedBy Geo B C E D B C B C
```

so the concrete square is recognized synthetically as a rectangle contained by BC, BC .

2.5.2 The binary form of II.1

The reusable distributive kernel is

```
euclid_proposition_2_1_two
```

whose geometric content is

$$\text{Rect}(BC, PQ) \simeq_{\text{sc}} \text{Rect}(BM, PQ) + \text{Rect}(MC, PQ)$$

when $B - M - C$ and the corresponding rectangular cut data are supplied.

For II.2 the fixed segment is specialized to $PQ = BC$. Thus the right-hand pieces naturally arrive in the order

$$\text{Rect}(BM, BC), \quad \text{Rect}(MC, BC).$$

2.5.3 Exchanging the containing-side order

Euclid’s statement uses the opposite order:

$$\text{Rect}(BC, BM), \quad \text{Rect}(BC, MC).$$

The current proof does not treat this as “commutativity of multiplication.” It calls

```
rectangle_contained_by_swap
```

on each arbitrary representative.

The theorem rotates a rectangle

$$A B C D \mapsto B C D A,$$

returns the corresponding `IsRectangleContainedBy` fact with the two containing segments exchanged, and simultaneously returns a `HilbertScissorsEq` relating the two cyclic presentations.

The higher-level theorem `rectangle_contained_by_comm` belongs to the same rectangle API, but the public II.2 proof uses `rectangle_contained_by_swap` directly because its returned data are exactly the swapped representatives and exact scissors transports needed for the subsequent II.1 call.

2.6 Proof Architecture of the Reconstruction

The Lean proof has four stages.

2.6.1 Step 1: recognize the square

The square hypothesis is converted to the “contained by” language:

```
have hSquareContained :
  IsRectangleContainedBy Geo
    B C E D B C B C :=
square_is_rectangle_contained_by_side
  Geo B C E D hSquare
```

The rectangle structure and the side congruence $CE \cong BC$ are then extracted from this fact.

2.6.2 Step 2: rotate the two arbitrary part representatives

The two hypotheses supplied in Euclid’s order are

```
hPart1 :
  IsRectangleContainedBy Geo
    V0 V1 V2 V3 B C B M

hPart2 :
  IsRectangleContainedBy Geo
    W0 W1 W2 W3 B C M C
```

The current source then applies `rectangle_contained_by_swap` separately to `hPart1` and `hPart2`. Each call returns a swapped `IsRectangleContainedBy` fact and an exact scissors equality for the cyclically rotated representative. The resulting representatives are $V_1V_2V_3V_0$ and $W_1W_2W_3W_0$.

The mathematical result is that the rotated representatives now have the side orders required by the II.1 binary theorem.

2.6.3 Step 3: apply II.1 with fixed segment BC

The central call is

```
have hSplit :
  HilbertScissorsEq Geo
    (hilbertParallelogramTerm Geo B C E D)
    (hilbertParallelogramTerm Geo V1 V2 V3 V0 +
      hilbertParallelogramTerm Geo W1 W2 W3 W0) :=
euclid_proposition_2_1_two
  Geo
  B C D E L M X
  B C
  hRect
```

```

hCE_BC
hBMC
hDLE
hDXC
hMXL
hLeftPar
hRightPar
B C E D
V1 V2 V3 V0
W1 W2 W3 W0
hSquareContained
hPart1Swap
hPart2Swap

```

This is the whole mathematical core of the reconstruction:

$$\text{Sq}(BC) \simeq_{\text{sc}} \text{Rect}(BM, BC) + \text{Rect}(MC, BC).$$

2.6.4 Step 4: return to Euclid's ordering

The two exact rotation equalities are added:

```

have hPartsBack :
  HilbertScissorsEq Geo
    (hilbertParallelogramTerm Geo V1 V2 V3 V0 +
     hilbertParallelogramTerm Geo W1 W2 W3 W0)
    (hilbertParallelogramTerm Geo V0 V1 V2 V3 +
     hilbertParallelogramTerm Geo W0 W1 W2 W3) :=
  HilbertScissorsEq.add
    (Geo := Geo)
    (HilbertScissorsEq.symm
     (Geo := Geo) hRotate1)
    (HilbertScissorsEq.symm
     (Geo := Geo) hRotate2)

```

Transitivity composes `hSplit` with `hPartsBack`, and the final symmetry reverses the result into the order used by the public theorem: sum of the two Euclidean rectangles on the left, square on the right.

2.7 Lean Formalization

The current public theorem is the following instantiated diagram theorem:

```

theorem euclid_proposition_2_2
  [HilbertIncidence Geo]
  [HilbertEuclideanPlane Geo]

  (B C D E L M X : Geo.Point)

  (hSquare : IsSquare Geo B C E D)

  (hBMC : Geo.Between B M C)

```

```

(hDLE : Geo.Between D L E)
(hDXC : Geo.Between D X C)
(hMXL : Geo.Between M X L)
(hLeftPar :
  IsParallelogram Geo L D B M)
(hRightPar :
  IsParallelogram Geo C E L M)

(V0 V1 V2 V3 : Geo.Point)
(W0 W1 W2 W3 : Geo.Point)

(hPart1 :
  IsRectangleContainedBy Geo
    V0 V1 V2 V3 B C B M)

(hPart2 :
  IsRectangleContainedBy Geo
    W0 W1 W2 W3 B C M C) :
HilbertScissorsEq Geo
  (hilbertParallelogramTerm Geo V0 V1 V2 V3 +
   hilbertParallelogramTerm Geo W0 W1 W2 W3)
  (hilbertParallelogramTerm Geo B C E D) := by

```

Three aspects of the signature are mathematically significant.

First, the theorem assumes an instantiated square and cut diagram. It is not an existence theorem constructing the square and the parallel cut from only a segment and a cut point. Euclid performs those constructions in the classical proof; the present public Lean theorem starts after they have been supplied.

Second, the two part rectangles are arbitrary representatives satisfying `IsRectangleContainedBy`. Their exact placement in the plane is not fixed by the statement.

Third, the conclusion is the exact relation `HilbertScissorsEq`. It is not merely a conclusion in the more general `HilbertScissorsEquicomplementable` relation. II.2 remains at the exact-dissection level because its proof is ultimately the binary exact split of II.1 plus exact representation transports.

2.8 Relationship with Proposition II.1

The contrast between the classical and formal dependency graphs is especially clear here.

Euclid's II.2 is proved directly from the square and the parallel cut. The current Lean file instead begins with

```
import CGJteamLab.Proposition2_1
```

and the public proof calls

```
euclid_proposition_2_1_two
```

once.

Thus II.2 is an explicit formal corollary of the binary II.1 API. No new dissection lemma is proved. The only additional mathematical work is conversion between the two possible orders of the containing segments.

This reuse also explains why the symmetric rectangle interface added at II.1 is not cosmetic. Without the side-order transport, the II.1 theorem would produce perfectly correct rectangles whose formal orientation does not match the language of II.2.

2.9 Relationship with Book I Infrastructure

The direct import is II.1, not a Book I proposition. Nevertheless the inherited chain underneath II.1 contains the Book I geometry on which the rectangle calculus rests.

The exact binary split is built over the scissors decomposition infrastructure developed around I.47. Rectangle structure uses Euclidean parallelogram theory and synthetic right angles; the square hypothesis uses the square infrastructure established in Book I. Consequently the formal proof of II.2 does not replay I.31 and I.46 as Euclid does. Those classical constructions have been absorbed into the established geometric API and into the hypotheses of the instantiated theorem.

2.10 Formalization Notes and Hidden Geometric Data

The visible square cut hides several obligations that are explicit parameters of the current theorem:

- $B - M - C$, fixing the orientation of the cut on the square side;
- $D - L - E$, locating the opposite endpoint of the cut;
- $D - X - C$ and $M - X - L$, recording the intersection X of the cut with the diagonal used by the inherited exact scissors split;
- `IsParallelogram Geo L D B M` for the left piece;
- `IsParallelogram Geo C E L M` for the right piece;
- two arbitrary representatives of the Euclidean part rectangles;
- two cyclic representative changes, each accompanied by an exact scissors equality.

The point X has no role in the elementary verbal statement. It belongs to the current exact-dissection interface inherited from the triangulated scissors proof. It is therefore documented as hidden formal geometry rather than presented as part of Euclid's original construction.

There is also an important negative observation. No `SameRay`, `SameSide`, or `OppositeSide` obligation appears in `Proposition2_2.lean` itself. The corresponding low-level geometry has already been discharged inside the reusable rectangle and II.1 theorems.

2.11 Geometry, Content, Representation, and Later Arithmetic

2.11.1 Geometry

The geometric layer consists of a square, betweenness on two opposite sides, the diagonal/cut witness, and two parallelogram pieces. Congruence of the square side with itself identifies the square as a rectangle contained by BC, BC .

2.11.2 Content / scissors

The conclusion and every transport used in the proof are exact `HilbertScissorsEq` relations. The proof uses symmetry, transitivity, and addition of exact scissors equalities. It does not weaken to the more general equicomplementability relation.

2.11.3 Representation

Representation is the distinctive issue in II.2. The theorem accepts arbitrary rectangles for

$$\text{Rect}(BC, BM) \quad \text{and} \quad \text{Rect}(BC, MC).$$

The binary II.1 theorem wants the factors in the reverse order. Cyclic rotation of each representative supplies the required conversion while preserving exact scissors content.

2.11.4 Later arithmetic / numerical area

Neither layer is used. In particular, the proof does not define a segment product, does not invoke Hilbert’s later segment field, and does not assign real areas to the square or rectangles.

2.12 Axiomatic and Semantic Status

The public theorem assumes

```
[HilbertIncidence Geo]
[HilbertEuclideanPlane Geo]
```

and therefore works in the same Euclidean project layer as II.1. No new proposition-specific geometric axiom or content axiom is declared in `Proposition2_2.lean`. Inspection of the current 162-line source shows one theorem declaration and no proposition-local `axiom`, `sorry`, or `admit`.

This should not be described as “axiom-free.” The theorem inherits the axioms and typeclass assumptions of the established Hilbert/Euclidean/scissors library transitively. No current `#print axioms` output was supplied with the source used for this documentation pass, so the transitive axiom list is not reproduced here.

The implementation checkpoint records that the II.2 Lean build had already been checked during theorem development. This documentation workspace does not contain the complete Lean project, so the Lean build is not re-run here; the LaTeX and Asymptote checks are reported separately.

2.13 Hartshorne’s Semantic Control

Hartshorne places the end of Book I and all of Book II inside Euclid’s theory of area, interpreted synthetically as equal content rather than as an unexplained numerical area function [5]. He also describes Book II as part of an “algebra of areas”: rectangles can play a product-like role because Euclid has no operation that multiplies two line segments to produce another segment.

II.2 fits that interpretation exactly, provided the levels are kept separate. The square on the whole and the two rectangles are geometric figures; their equality is a content statement; the modern identity $x^2 = xy + xz$ is only an interpretive shadow.

The current Lean theorem is locally stronger than the weakest equal-content reading because its configured proof stays at exact scissors equality. That strength comes from the explicit binary dissection inherited from II.1, not from a general principle identifying every equal-content pair with an exact dissection.

2.14 Hilbert’s Layering

Hilbert supplies a second semantic guardrail. His theory separates synthetic plane content from the later arithmetic of segments. Segment multiplication is introduced only in a later arithmetic construction, with a fixed unit segment and parallel geometry; it is not primitive rectangle notation.

The II.2 proof remains entirely on the earlier side of this boundary:

square and rectangle geometry
 $\Downarrow$
 exact scissors dissection
 $\Downarrow$
 representative rotation and transport.

The apparent commutativity required by the formula $xy = yx$ is realized geometrically by `rectangle_contained_by_swap`, not imported from Hilbert’s later segment arithmetic.

2.15 Dependency Summary

The actual Lean dependency chain is

Hilbert incidence + Euclidean plane layer
 $\Downarrow$
 Book I right-angle, square, parallelogram, and scissors infrastructure
 $\Downarrow$
`HilbertRectangle : IsRectangleContainedBy, rectangle_contained_by_swap`
 $\Downarrow$
 Proposition II.1 binary API
 $\Downarrow$
`euclid_proposition_2_1_two`
 $\Downarrow$
 two representative rotations + exact scissors addition/transitivity
 $\Downarrow$
`euclid_proposition_2_2.`

The classical local chain is different:

$$\text{I.46} \longrightarrow \text{I.31} \longrightarrow \text{square cut into two rectangles} \longrightarrow \text{II.2.}$$

New geometric axiom in II.2:	no
New content axiom in II.2:	no
New proposition-local helper theorem:	no
New reusable helper theorem:	no
New <code>HilbertRectangle</code> theorem introduced by II.2:	no
New Interface / Book Zero / Scissors theorem:	no
Reuses an earlier Book II proposition:	yes, II.1
Direct side-order transport used:	yes, <code>rectangle_contained_by_swap</code>
Exact <code>HilbertScissorsEq</code> :	yes
General equicomplementability required:	no
Numerical area:	no
Segment multiplication:	no
Public Lean theorem build:	previously checked; not re-run here
Full relevant Lean build:	previously checked; not re-run here
Proposition-local <code>sorry/admit</code> :	no

2.16 Conclusion

Proposition II.2 is mathematically a square cut into two rectangles, but formally it does more than repeat the picture. It demonstrates that the II.1 rectangle calculus is already reusable: the binary dissection theorem supplies additivity, and cyclic rectangle rotation reconciles the two possible orders of the containing sides.

This is also the first sharp divergence between Euclid’s local proof graph and the formal Book II graph. Euclid proves II.2 directly from I.46 and I.31. Lean uses II.1 as a library theorem and turns II.2 into a short transport-and-reuse argument. The theorem is therefore not a literal transcription of the classical proof, but it remains entirely synthetic and geometric.

The result can be remembered by the modern shadow

$$x^2 = xy + xz, \quad x = y + z,$$

but the formal content is more precise: arbitrary representatives of the two rectangles contained by the whole side and the two parts are exactly scissors-equal, in sum, to the concrete square on the whole side. No numerical area and no multiplication of segment lengths is required.

Chapter 3

Proposition II.3

3.1 Introduction

Proposition II.3 is the second immediate reuse of the rectangle calculus established in Proposition II.1. Its classical form is again a statement about a straight line cut at an arbitrary point, but now one of the two resulting segments plays a distinguished role. The rectangle contained by the whole line and that segment is decomposed into a rectangle contained by the two parts and the square on the distinguished segment.

The current Lean reconstruction makes this structure unusually transparent. Its only direct import is the II.1 module:

```
import CGJteamLab.Proposition2_1
```

The proof first recognizes the supplied square on MC as a rectangle contained by MC, MC , and then applies the binary form of II.1 with the fixed segment specialized to MC . No new dissection theorem, side-order swap, or proposition-local helper theorem is needed.

This is an important contrast with II.2. There the Euclidean order of the two containing segments did not match the order produced by the binary II.1 theorem, so cyclic rectangle rotation was required. In II.3 the order already matches:

$$\text{Rect}(BC, MC) \simeq_{\text{sc}} \text{Rect}(BM, MC) + \text{Rect}(MC, MC).$$

The square on MC is then simply a chosen representative of the final $\text{Rect}(MC, MC)$ term.

Thus II.3 is not formalized through multiplication of segment lengths or through a numerical area function. Its apparently algebraic shape is realized by an exact geometric cut together with representative independence in the rectangle API.

3.2 Euclid's Statement and Classical Configuration

3.2.1 The statement

In Heath's wording, Proposition II.3 states:

If a straight line be cut at random, the rectangle contained by the whole and one of

the segments is equal to the rectangle contained by the segments and the square on the aforesaid segment.

If AB is cut at C and the distinguished part is BC , the geometric identity is

$$\text{Rect}(AB, BC) = \text{Rect}(AC, CB) + \text{Sq}(BC).$$

As throughout the Book II documentation, Rect and Sq are expository shorthand for geometric figures and their content. They are not number-valued functions.

3.2.2 The classical diagram

Euclid constructs the square $CDEB$ on the distinguished segment CB by Proposition I.46. The side ED is produced, and through A a line is drawn parallel to the corresponding sides of the square by Proposition I.31 [1]. The resulting larger rectangle has adjacent side magnitudes AB and BC . The line through C separates it into a rectangle with side magnitudes AC, CB and the square on CB .

The current Lean lettering is related to the classical roles by

classical role	Lean role
AB	BC
C	M
AC	BM
CB	MC

so the formal cut relation is $B - M - C$ and the distinguished segment is MC . The configured outer rectangle is $BCED$, with $CE \cong MC$, while the cut through M meets the opposite side at L .

Figure 3.1 shows the geometry used by the reconstruction. The rectangle $BCED$ is drawn below the segment BC . Its right-hand piece has width MC and height congruent to MC , so it realizes the square-like $\text{Rect}(MC, MC)$ part of the dissection. The supplied square $MCFG$ is drawn on the opposite side of MC to emphasize a formal point: the conclusion may use a square representative different from the concrete piece created by the cut.

3.3 Classical Proof

Let AB be cut at C , and choose BC as the segment named in the proposition. Construct the square $CDEB$ on CB by I.46. Produce one side of the square and through A draw a line parallel to the opposite sides by I.31. This forms a larger rectangle whose adjacent side magnitudes are AB and BC .

The line through C partitions that larger rectangle into two pieces. The first has adjacent side magnitudes AC and CB , and the second is precisely the square on CB . Hence

$$\text{Rect}(AB, BC) = \text{Rect}(AC, CB) + \text{Sq}(BC).$$

This is Euclid's entire content argument: the whole rectangle is constituted by the two displayed subfigures.




Figure 3.1: The geometric mechanism of II.3. The outer rectangle $BCED$ has height $CE \cong MC$ and is split by ML at the point $B - M - C$. The dashed diagonal DC meets the cut at the hidden witness X . The right-hand cut piece represents $\text{Rect}(MC, MC)$; the separately drawn square $MCFG$ is the square representative used in the public theorem.

The local classical dependency graph is therefore

$$\begin{array}{c}
 \text{I.46: construct the square on } CB \\
 \Downarrow \\
 \text{I.31: draw the required parallel through } A \\
 \Downarrow \\
 \text{Book II Definition 1: identify the rectangles} \\
 \Downarrow \\
 \text{whole rectangle} = \text{rectangle} + \text{square} \\
 \Downarrow \\
 \text{II.3.}
 \end{array}$$

Euclid does not cite Proposition II.1 in this proof. The current Lean proof, by contrast, is explicitly organized as a corollary of the reusable II.1 binary theorem. The classical and formal dependency graphs therefore prove the same geometric identity through different library organizations.

3.4 The Geometric Identity

In the current Lean lettering, $B - M - C$. The theorem fixes the segment MC and applies it to the whole BC and the two parts BM, MC . Its exact geometric content is

$$\text{Rect}(BC, MC) \simeq_{\text{sc}} \text{Rect}(BM, MC) + \text{Sq}(MC).$$

A modern algebraic shadow is obtained by writing

$$x = y + z,$$

where x corresponds to BC , y to BM , and z to MC . Then the pattern is

$$xz = yz + z^2,$$

or equivalently

$$(y + z)z = yz + z^2.$$

This formula is useful only as a mnemonic. The Lean proof never forms the products xz or yz as segment-arithmetic objects. The actual objects are rectangle representatives and a square, related by exact scissors equality.

3.5 Reusable Book II Infrastructure

The current file begins with

```
import CGJteamLab.Proposition2_1
```

and introduces no new rectangle abstraction.

3.5.1 The binary form of II.1

The reusable theorem is

```
euclid_proposition_2_1_two
```

with geometric content

$$\text{Rect}(BC, PQ) \simeq_{\text{sc}} \text{Rect}(BM, PQ) + \text{Rect}(MC, PQ)$$

whenever $B - M - C$ and the required rectangular cut data are supplied.

For II.3 the fixed segment is specialized directly to

$$PQ = MC.$$

Therefore II.1 returns

$$\text{Rect}(BC, MC) \simeq_{\text{sc}} \text{Rect}(BM, MC) + \text{Rect}(MC, MC),$$

which already has exactly the containing-side order required by the public II.3 theorem.

3.5.2 A square as $\text{Rect}(MC, MC)$

The square hypothesis

```
hSquare : IsSquare Geo M C F G
```

is converted by

```
square_is_rectangle_contained_by_side
```

into

```
IsRectangleContainedBy Geo M C F G M C M C
```

so the supplied square is a concrete representative of the final $\text{Rect}(MC, MC)$ term.

3.5.3 What II.3 does not use

Unlike II.2, the current proof does not call

```
rectangle_contained_by_swap  
rectangle_contained_by_comm
```

The absence is mathematically informative. The binary II.1 theorem already places BM and MC first and the fixed segment MC second, exactly as II.3 needs. No geometric analogue of exchanging factors is required.

3.6 Proof Architecture of the Reconstruction

The public proof has only two mathematical stages.

3.6.1 Step 1: recognize the square

The first local fact is

```
have hSquareContained :  
  IsRectangleContainedBy Geo  
    M C F G M C M C :=  
  square_is_rectangle_contained_by_side  
    Geo M C F G hSquare
```

This moves from the square-specific geometric predicate to the general “rectangle contained by” language used by II.1.

3.6.2 Step 2: instantiate II.1 with the fixed segment MC

The remainder of the proof is one direct call:

```
exact
  euclid_proposition_2_1_two
    Geo
    B C D E L M X
    M C
    hRect
    hCE_MC
    hBMC
    hDLE
    hDXC
    hMXL
    hLeftPar
    hRightPar
    U0 U1 U2 U3
    V0 V1 V2 V3
    M C F G
    hWhole
    hProduct
    hSquareContained
```

No scissors transitivity, addition, symmetry, or representative rotation is performed locally in II.3. Those mechanisms have already been packaged inside the representative-independent binary theorem of II.1.

This makes the formal DAG exceptionally short:

$$\begin{array}{c}
 \text{square } MCFG \\
 \Downarrow \\
 \text{square_is_rectangle_contained_by_side} \\
 \Downarrow \\
 \text{Rect}(MC, MC) \text{ representative} \\
 \Downarrow \\
 \text{euclid_proposition_2_1_two with } PQ = MC \\
 \Downarrow \\
 \text{euclid_proposition_2_3.}
 \end{array}$$

3.7 Lean Formalization

The current public theorem is:

```
theorem euclid_proposition_2_3
  [HilbertIncidence Geo]
  [HilbertEuclideanPlane Geo]

  (B C D E L M X : Geo.Point)
  (hRect : IsRectangle Geo B C E D)
  (hCE_MC : Geo.Congruent C E M C)

  (hBMC : Geo.Between B M C)
```

```

(hDLE : Geo.Between D L E)
(hDXC : Geo.Between D X C)
(hMXL : Geo.Between M X L)
(hLeftPar :
  IsParallelogram Geo L D B M)
(hRightPar :
  IsParallelogram Geo C E L M)

(F G : Geo.Point)
(hSquare : IsSquare Geo M C F G)

(U0 U1 U2 U3 : Geo.Point)
(V0 V1 V2 V3 : Geo.Point)

(hWhole :
  IsRectangleContainedBy Geo
    U0 U1 U2 U3 B C M C)

(hProduct :
  IsRectangleContainedBy Geo
    V0 V1 V2 V3 B M M C) :
HilbertScissorsEq Geo
  (hilbertParallelogramTerm Geo U0 U1 U2 U3)
  (hilbertParallelogramTerm Geo V0 V1 V2 V3 +
    hilbertParallelogramTerm Geo M C F G) := by

```

Several features of this signature are mathematically significant.

First, the theorem assumes a configured outer rectangle and its cut. It does not construct that diagram from the bare data $B - M - C$. Euclid performs the square and parallel constructions in his proof; the formal theorem receives the corresponding geometry as explicit hypotheses.

Second, $U_0U_1U_2U_3$ and $V_0V_1V_2V_3$ are arbitrary representatives of

$$\text{Rect}(BC, MC) \quad \text{and} \quad \text{Rect}(BM, MC).$$

The conclusion is therefore not restricted to the literal pieces drawn in the configured cut.

Third, the square $MCFG$ is concrete, but the II.1 representative-independence machinery permits it to stand for the $\text{Rect}(MC, MC)$ piece even when it is not the internal right-hand rectangle of the supplied cut diagram.

Fourth, the conclusion is exactly

```
HilbertScissorsEq
```

rather than the more general relation

```
HilbertScissorsEquicomplementable
```

and it is not equality of real numbers.

3.8 Relationship with Earlier Book II Propositions

3.8.1 Proposition II.1

II.3 is a direct formal corollary of II.1. The import is

```
import CGJteamLab.Proposition2_1
```

and the only substantial theorem called in the final proof is

```
euclid_proposition_2_1_two
```

The specialization $PQ = MC$ converts the general binary distributive pattern into II.3 immediately.

The relationship can be written schematically as

$$\text{Rect}(BC, PQ) = \text{Rect}(BM, PQ) + \text{Rect}(MC, PQ)$$

followed by

$$PQ := MC.$$

The final term then becomes $\text{Rect}(MC, MC)$, represented by the square on MC .

3.8.2 Proposition II.2

The current Lean file does *not* import Proposition II.2, and its public theorem is not used in the proof. II.2 and II.3 are therefore sibling corollaries of the II.1 binary API rather than a formal chain $\text{II.1} \rightarrow \text{II.2} \rightarrow \text{II.3}$.

The contrast is nevertheless useful. II.2 required cyclic representative rotation because its chosen side order was opposite to the order returned by II.1. II.3 chooses the fixed segment MC in precisely the position expected by II.1, so even that transport disappears. This is evidence that the II.1 API is not merely sufficient but well aligned with the geometry of the next Book II identities.

3.9 Relationship with Book I Infrastructure

Euclid's local proof explicitly calls I.46 to construct the square and I.31 to draw the parallel. The current Lean theorem does not replay those constructions. Instead it assumes the resulting rectangle, cut, parallelograms, side congruence, and square as data.

Below II.1, however, the formal dependency chain still rests on the Book I infrastructure for rectangles, right angles, squares, parallelograms, and exact scissors decomposition. The binary split ultimately uses the exact rectangle cut machinery developed over the Book I scissors layer. Thus the difference is one of modular organization, not a change from synthetic to numerical geometry.

3.10 Formalization Notes and Hidden Geometric Data

The elementary picture hides several obligations that appear explicitly in the current theorem signature:

- $B - M - C$, fixing the cut point and its orientation on the whole side;
- $D - L - E$, locating the opposite endpoint of the cut;
- $D - X - C$ and $M - X - L$, recording the intersection X of the diagonal and the cut line used by the inherited exact scissors theorem;
- `IsParallelogram Geo L D B M` for the left piece;
- `IsParallelogram Geo C E L M` for the right piece;
- $CE \cong MC$, making the outer rectangle have the required fixed second side magnitude;
- a supplied square $MCFG$;
- arbitrary representatives of the whole rectangle and the BM -by- MC product rectangle.

The point X is invisible in Euclid’s verbal statement. It belongs to the triangulated exact-dissection interface inherited from II.1. It should therefore be read as hidden formal geometry rather than as an additional step in Euclid’s classical construction.

There are also useful negative observations. No `SameRay`, `SameSide`, or `OppositeSide` hypothesis occurs in `Proposition2_3.lean` itself, and no side-order swap is performed. Those lower-level obligations have already been absorbed by the reusable rectangle and II.1 layers.

3.11 Geometry, Content, Representation, and Later Arithmetic

3.11.1 Geometry

The geometric layer consists of the rectangle $BCED$, the cut point M , the opposite cut point L , the diagonal witness X , the two parallelogram pieces, the congruence $CE \cong MC$, and the square $MCFG$.

3.11.2 Content / scissors

The conclusion is the exact relation `HilbertScissorsEq`. The public proof does not weaken the result to general equicomplementability. Exactness is inherited from the binary rectangle split and representative transports already proved in II.1.

3.11.3 Representation

Representation is present but simpler than in II.2. The theorem accepts an arbitrary rectangle representative for $\text{Rect}(BC, MC)$ and an arbitrary representative for $\text{Rect}(BM, MC)$, together with a concrete square on MC . The binary II.1 theorem is already independent of the particular concrete rectangles appearing in the cut, so no additional representation transport is required locally.

3.11.4 Later arithmetic / numerical area

Neither layer is used. No segment product is defined, no fixed unit segment is introduced, and no real-valued area function appears. The expression $xz = yz + z^2$ is only the modern algebraic shadow of a geometric scissors identity.

3.12 Axiomatic and Semantic Status

The public theorem assumes

```
[HilbertIncidence Geo]
[HilbertEuclideanPlane Geo]
```

and therefore lives in the same Euclidean project layer as II.1 and II.2. Inspection of the current 96-line source shows one public theorem declaration and no proposition-local `axiom`, `sorry`, or `admit`.

This should not be described as “axiom-free.” The result inherits the axioms and typeclass assumptions of the established Hilbert/Euclidean/scissors library transitively. No current `#print axioms` output for II.3 was supplied with the documentation sources, so a transitive axiom list is not asserted here.

The implementation checkpoint records that the II.3 Lean theorem had already been build-checked during formalization. The present documentation pass uses the current source as authoritative but does not contain the full Lean project needed to repeat that build locally.

3.13 Hartshorne’s Semantic Control

Hartshorne’s general analysis of Euclid’s area theory states that the results of Book II are valid when Euclid’s equality of figures is read as equality of content [5]. His condensed list of selected Book II propositions does not separately single out II.3, so the source should not be cited as providing a special proof of this proposition.

The general semantic point is sufficient. II.3 compares a rectangle with the union of another rectangle and a square. In the present reconstruction this comparison is sharpened, for this configured case, to exact scissors equality. The modern identity

$$xz = yz + z^2$$

therefore belongs to Hartshorne’s “algebra of areas” only as an interpretive shadow of operations on geometric figures.

II.3 also clarifies the representative issue. The term called “the square on MC ” need not be the literal internal square-shaped piece of the cut; the formal theorem accepts the separately supplied square $MCFG$ because the rectangle calculus already proves independence of representatives at the scissors level.

3.14 Hilbert's Layering

Hilbert provides the complementary warning against reading later segment arithmetic into the proof. His synthetic theory of plane content distinguishes finite equidecomposition from the more general content relations, while his arithmetic of segments is introduced later and requires a fixed unit segment and additional parallel constructions [2].

II.3 remains entirely below that arithmetic boundary. Its formal chain is

$$\begin{array}{c} \text{rectangle and square geometry} \\ \Downarrow \\ \text{exact binary dissection from II.1} \\ \Downarrow \\ \text{representative-independent exact scissors equality.} \end{array}$$

In fact II.3 is even more direct than II.2: no geometric commutativity theorem is needed because the ordered containing segments already agree with the II.1 binary interface.

3.15 Dependency Summary

The actual Lean dependency chain is

$$\begin{array}{c} \text{Hilbert incidence + Euclidean plane layer} \\ \Downarrow \\ \text{Book I right-angle, square, parallelogram, and scissors infrastructure} \\ \Downarrow \\ \text{HilbertRectangle} \\ \text{IsRectangleContainedBy} \\ \text{square_is_rectangle_contained_by_side} \\ \Downarrow \\ \text{Proposition II.1 binary API} \\ \text{euclid_proposition_2_1_two} \\ \text{specialized by } PQ = MC \\ \Downarrow \\ \text{euclid_proposition_2_3.} \end{array}$$

The classical local chain is different:

$$\text{I.46} \longrightarrow \text{I.31} \longrightarrow \text{rectangle cut into rectangle + square} \longrightarrow \text{II.3.}$$

New geometric axiom in II.3:	no
New content axiom in II.3:	no
New proposition-local helper theorem:	no
New reusable helper theorem:	no
New <code>HilbertRectangle</code> theorem introduced by II.3:	no
New Interface / Book Zero / Scissors theorem:	no
Reuses an earlier Book II proposition:	yes, II.1
Imports II.2:	no
Direct side-order transport used:	no
Exact <code>HilbertScissorsEq</code> :	yes
General equicomplementability required:	no
Numerical area:	no
Segment multiplication:	no
Public Lean theorem build:	previously checked; not re-run here
Full relevant Lean build:	previously checked; not re-run here
Proposition-local <code>sorry/admit</code> :	no

3.16 Conclusion

Proposition II.3 is one of the cleanest tests of the Book II rectangle API. Classically, Euclid constructs a square on one segment and enlarges it by a parallel construction until the whole rectangle contained by the original line and that segment is obtained. Formally, the same identity is already contained in the binary theorem of II.1.

The decisive specialization is simply

$$PQ = MC.$$

It turns the second part rectangle of II.1 into $\text{Rect}(MC, MC)$, and the supplied square on MC provides exactly that representative. Nothing else is needed.

This also explains the precise relation between II.2 and II.3. Both are formal corollaries of II.1, but II.2 needs geometric side-order transport while II.3 does not. The result is therefore a very short Lean proof without being an algebraic proof in the modern numerical sense: the content remains rectangles, squares, congruent sides, and exact scissors equality throughout.

Chapter 4

Proposition II.4

4.1 Introduction

Proposition II.4 is the first proposition of Book II in the present reconstruction whose formal proof is naturally expressed as a composition of *two different earlier Book II propositions*. The classical theorem says that when a segment is cut into two parts, the square on the whole is equal to the two squares on the parts together with twice the rectangle contained by the parts.

The modern algebraic shadow is the familiar identity

$$(y + z)^2 = y^2 + z^2 + 2yz.$$

The formal proof, however, does not define a product of segment lengths and does not evaluate a numerical area. It remains entirely inside the rectangle and exact-scissors layer established at the beginning of Book II.

The current Lean source begins with

```
import CGJteamLab.Proposition2_2
import CGJteamLab.Proposition2_3
```

and the proof uses those imports exactly as the architecture suggests:

1. Proposition II.2 splits the square on the whole segment into two rectangles pairing the whole with the two parts;
2. Proposition II.3 expands each of those rectangles into one square and one cross rectangle;
3. representation transport identifies the two cross rectangles as two representatives of the same “rectangle contained by” the two parts;
4. addition and transitivity in `HilbertScissorsEq` combine the results.

Thus the substantial new fact at II.4 is not a new geometric construction or a new arithmetic law. It is the observation that the Book II rectangle API is already compositional enough to express the square-of-a-sum pattern using previously proved exact scissors theorems.

4.2 Euclid's Statement and Classical Configuration

4.2.1 The statement

In Heath's wording, Proposition II.4 states:

If a straight line be cut at random, the square on the whole is equal to the squares on the segments and twice the rectangle contained by the segments.

Let AB be cut at C . In the geometric shorthand used throughout this volume, the statement is

$$\text{Sq}(AB) = \text{Sq}(AC) + \text{Sq}(CB) + 2 \text{Rect}(AC, CB).$$

Here Sq and Rect are expository names for geometric figures and their content. The symbol 2 means that two equal rectangle contents occur in the decomposition; it is not scalar multiplication inside the Lean segment language.

4.2.2 The classical diagram

Euclid describes the square $ADEB$ on AB by I.46 and joins the diagonal BD . Through the cut point C he draws CF parallel to the vertical sides of the square, and through the intersection G of CF with BD he draws HK parallel to AB and DE by I.31 [1]. The square is thereby divided into four subfigures named AG , CK , HF , and GE .

The figure is the classical Euclidean configuration. It is deliberately not presented as the literal Lean input diagram. The formal theorem receives three configured rectangle-cut diagrams inherited from II.2 and II.3, rather than reconstructing this single classical picture point by point.

4.3 Classical Proof

Euclid's proof first identifies the lower-right parallelogram $CGKB$ as a square on CB . Since $CF \parallel AD$, the diagonal BD gives the relevant angle equality by I.29. The square has $BA = AD$, so I.5 gives equal base angles in triangle ADB . Combining these angle equalities and using I.6 yields

$$BC = CG.$$

Opposite sides in the surrounding parallelograms are equal by I.34, so the four sides of $CGKB$ are equal. Right-angle information from the parallels and the square then shows that $CGKB$ is a square on CB [1].

The same reasoning identifies HF as the square on AC . The two remaining subfigures AG and GE are complements about the diagonal of the outer square, and Proposition I.43 gives

$$AG = GE.$$

Moreover AG has adjacent side magnitudes AC and CG , while $CG = CB$. Hence AG is the rectangle contained by AC, CB , and $AG + GE$ is twice that rectangle. Since the four subfigures constitute the whole square $ADEB$, Euclid obtains the required equality [1].




Figure 4.1: The classical configuration of Euclid II.4. The square $ADEB$ is cut at C by CF , the diagonal BD meets it at G , and the parallel HK through G produces four subfigures. Euclid proves that HF and CK are the squares on AC and CB , while the complements AG and GE are equal; AG is the rectangle contained by AC, CB .

The local classical dependency graph is therefore

$$\begin{array}{c}
 \text{I.46: square on } AB \quad + \quad \text{I.31: required parallels} \\
 \Downarrow \\
 \text{I.29} + \text{I.5} + \text{I.6: } BC = CG \\
 \Downarrow \\
 \text{I.34: recognize the two smaller squares} \\
 \Downarrow \\
 \text{I.43: complements } AG = GE \\
 \Downarrow \\
 \text{Sq}(AB) = \text{Sq}(AC) + \text{Sq}(CB) + 2\text{Rect}(AC, CB) \\
 \Downarrow \\
 \text{II.4.}
 \end{array}$$

This classical DAG is important because it is *not* the DAG of the Lean proof. In particular, the formal reconstruction does not locally call I.43 and does not prove the two cross rectangles equal by diagonal complements. That equality has already been absorbed into the representative-independent rectangle calculus developed after Book I.

4.4 The Geometric Identity in the Current Formalization

The public Lean theorem starts with a strict cut

$$A - C - B.$$

It receives a square on AB , two square representatives on the two parts, and two arbitrary representatives of the cross rectangle contained by AC and CB .

At the level of exact scissors terms the final conclusion has the form

$$\text{Sq}(AB) \simeq_{\text{sc}} (\text{Rect}(AC, CB) + \text{Sq}(CA)) + (\text{Rect}(AC, CB) + \text{Sq}(CB)).$$

The use of CA rather than AC in the first square is only an endpoint orientation choice. Segments are unoriented for congruence purposes, so this is the same geometric side magnitude.

The two occurrences of the cross rectangle are deliberately represented by independent quadrilaterals. The theorem therefore says more than “the same literal rectangle occurs twice in one drawing.” It says that any two representatives satisfying

```
IsRectangleContainedBy Geo ... A C C B
```

may occupy the two cross-term positions, because the rectangle layer has already proved representative independence at the exact scissors level.

A modern mnemonic is

$$x = y + z \quad \implies \quad x^2 = y^2 + z^2 + 2yz.$$

This is useful for recognizing the combinatorial pattern only. No object corresponding to yz is constructed by `HilbertSegmentMul` in the II.4 proof.

4.5 Reusable Book II Infrastructure

4.5.1 Proposition II.2: split the square on the whole

The first reusable theorem gives

$$\text{Rect}(AB, AC) + \text{Rect}(AB, CB) \simeq_{\text{sc}} \text{Sq}(AB).$$

In the current source it is called exactly as

```
have hBCA : Geo.Between B C A :=
  (HilbertOrder.between_incidence A C B hACB).2.2.2.2

-----

-- II.2:
--
--   Rect(AB,AC) + Rect(AB,CB) = Square(AB) .
-----

have hII2 :
  HilbertScissorsEq Geo
```

```

(hilbertParallelogramTerm Geo U0 U1 U2 U3 +
 hilbertParallelogramTerm Geo V0 V1 V2 V3)
(hilbertParallelogramTerm Geo A B E0 D0) :=
euclid_proposition_2_2
Geo
A B D0 E0 L0 C X0
hSquareAB
hACB
hD0L0E0
hD0X0B
hCX0L0
hII2LeftPar
hII2RightPar
U0 U1 U2 U3
V0 V1 V2 V3
hAB_AC
hAB_CB

```

The final II.4 proof uses the symmetry of this relation, so the square becomes the starting point and the two whole-by-part rectangles become the two branches to be expanded.

4.5.2 Proposition II.3: expand a whole-by-part rectangle

Proposition II.3 has the generic geometric pattern

$$\text{Rect}(XY, ZY) \simeq_{\text{sc}} \text{Rect}(XZ, ZY) + \text{Sq}(ZY)$$

for a strict division point Z on XY , with the configured rectangle cut supplied explicitly.

II.4 calls this theorem twice. The right branch uses the original orientation $A - C - B$ directly:

$$\text{Rect}(AB, CB) \simeq_{\text{sc}} \text{Rect}(AC, CB) + \text{Sq}(CB).$$

The left branch reads the same line in the reverse direction $B - C - A$:

$$\text{Rect}(BA, CA) \simeq_{\text{sc}} \text{Rect}(BC, CA) + \text{Sq}(CA).$$

After endpoint reversal and rectangle rotation, its cross term is another representative of $\text{Rect}(AC, CB)$.

4.5.3 Rectangle rotation and endpoint reversal

The only side-order normalization that is mathematically visible in II.4 is

```
rectangle_contained_by_swap
```

applied to the second cross-rectangle representative. It cyclically rotates the concrete rectangle and exchanges the roles of its containing sides while returning an exact scissors certificate for the change of presentation.

Endpoint reversal is handled by the already established congruence theorem

```
CongruentSwapSecond
```

so that AB may be read as BA , AC as CA , and CB as BC when the reversed II.3 invocation requires that orientation. These steps are representation transport, not new metric statements.

4.5.4 Exact scissors addition

Once the two II.3 branches are available, the constructor

```
HilbertScissorsEq.add
```

combines them componentwise. Transitivity then attaches the resulting four-term expression to the square through the symmetric II.2 equality.

No subtraction, cancellation, positivity, content order, or numerical area operation occurs.

4.6 Proof Architecture of the Reconstruction

4.6.1 Step 1: reverse the cut orientation for the left branch

From $A - C - B$ the proof derives $B - C - A$:

```
have hBCA : Geo.Between B C A :=
  (HilbertOrder.between_incidence A C B hACB).2.2.2.2
```

This is the only explicit betweenness reversal performed locally.

4.6.2 Step 2: apply II.2 to the square on the whole

The theorem `euclid_proposition_2_2` produces the exact decomposition of the square into the two rectangles $\text{Rect}(AB, AC)$ and $\text{Rect}(AB, CB)$. Nothing about the smaller squares is used yet.

4.6.3 Step 3: prepare the reversed left II.3 call

The first II.2 rectangle is formally contained by AB, AC , while the reversed II.3 call expects BA, CA . The same concrete rectangle is repackaged using endpoint reversal. The second final cross-rectangle representative is then rotated from the side order AC, CB to CB, AC and read, after endpoint reversal, as BC, CA .

The current source makes these transports explicit:

```
have hAB_AC_rev :
  IsRectangleContainedBy Geo
    U0 U1 U2 U3 B A C A :=
And.intro
  hAB_AC.1
  (And.intro
    (CongruentSwapSecond
      Geo U0 U1 A B hAB_AC.2.1)
```

```

      (CongruentSwapSecond
        Geo U1 U2 A C hAB_AC.2.2))

-----

-- Rotate the second cross rectangle:
--
--   Rect(AC,CB) -> Rect(CB,AC).
--
-- Endpoint reversal then reads the rotated representative as
-- Rect(BC,CA), exactly what the reversed II.3 needs.
-----

have hSwapResult :=
  rectangle_contained_by_swap
    Geo
    T20 T21 T22 T23
    A C C B
    hCross2

have hCross2Swap :=
  hSwapResult.1

have hRotateCross2 :=
  hSwapResult.2

have hCross2ForLeft :
  IsRectangleContainedBy Geo
    T21 T22 T23 T20 B C C A :=
And.intro
  hCross2Swap.1
  (And.intro
    (CongruentSwapSecond
      Geo T21 T22 C B hCross2Swap.2.1)
    (CongruentSwapSecond
      Geo T22 T23 A C hCross2Swap.2.2))

```

4.6.4 Step 4: expand the left rectangle by II.3

The reversed invocation gives

$$\text{Rect}(BA, CA) \simeq_{\text{sc}} \text{Rect}(BC, CA) + \text{Sq}(CA).$$

The rotated cross term is then transported back to the arbitrary final AC, CB representative. The exact source block is:

```

have hLeftRaw :
  HilbertScissorsEq Geo
    (hilbertParallelogramTerm Geo U0 U1 U2 U3)
    (hilbertParallelogramTerm Geo T21 T22 T23 T20 +
      hilbertParallelogramTerm Geo C A F G) :=
euclid_proposition_2_3
  Geo
  B A D1 E1 L1 C X1
  hRectLeft
  hAE1_CA

```

```

hBCA
hD1L1E1
hD1X1A
hCX1L1
hLeftLeftPar
hLeftRightPar
F G
hSquareCA
U0 U1 U2 U3
T21 T22 T23 T20
hAB_AC_rev
hCross2ForLeft

-----
-- Return the left cross term to the original arbitrary
-- representative T20-T21-T22-T23.
-----

have hRotateCross2Back :
  HilbertScissorsEq Geo
    (hilbertParallelogramTerm Geo T21 T22 T23 T20)
    (hilbertParallelogramTerm Geo T20 T21 T22 T23) :=
  HilbertScissorsEq.symm
    (Geo := Geo)
  hRotateCross2

have hLeftTarget :
  HilbertScissorsEq Geo
    (hilbertParallelogramTerm Geo U0 U1 U2 U3)
    (hilbertParallelogramTerm Geo T20 T21 T22 T23 +
      hilbertParallelogramTerm Geo C A F G) :=
  HilbertScissorsEq.trans
    (Geo := Geo)
  hLeftRaw
  (HilbertScissorsEq.add
    (Geo := Geo)
    hRotateCross2Back
    (HilbertScissorsEq.refl
      (Geo := Geo)
      (hilbertParallelogramTerm Geo C A F G)))

```

4.6.5 Step 5: expand the right rectangle by II.3

The right branch needs no reversal of the whole segment. It directly produces

$$\text{Rect}(AB, CB) \simeq_{\text{sc}} \text{Rect}(AC, CB) + \text{Sq}(CB).$$

4.6.6 Step 6: add the branches and attach them to the square

The two II.3 relations are added. Finally the result is composed with the symmetric II.2 relation. The closing source is:

```

have hRight :
  HilbertScissorsEq Geo

```



```

      (hilbertParallelogramTerm Geo V0 V1 V2 V3)
      (hilbertParallelogramTerm Geo T10 T11 T12 T13 +
        hilbertParallelogramTerm Geo C B H K) :=
euclid_proposition_2_3
  Geo
  A B D2 E2 L2 C X2
  hRectRight
  hBE2_CB
  hACB
  hD2L2E2
  hD2X2B
  hCX2L2
  hRightLeftPar
  hRightRightPar
  H K
  hSquareCB
  V0 V1 V2 V3
  T10 T11 T12 T13
  hAB_CB
  hCross1

-----
-- Expand both II.2 summands simultaneously.
-----


have hExpanded :
  HilbertScissorsEq Geo
    (hilbertParallelogramTerm Geo U0 U1 U2 U3 +
      hilbertParallelogramTerm Geo V0 V1 V2 V3)
    ((hilbertParallelogramTerm Geo T20 T21 T22 T23 +
      hilbertParallelogramTerm Geo C A F G) +
      (hilbertParallelogramTerm Geo T10 T11 T12 T13 +
        hilbertParallelogramTerm Geo C B H K)) :=
  HilbertScissorsEq.add
    (Geo := Geo)
    hLeftTarget
    hRight

-----
-- Square(AB)
--   = Rect(AB,AC) + Rect(AB,CB)
--   = the four terms of II.4.
-----

exact
  HilbertScissorsEq.trans
    (Geo := Geo)
    (HilbertScissorsEq.symm
      (Geo := Geo)
      hII2)
    hExpanded

```

The actual Lean DAG is therefore



This is substantially shorter conceptually than Euclid’s local geometric DAG, but it is not an algebraic shortcut. The earlier propositions already package the relevant geometric dissections as reusable exact-scissors theorems.

4.7 Lean Formalization

The current public theorem signature is copied from the compiling `Proposition2_4.lean` source:

```

theorem euclid_proposition_2_4
  [HilbertIncidence Geo]
  [HilbertEuclideanPlane Geo]

  (A B C : Geo.Point)

  -----
  -- II.2 diagram: square on AB cut at C.
  -----

  (D0 E0 L0 X0 : Geo.Point)
  (hSquareAB : IsSquare Geo A B E0 D0)

  (hACB : Geo.Between A C B)
  (hD0LOE0 : Geo.Between D0 L0 E0)
  (hD0X0B : Geo.Between D0 X0 B)
  (hCX0LO : Geo.Between C X0 L0)

  (hII2LeftPar :
    IsParallelogram Geo L0 D0 A C)
  (hII2RightPar :
    IsParallelogram Geo B E0 L0 C)

  -- Arbitrary representatives of Rect(AB,AC) and Rect(AB,CB).
  (U0 U1 U2 U3 : Geo.Point)
  (V0 V1 V2 V3 : Geo.Point)

  (hAB_AC :
    IsRectangleContainedBy Geo
      U0 U1 U2 U3 A B A C)

  (hAB_CB :
```

```

IsRectangleContainedBy Geo
  V0 V1 V2 V3 A B C B)

-----

-- II.3 diagram for the left term.
--
-- We read the divided whole segment in the reverse direction:
--
--   B-C-A.
--
-- Thus II.3 gives
--
--   Rect(BA,CA) = Rect(BC,CA) + Square(CA).
-----

(D1 E1 L1 X1 : Geo.Point)

(hRectLeft :
  IsRectangle Geo B A E1 D1)

(hAE1_CA :
  Geo.Congruent A E1 C A)

(hD1L1E1 : Geo.Between D1 L1 E1)
(hD1X1A : Geo.Between D1 X1 A)
(hCX1L1 : Geo.Between C X1 L1)

(hLeftLeftPar :
  IsParallelogram Geo L1 D1 B C)
(hLeftRightPar :
  IsParallelogram Geo A E1 L1 C)

-- Square on CA, i.e. on the same unoriented segment as AC.
(F G : Geo.Point)
(hSquareCA : IsSquare Geo C A F G)

-----

-- II.3 diagram for the right term:
--
--   Rect(AB,CB) = Rect(AC,CB) + Square(CB).
-----

(D2 E2 L2 X2 : Geo.Point)

(hRectRight :
  IsRectangle Geo A B E2 D2)

(hBE2_CB :
  Geo.Congruent B E2 C B)

(hD2L2E2 : Geo.Between D2 L2 E2)
(hD2X2B : Geo.Between D2 X2 B)
(hCX2L2 : Geo.Between C X2 L2)

(hRightLeftPar :
  IsParallelogram Geo L2 D2 A C)
(hRightRightPar :
  IsParallelogram Geo B E2 L2 C)

```

```

-- Square on CB.
(H K : Geo.Point)
(hSquareCB : IsSquare Geo C B H K)

-----

-- Two arbitrary representatives of the repeated cross rectangle
-- Rect(AC,CB).
-----

(T10 T11 T12 T13 : Geo.Point)
(T20 T21 T22 T23 : Geo.Point)

(hCross1 :
  IsRectangleContainedBy Geo
    T10 T11 T12 T13 A C C B)

(hCross2 :
  IsRectangleContainedBy Geo
    T20 T21 T22 T23 A C C B) :

HilbertScissorsEq Geo
  (hilbertParallelogramTerm Geo A B E0 D0)
  ((hilbertParallelogramTerm Geo T20 T21 T22 T23 +
    hilbertParallelogramTerm Geo C A F G) +
    (hilbertParallelogramTerm Geo T10 T11 T12 T13 +
    hilbertParallelogramTerm Geo C B H K)) := by

```

Several aspects of the signature deserve explicit interpretation.

First, one strict division relation $A - C - B$ is shared by the theorem, but the public statement receives *three* configured cut diagrams:

- the II.2 square-cut diagram, with points D_0, E_0, L_0, X_0 ;
- the left II.3 rectangle-cut diagram, with points D_1, E_1, L_1, X_1 ;
- the right II.3 rectangle-cut diagram, with points D_2, E_2, L_2, X_2 .

The theorem does not construct these configurations from A, C, B alone. Rather, it proves the content identity once the corresponding geometric witnesses have been supplied.

Second, $U_0U_1U_2U_3$ and $V_0V_1V_2V_3$ are arbitrary representatives of the two rectangles produced at the II.2 level. They need not be literal subrectangles of the square ABE_0D_0 .

Third, the cross terms are represented independently by $T_{10}T_{11}T_{12}T_{13}$ and $T_{20}T_{21}T_{22}T_{23}$. Both satisfy the same `IsRectangleContainedBy` predicate for AC, CB . This makes the formal “twice the rectangle” statement representative-independent rather than tied to one duplicated quadrilateral.

Fourth, the conclusion is exactly

```
HilbertScissorsEq Geo ...
```

and not the weaker

```
HilbertScissorsEquicomplementtable Geo ...
```

relation. Exactness is inherited from II.2, II.3, rectangle rotation, formal addition, and transitivity.

4.8 Relationship with Earlier Book II Propositions

4.8.1 Proposition II.1

II.4 does not import `Proposition2_1` directly. Nevertheless II.1 is the underlying distributive core because both II.2 and II.3 are themselves built from its binary rectangle split. The dependency is therefore transitive:

$$\text{II.1} \longrightarrow \text{II.2} \text{ and } \text{II.3} \longrightarrow \text{II.4}.$$

This is the first point where the investment in II.1 becomes visibly compositional at the proposition level. II.2 and II.3 were sibling specializations of the binary II.1 API; II.4 uses both siblings together.

4.8.2 Proposition II.2

II.2 supplies exactly the outer decomposition needed by II.4:

$$\text{Sq}(AB) \simeq_{\text{sc}} \text{Rect}(AB, AC) + \text{Rect}(AB, CB).$$

The current II.4 source imports II.2 explicitly and calls its public theorem once.

4.8.3 Proposition II.3

II.3 supplies the expansion of each whole-by-part rectangle. The current proof calls the same public theorem twice, once after reversing the line orientation and once directly. Thus II.4 is a genuine composition theorem over the public Book II API rather than a restatement of one earlier helper.

4.9 Relationship with Book I Infrastructure

Euclid’s own local proof depends explicitly on I.46, I.31, I.29, I.5, I.6, I.34, and I.43. The formal II.4 theorem does not replay any of those arguments locally. Their geometric content has entered the current project through the established square, parallel, parallelogram, rectangle, and scissors layers and through the already proved II.2 and II.3 theorems.

The contrast with I.43 is especially instructive. Euclid needs I.43 to prove that the two complements AG and GE in one concrete square are equal. Lean instead receives two arbitrary cross-rectangle representatives and normalizes one of them with `rectangle_contained_by_swap`. Representative independence is already built into the rectangle API, so no local complement theorem is required.

The formal proof is therefore not a transcription of Euclid’s diagram. It is a proof over a stronger reusable library whose theorems were themselves proved synthetically from the established Hilbert/Book I geometry.

4.10 Formalization Notes and Hidden Geometric Data

The polished identity hides a substantial amount of explicit positional data. The current signature contains:

- $A - C - B$, fixing the strict order of the original segment;
- a complete square-cut configuration for the II.2 call;
- two complete rectangle-cut configurations for the two II.3 calls;
- three diagonal/cut intersection witnesses X_0, X_1, X_2 ;
- six parallelogram hypotheses certifying the two pieces in each cut;
- one square on the whole and one square on each part;
- two arbitrary II.2 rectangle representatives;
- two arbitrary representatives of the repeated cross rectangle.

The proof then adds two representational obligations not visible in Euclid's statement:

- reverse $A - C - B$ to $B - C - A$ for the left II.3 call;
- rotate one cross-rectangle representative and reverse segment endpoints so that the ordered side data match the reversed II.3 interface.

There is also useful negative information. No `SameRay`, `SameSide`, or `OppositeSide` datum occurs in the public II.4 proof. No new right-angle propagation theorem, no new parallel theorem, and no new scissors constructor is introduced. Those obligations are encapsulated inside the imported rectangle and Book II layers.

4.11 Geometry, Content, Representation, and Later Arithmetic

4.11.1 Geometry

The geometry consists of one segment cut $A - C - B$, one square on the whole, two squares on the parts, and three configured rectangle dissections. The classical source packages these ideas in one square diagram; the Lean theorem packages them as three reusable instances of already formalized cut theorems.

4.11.2 Content / scissors

The conclusion is exact `HilbertScissorsEq`. The proof only uses reflexivity, symmetry, transitivity, compatibility with addition, and exact scissors certificates supplied by II.2, II.3, and rectangle rotation. General equicomplementability is not required.

4.11.3 Representation

Representation is the genuinely nontrivial local layer. Endpoint order on the containing segments must be normalized for the reversed II.3 branch, and one cross rectangle must be cyclically rotated before it can be used in that branch. The final theorem nevertheless permits two completely independent representatives of $\text{Rect}(AC, CB)$.

This is the precise formal content of “twice the rectangle” in the current architecture: two scissors terms representing the same geometric rectangle magnitude occur as separate summands.

4.11.4 Later arithmetic / numerical area

Neither later layer is used. The proof imports no segment multiplication module, chooses no unit segment, proves no distributive law for a product of segment classes, and evaluates no real-valued area function.

The formula

$$(y + z)^2 = y^2 + z^2 + 2yz$$

should therefore remain explicitly labelled as the modern algebraic shadow of the synthetic construction.

4.12 Axiomatic and Semantic Status

The public theorem assumes

```
[HilbertIncidence Geo]
[HilbertEuclideanPlane Geo]
```

and therefore belongs to the same Euclidean project layer as II.1–II.3.

The current `Proposition2_4.lean` file introduces no proposition-local `axiom`, `sorry`, or `admit`. The theorem was compiled successfully in the project. The supplied transitive axiom audit is:

```
#print axioms Geometry.euclid_proposition_2_4

'Geometry.euclid_proposition_2_4' depends on axioms:
[propext, Classical.choice, bookZero_nullSegment1, Quot.sound]
```

Accordingly II.4 should not be described as “axiom-free.” The precise statement is that it introduces no new proposition-specific axiom and no `sorry`; its transitive dependencies are exactly the four declarations listed above. In particular the list contains no `sorryAx`.

The compile result reported for the current theorem is clean. A separate full `lake build` result was not part of the documentation input used here, so the status table below does not claim that stronger check.

4.13 Hartshorne’s Semantic Control

Hartshorne’s Section 22 supplies the appropriate general semantic control for Book II: Euclid’s equalities of plane figures can be interpreted through equal content without first assigning numerical areas [5]. He also explains why the algebraic-looking manipulations of rectangles and squares may be regarded as a kind of geometrical algebra.

II.4 is a particularly vivid example of that interpretation. Its modern shadow is a quadratic identity, but Euclid’s objects are still a large square, two smaller squares, and two rectangles. The current formalization sharpens the configured result from general equal content to exact scissors equality, because every step is built from exact dissection and exact representation transport.

Hartshorne’s “Brief Euclid” explicitly lists II.4 in the form that the square on the whole line equals the squares on its two segments plus twice the rectangle on the two segments [5]. Thus II.4 has a more direct Hartshorne-side textual counterpart than II.2 or II.3. The surrounding Section 22 discussion still supplies the essential semantic interpretation: Euclid’s equality is read as equal content, not as prior equality of numerical areas.

The stronger exact-scissors certificate remains a fact about the present Lean architecture. It should not be attributed to Hartshorne’s general content relation.

4.14 Hilbert’s Layering

Hilbert provides the complementary architectural warning. His plane-content theory and his later arithmetic of segments are distinct layers. Segment multiplication is introduced only later, using a fixed unit segment and parallel constructions, after which commutativity and distributivity become actual arithmetic theorems [2].

II.4 may look like the first place where one should invoke those later laws, because the shadow

$$(y + z)^2 = y^2 + z^2 + 2yz$$


contains both distributivity and commutativity. The compiling source shows that this move is unnecessary. The proof stays below segment arithmetic:

rectangle/square geometry
 $\Downarrow$
 exact II.2 and II.3 scissors identities
 $\Downarrow$
 geometric representation transport
 $\Downarrow$
 exact II.4 scissors identity.

This is a useful architectural checkpoint. Even a full quadratic-looking Book II identity can be derived from the geometric rectangle calculus without crossing into Hilbert’s later segment field.

4.15 Dependency Summary

The actual Lean dependency chain is



The classical local chain is different:

$$\text{I.46} + \text{I.31} + \text{I.29} + \text{I.5} + \text{I.6} + \text{I.34} + \text{I.43} \longrightarrow \text{II.4.}$$

New geometric axiom in II.4:	no
New content axiom in II.4:	no
New proposition-local helper theorem:	no
New reusable helper theorem:	no
New HilbertRectangle theorem introduced by II.4:	no
New Interface / Book Zero / Scissors theorem:	no
Reuses earlier Book II propositions:	yes, II.2 and II.3
Direct import of II.1:	no; inherited through II.2 and II.3
Direct side-order transport used:	yes, rectangle_contained_by_swap
Endpoint-reversal transport used:	yes
Exact HilbertScissorsEq:	yes
General equicomplementability required:	no
Numerical area:	no
Segment multiplication:	no
Public Lean theorem compile:	yes, clean
Full relevant lake build:	not separately reported here
Proposition-local sorry/admit:	no
sorryAx in reported axiom list:	no

4.16 Conclusion

Proposition II.4 is the first strong compositional test of the Book II API. Euclid proves the theorem inside one carefully constructed square: two subfigures are shown to be squares, and I.43 identifies the two remaining complements. The current Lean proof instead works one level higher. It takes II.2 as the outer square decomposition and applies II.3 to each of the two resulting rectangles.

The only local difficulty is representational. One II.3 call uses the reversed order $B - C - A$, and one cross rectangle must be rotated so that its ordered containing sides match that call. Once these transports are made, `HilbertScissorsEq.add` and transitivity close the theorem.

Consequently II.4 requires no new rectangle theorem, no new scissors theorem, no new proposition-local helper, and no segment multiplication. Its apparent quadratic algebra is already present in the synthetic rectangle calculus as a composition of exact geometric decompositions. The reported axiom audit adds no new proposition-specific debt and contains no `sorryAx`.

Chapter 5

Proposition II.5

5.1 Introduction

Proposition II.5 is the first Book II theorem in the present reconstruction to combine a *midpoint condition* with the rectangle/scissors calculus. Let C bisect AB , and let D be a second cut point. In the ordering chosen by the Lean theorem,

$$A - D - C - B,$$

so that C is the midpoint of AB and D lies in the left half. The theorem asserts, in geometric content language,

$$\text{Rect}(AD, DB) + \text{Sq}(CD) = \text{Sq}(CB).$$

The familiar modern algebraic shadow is obtained by writing the half-length as a and the displacement of the unequal cut from the midpoint as x :

$$(a - x)(a + x) + x^2 = a^2.$$

This formula is only an interpretive mnemonic. The Lean proof does not define multiplication of segment lengths and does not evaluate numerical area. Its objects are concrete rectangle and square representatives, while equality is expressed by exact scissors equivalence.

The current source begins with

```
import CGJteamLab.Proposition2_4
```

but this import fact should not be confused with the proof DAG. The proof does not invoke `euclid_proposition_2_4`. It directly reuses the binary rectangle theorem from II.1 and Proposition II.3, together with the reusable rectangle representation API. Thus II.5 provides an especially useful example of the difference between module dependency and mathematical proof dependency.

5.2 Euclid's Statement and Classical Configuration

5.2.1 The statement

In the notation used here, Euclid's proposition may be stated as follows. A straight line AB is divided equally at C and unequally at D . Then the rectangle contained by AD and DB , together

with the square on CD , is equal in content to the square on the half CB [1].

The phrase “rectangle contained by AD, DB ” names a plane figure whose adjacent sides have those magnitudes. It is not a product already formed in a segment field. Likewise, the addition in the displayed identity describes the content of two figures taken together.

5.2.2 The classical diagram

The principal figure below follows the standard Euclidean configuration with the unequal cut D chosen in the right half, so its baseline order is $A - C - D - B$. This is the mirror version of the Lean convention $A - D - C - B$; the content identity is unchanged. Euclid erects the square $CEFB$ on the half CB , joins BE , draws the parallel through D , then the horizontal through its intersection H with the diagonal, and finally the parallel through A [1].




Figure 5.1: The classical mechanism of Euclid II.5. The square $CEFB$ is built on the half CB . The point D is the unequal cut, H lies on diagonal BE , and LG is the small square on CD . The remaining part of $CEFB$ is the gnomon used in Euclid’s content argument. The formal Lean proof does not introduce a primitive gnomon object.

5.3 Classical Proof

The classical proof uses the gnomon structure of Figure 5.1. The diagonal BE and the parallels create complementary parallelograms. By I.43 the complements CH and HF are equal. Adding the same rectangle DM to the relevant figures yields an equality in which the rectangle CM can be replaced by AL : these are equal parallelograms on equal bases AC and CB and in the same parallels, using I.36 and the midpoint condition $AC = CB$ [1].

The construction also gives $DH = DB$ by the square-of-a-sum geometry already available at II.4 (in Euclid’s classical dependency chain). Consequently the figure AH is the rectangle contained by the unequal parts AD and DB . Euclid identifies that rectangle with the gnomon obtained by removing the small corner square LG from the square $CEFB$. Finally LG is the square on the segment CD , and adjoining it to the gnomon recovers the entire square on CB .

Thus the classical local DAG is schematically

$$\begin{array}{c}
\text{I.43: equal complements} \quad + \quad \text{I.36: equal parallelograms on equal bases} \\
\text{together with the II.4 corollary} \\
\Downarrow \\
\text{rectangle on } AD, DB = \text{gnomon in the square on } CB \\
\Downarrow \\
\text{add the square on } CD \\
\Downarrow \\
\text{Rect}(AD, DB) + \text{Sq}(CD) = \text{Sq}(CB).
\end{array}$$

This is *not* the Lean DAG. In particular, the current formal proof does not reconstruct the classical gnomon, does not call I.43 locally, and does not call II.4 despite importing the II.4 module.

5.4 The Geometric Identity in the Current Formalization

The public Lean theorem chooses the strict order

$$A - D - C - B$$

through two hypotheses:

```
(hMidC : HilbertIsMidpoint Geo C A B)
(hADC : Geo.Between A D C)
```

The midpoint hypothesis supplies $A - C - B$ and $AC \cong CB$. From $A - D - C$ and $A - C - B$, a private order helper derives $D - C - B$ and $A - D - B$. Thus the full four-point order is a theorem of the hypotheses, not an additional assumption.

At the exact-scissors level the target is

$$\text{Rect}(DB, AD) + \text{Sq}(DC) \simeq_{\text{sc}} \text{Sq}(CB).$$

The reversed order DB, AD rather than AD, DB is immaterial at the geometric rectangle level because side-order exchange is itself proved by rectangle rotation/transport. It is not treated as primitive commutativity of a segment product.

A notable semantic point is that no gnomon type occurs in the source. The classical gnomon is represented extensionally by formal sums of rectangle and square scissors terms. The proof then rearranges and composes those sums by `HilbertScissorsEq.add`, symmetry, transitivity, and the existing rectangle transport lemmas.

5.5 Reusable Book II Infrastructure

5.5.1 The binary rectangle split from II.1

The theorem `euclid_proposition_2_1_two` is used twice. First it splits DB at C , with AD held as the other containing side:

$$\text{Rect}(DB, AD) \simeq_{\text{sc}} \text{Rect}(DC, AD) + \text{Rect}(CB, AD).$$

Later the same binary theorem, now applied to AC split at D , assembles

$$\text{Rect}(CB, AD) + \text{Rect}(CB, DC) \simeq_{\text{sc}} \text{Rect}(CB, AC),$$

after geometric side-order transports.

5.5.2 Proposition II.3

Proposition II.3 supplies the first quadratic-looking decomposition:

$$\text{Rect}(AC, DC) \simeq_{\text{sc}} \text{Rect}(AD, DC) + \text{Sq}(DC).$$

The current source calls `euclid_proposition_2_3` directly. The midpoint congruence $AC \cong CB$ is then used to transport the left-hand rectangle to a representative contained by CB, DC .

5.5.3 Rectangle representation and transport

The proof relies on the established Book II rectangle API rather than numerical products. In particular it uses

- `rectangle_transport_scissorsEq` to change containing segments along segment congruences;
- `rectangle_contained_by_comm` and `rectangle_contained_by_swap` to exchange the two adjacent side roles geometrically;
- `rectangle_contained_by_unique` to compare arbitrary representatives of the same contained-by data;
- `square_is_rectangle_contained_by_side` to view a square as a rectangle contained by one side twice.

5.6 Proof Architecture of the Reconstruction

The production file preserves the private helper names created during the incremental proof search. The names contain `testXX`, but all these lemmas are `private`; they are implementation history, not public API. Their mathematical roles are stable and form the following chain.

5.6.1 Step 1: recover the full order and midpoint congruence

`proposition2_5_test01_order` derives

$$D - C - B, \quad A - D - B, \quad AC \cong CB, \quad CA \cong CB.$$

This is the only place where the four-point order is consolidated.

5.6.2 Step 2: apply II.3 on the left half

`proposition2_5_test02_left_decomposition` proves

$$\text{Rect}(AC, DC) \simeq_{\text{sc}} \text{Rect}(AD, DC) + \text{Sq}(DC).$$

5.6.3 Step 3: transport the midpoint side

`proposition2_5_test03_midpoint_rectangle_transport` uses $AC \cong CB$ to prove

$$\text{Rect}(AC, DC) \simeq_{\text{sc}} \text{Rect}(CB, DC).$$

By symmetry and transitivity, `proposition2_5_test04_first_assembled_identity` obtains

$$\text{Rect}(CB, DC) \simeq_{\text{sc}} \text{Rect}(AD, DC) + \text{Sq}(DC).$$

5.6.4 Step 4: split the unequal part DB

`proposition2_5_test05_db_decomposition` applies the binary II.1 theorem to $D - C - B$:

$$\text{Rect}(DB, AD) \simeq_{\text{sc}} \text{Rect}(DC, AD) + \text{Rect}(CB, AD).$$

The next two helpers use geometric commutativity to normalize the first term:

$$\text{Rect}(DB, AD) \simeq_{\text{sc}} \text{Rect}(AD, DC) + \text{Rect}(CB, AD).$$

5.6.5 Step 5: the scissors bridge

The private lemma `proposition2_5_test08_scissors_bridge` is an abstract content calculation. Given

$$P \simeq X + Y, \quad Z \simeq X + S,$$

it proves

$$P + S \simeq Y + Z.$$

Crucially, this is *not* a cancellation principle. It is obtained by adding exact scissors equalities, reassociating and commuting the formal multiset sum, and using transitivity. Applying it to the two preceding identities gives

$$\boxed{\text{Rect}(DB, AD) + \text{Sq}(DC) \simeq_{\text{sc}} \text{Rect}(CB, AD) + \text{Rect}(CB, DC)}.$$

5.6.6 Step 6: assemble the right side

`proposition2_5_test09_assemble_right_side` rotates the two right-hand rectangle representatives, applies the binary II.1 split to $A - D - C$, and rotates the result back. It proves

$$\text{Rect}(CB, AD) + \text{Rect}(CB, DC) \simeq_{\text{sc}} \text{Rect}(CB, AC).$$

5.6.7 Step 7: recognize the half-square

Finally `proposition2_5_test10_half_rectangle_is_square` uses $AC \cong CB$ and representative uniqueness to identify

$$\text{Rect}(CB, AC) \simeq_{\text{sc}} \text{Sq}(CB).$$

This is where the midpoint hypothesis enters a second time: first it controlled order, and now it turns the final rectangle into the square on the half.

5.7 Lean Formalization

5.7.1 The public theorem signature

The following is the current compiling public signature. It is intentionally large because the theorem receives the concrete rectangle-cut configurations and representatives required by the earlier Book II APIs rather than hiding them behind choice constructions.

```

theorem euclid_proposition_2_5
  [HilbertIncidence Geo]
  [HilbertEuclideanPlane Geo]

  (A B C D : Geo.Point)
  (hMidC : HilbertIsMidpoint Geo C A B)
  (hADC : Geo.Between A D C)

  -----

  -- Data used by test 04 inside test 08.
  -----

  (E4 H4 L4 X4 : Geo.Point)
  (hRect4 : IsRectangle Geo A C H4 E4)
  (hCH_DC : Geo.Congruent C H4 D C)
  (hELE4 : Geo.Between E4 L4 H4)
  (hEXC4 : Geo.Between E4 X4 C)
  (hDXL4 : Geo.Between D X4 L4)
  (hLeftPar4 : IsParallelogram Geo L4 E4 A D)
  (hRightPar4 : IsParallelogram Geo C H4 L4 D)

  -- Square on DC.
  (F G : Geo.Point)
  (hSquareDC : IsSquare Geo D C F G)

  -- Rect(AC,DC).
  (R0 R1 R2 R3 : Geo.Point)
  (hAC_DC :
    IsRectangleContainedBy Geo
      R0 R1 R2 R3 A C D C)

  -- Common Rect(AD,DC).
  (T0 T1 T2 T3 : Geo.Point)
  (hAD_DC :
    IsRectangleContainedBy Geo
      T0 T1 T2 T3 A D D C)

  -- Rect(CB,DC).
  (Z0 Z1 Z2 Z3 : Geo.Point)
  (hCB_DC :
    IsRectangleContainedBy Geo
      Z0 Z1 Z2 Z3 C B D C)

  -----

  -- Data used by test 07 inside test 08.
  -----

  (E7 H7 L7 X7 : Geo.Point)

```



```

(hRect7 : IsRectangle Geo D B H7 E7)
(hBH_AD : Geo.Congruent B H7 A D)
(hELE7 : Geo.Between E7 L7 H7)
(hEXB7 : Geo.Between E7 X7 B)
(hCXL7 : Geo.Between C X7 L7)
(hLeftPar7 : IsParallelogram Geo L7 E7 D C)
(hRightPar7 : IsParallelogram Geo B H7 L7 C)

-- Rect(DB,AD), the unequal-parts rectangle.
(P0 P1 P2 P3 : Geo.Point)
(hDB_AD :
  IsRectangleContainedBy Geo
    P0 P1 P2 P3 D B A D)

-- Rect(DC,AD), before commutativity normalization.
(Q0 Q1 Q2 Q3 : Geo.Point)
(hDC_AD :
  IsRectangleContainedBy Geo
    Q0 Q1 Q2 Q3 D C A D)

-- Rect(CB,AD).
(Y0 Y1 Y2 Y3 : Geo.Point)
(hCB_AD :
  IsRectangleContainedBy Geo
    Y0 Y1 Y2 Y3 C B A D)

-----

-- Data for test 09:
-- assemble Rect(CB,AD) + Rect(CB,DC) into Rect(CB,AC).
-----

(E9 H9 L9 X9 : Geo.Point)
(hRect9 : IsRectangle Geo A C H9 E9)
(hCH_CB9 : Geo.Congruent C H9 C B)
(hELE9 : Geo.Between E9 L9 H9)
(hEXC9 : Geo.Between E9 X9 C)
(hDXL9 : Geo.Between D X9 L9)
(hLeftPar9 : IsParallelogram Geo L9 E9 A D)
(hRightPar9 : IsParallelogram Geo C H9 L9 D)

-- Rect(CB,AC).
(W0 W1 W2 W3 : Geo.Point)
(hCB_AC :
  IsRectangleContainedBy Geo
    W0 W1 W2 W3 C B A C)

-----

-- Data for test 10: square on CB.
-----

(S0 S1 S2 S3 : Geo.Point)
(hSquareCB : IsSquare Geo S0 S1 S2 S3)
(hS01_CB : Geo.Congruent S0 S1 C B) :

HilbertScissorsEq Geo
  (hilbertParallelogramTerm Geo P0 P1 P2 P3 +
    hilbertParallelogramTerm Geo D C F G)
  (hilbertParallelogramTerm Geo S0 S1 S2 S3) := by

```

Several layers of data are visible in the signature.

1. A, B, C, D and the midpoint/betweenness hypotheses encode the one-dimensional configuration.
2. The groups labelled in the source as data for tests 04, 07, and 09 are concrete rectangle-cut witnesses needed by II.3 or II.1.
3. F, G give a square on DC .
4. $P_0P_1P_2P_3$ represents the rectangle contained by DB, AD .
5. $W_0W_1W_2W_3$ represents the intermediate rectangle contained by CB, AC .
6. $S_0S_1S_2S_3$ is an arbitrary square representative whose side is congruent to CB .

Thus the theorem proves representative-independent content equality for supplied geometric witnesses; it does not claim that all of those witnesses are constructed internally from four points alone.

5.7.2 The final three-stage assembly

After all private helpers have been established, the public theorem is short in conceptual structure. The current source builds exactly three equations:

1. h08: unequal-parts rectangle plus the DC square equals two rectangles with common side CB ;
2. h09: those two rectangles assemble into the rectangle $\text{Rect}(CB, AC)$;
3. h10: midpoint congruence turns that rectangle into the square on CB .

The final transitivity block is:

```

have h08 :
  HilbertScissorsEq Geo
    (hilbertParallelogramTerm Geo P0 P1 P2 P3 +
     hilbertParallelogramTerm Geo D C F G)
    (hilbertParallelogramTerm Geo Y0 Y1 Y2 Y3 +
     hilbertParallelogramTerm Geo Z0 Z1 Z2 Z3) :=
  proposition2_5_test08_core_identity
  Geo
  A B C D
  hMidC hADC
  E4 H4 L4 X4
  hRect4 hCH_DC hELE4 hEXC4 hDXL4
  hLeftPar4 hRightPar4
  F G hSquareDC
  R0 R1 R2 R3
  hAC_DC
  T0 T1 T2 T3
  hAD_DC
  Z0 Z1 Z2 Z3
  hCB_DC
  E7 H7 L7 X7

```

```

hRect7 hBH_AD hELE7 hEXB7 hCXL7
hLeftPar7 hRightPar7
P0 P1 P2 P3
hDB_AD
Q0 Q1 Q2 Q3
hDC_AD
Y0 Y1 Y2 Y3
hCB_AD

-----

-- test09:
-- Rect(CB,AD) + Rect(CB,DC) = Rect(CB,AC).
-----

have h09 :
  HilbertScissorsEq Geo
    (hilbertParallelogramTerm Geo Y0 Y1 Y2 Y3 +
     hilbertParallelogramTerm Geo Z0 Z1 Z2 Z3)
    (hilbertParallelogramTerm Geo W0 W1 W2 W3) :=
proposition2_5_test09_assemble_right_side
  Geo
  A B C D
  hADC
  E9 H9 L9 X9
  hRect9 hCH_CB9 hELE9 hEXC9 hDXL9
  hLeftPar9 hRightPar9
  Y0 Y1 Y2 Y3
  hCB_AD
  Z0 Z1 Z2 Z3
  hCB_DC
  W0 W1 W2 W3
  hCB_AC

-----

-- test10:
-- Rect(CB,AC) = Square(CB).
-----

have h10 :
  HilbertScissorsEq Geo
    (hilbertParallelogramTerm Geo W0 W1 W2 W3)
    (hilbertParallelogramTerm Geo S0 S1 S2 S3) :=
proposition2_5_test10_half_rectangle_is_square
  Geo
  A B C D
  hMidC hADC
  W0 W1 W2 W3
  hCB_AC
  S0 S1 S2 S3
  hSquareCB
  hS01_CB

-----

-- Full II.5.
-----

exact
  HilbertScissorsEq.trans

```

```
(Geo := Geo)
h08
(HilbertScissorsEq.trans
  (Geo := Geo)
  h09
```

The formal result is therefore exact `HilbertScissorsEq`, not the weaker relation `HilbertScissorsEquicomplementable`, and not equality of real numbers.

5.8 Relationship with Earlier Book II Propositions

5.8.1 Proposition II.1

II.1 is the basic additive engine of the proof. Its binary theorem is called directly in the decomposition of DB and again in the assembly of AC . Thus II.5 genuinely reuses the rectangle split calculus rather than reproving the corresponding dissections from Book I geometry.

5.8.2 Proposition II.3

II.3 is also called directly. It supplies the identity

$$\text{Rect}(AC, DC) = \text{Rect}(AD, DC) + \text{Sq}(DC),$$

at the exact scissors level. This is the step that inserts the square on the difference between the two section points.

5.8.3 Proposition II.4

The source file imports `CGJteamLab.Proposition2_4`, so II.4 is a direct *module* dependency. Nevertheless the theorem `euclid_proposition_2_4` is not invoked in the proof term. This sharply contrasts with Euclid’s classical proof, where a corollary of II.4 participates in the identification of the relevant rectangle and small square.

This distinction should be recorded explicitly:

module chain: II.4 module $\rightarrow$ II.5 module,
Lean theorem calls: II.1 binary API + II.3 + rectangle API.

5.9 Relationship with Book I Infrastructure

The Book I geometry is present below the reusable rectangle layer. Rectangles, squares, right angles, parallel sides, parallelograms, and finite scissors terms all ultimately depend on the established Hilbert/Book I development. II.5 does not locally replay Euclid’s diagonal-complement argument from I.43. Instead, the effect of those earlier geometric facts has already been packaged into `HilbertRectangle`, the scissors API, and the earlier Book II theorems.

This is an architectural gain: a proposition whose classical proof is diagram specific becomes a composition of stable content-level interfaces.

5.10 Formalization Notes and Hidden Geometric Data

5.10.1 Order orientation

The classical diagram commonly places the unequal cut in the right half $A - C - D - B$, whereas the Lean theorem fixes $A - D - C - B$. These are mirror cases. The source makes the chosen orientation explicit through `hADC` and then derives $D - C - B$ from the midpoint betweenness. No appeal to a picture is allowed to supply this ordering silently.

5.10.2 Midpoint as geometry, not arithmetic

`HilbertIsMidpoint` contributes both betweenness and segment congruence. The proof uses those components separately. Betweenness creates the required subsegment orders; congruence transports rectangle sides and eventually recognizes the square on the half. There is no variable standing for a numerical half-length.

5.10.3 Representative independence

The objects named “the rectangle contained by” particular segments are concrete quadrilateral representatives. Different helpers may receive different representatives of the same pair of side magnitudes. `rectangle_transport_scissorsEq`, `rectangle_contained_by_unique`, and the side-order transport theorems are therefore mathematically substantive parts of the formal statement, not mere syntactic rewrites.

5.10.4 No gnomon primitive

Euclid’s proof naturally names a gnomon. The present library does not define a `GnomonTerm` or another primitive gnomon object for II.5. No such new abstraction is needed here: the required content is expressed as a formal sum of scissors terms. The private scissors bridge then performs the exact content rearrangement that the classical gnomon packages visually.

5.10.5 No cancellation axiom

At an intermediate stage the two identities contain the common rectangle $\text{Rect}(AD, DC)$. It might be tempting to introduce a cancellation principle for content. The actual source deliberately avoids this. The scissors bridge combines equalities additively and rearranges formal sums, so no new positivity or cancellation theorem is required.

5.11 Axiomatic and Semantic Status

The public theorem assumes

```
[HilbertIncidence Geo]
[HilbertEuclideanPlane Geo]
```

and the production file introduces no new proposition-specific `axiom`, `sorry`, or `admit`. The user-verified module build is clean. The reported transitive axiom audit is


```
#print axioms Geometry.euclid_proposition_2_5

'Geometry.euclid_proposition_2_5' depends on axioms: [propext,
Classical.choice,
Geometry.bookZero_nullSegment1,
Quot.sound]
```

Accordingly II.5 should not be described as “axiom-free.” The precise claim is that II.5 introduces no new proposition-specific axiom and no `sorry`; its transitive dependencies are the four declarations shown above, and the list contains no `sorryAx`.

5.12 Dependency Summary

The actual Lean dependency chain is best displayed as



The classical local chain is different:

$$\text{I.43} + \text{I.36} + \text{II.4 corollary} \longrightarrow \text{gnomon identification} \longrightarrow \text{II.5.}$$

The formal proof is therefore not a line-by-line translation of the classical one. It reconstructs the same geometric content through a different, reusable DAG.

New geometric axiom in II.5:	no
New content axiom in II.5:	no
New proposition-local helper theorem:	yes, private helpers
New reusable public helper theorem:	no
New <code>HilbertRectangle</code> theorem:	no
New Interface / Book Zero / Scissors theorem:	no
Directly reuses earlier Book II theorem:	yes, II.1 and II.3
Imports II.4 module:	yes
Calls <code>euclid_proposition_2_4</code> :	no
Uses exact <code>HilbertScissorsEq</code> :	yes
Uses general equicomplementability:	no
Uses numerical area:	no
Uses segment multiplication:	no
Public theorem compiles:	yes
Module build:	clean (user verified)
Proposition-local <code>sorry/admit</code> :	no
Transitive <code>sorryAx</code> :	absent in reported audit

5.13 Conclusion

Proposition II.5 is an important architectural checkpoint for Book II. The classical proof organizes the equality around a gnomon inside the square on the half. The formal proof shows that a separate gnomon primitive is not needed to recover the theorem: the same content can be assembled from the binary II.1 rectangle split, II.3, midpoint congruence, exact rectangle transport, and the additive structure of `HilbertScissorsEq`.

The result also clarifies the role of the modern quadratic identity. It is a useful shadow for recognizing the proposition, but the verified argument stays synthetic throughout. Geometry supplies the order and congruences; representation supplies rectangles and squares; scissors equality supplies the content calculus. Numerical area and segment multiplication remain outside the proof.

Chapter 6

Proposition II.6

6.1 Introduction

Proposition II.6 is the external-section companion of Proposition II.5. A straight line AB is bisected at C and then produced beyond B to D . Thus the one-dimensional order used by the formal theorem is

$$A - C - B - D.$$

The proposition asserts, in geometric content language,

$$\text{Sq}(BC) + \text{Rect}(AD, BD) = \text{Sq}(CD).$$

The familiar modern algebraic shadow is obtained by writing the half-length $AC = CB$ as a and the added segment BD as x :

$$a^2 + (2a + x)x = (a + x)^2.$$

Equivalently, if $y = CD$ and $z = CB$, one sees the difference-of-squares pattern

$$(y + z)(y - z) + z^2 = y^2.$$

Both formulas are documentary mnemonics only. The Lean proof does not define a multiplication of segment lengths and does not evaluate a numerical area. Its objects are concrete rectangle and square representatives, and the conclusion is exact scissors equality.

The production source begins with

```
import CGJteamLab.Proposition2_5
```

so Proposition II.5 is a module dependency. It is not, however, a theorem dependency of the proof term. The current II.6 proof directly calls the binary theorem from II.1, Proposition II.3, Proposition II.4, and the rectangle-transport/scissors API. This distinction between the module chain and the actual theorem-call DAG is important in the present reconstruction.

6.2 Euclid's Statement and Classical Configuration

6.2.1 The statement

In Heath's wording, the proposition begins:

If a straight line be bisected and a straight line be added to it in a straight line, the rectangle contained by the whole with the added straight line and the added straight line together with the square on the half is equal to the square on the straight line made up of the half and the added straight line.

With AB bisected at C and BD added in a straight line, the concrete statement is

$$\text{Rect}(AD, DB) + \text{Sq}(CB) = \text{Sq}(CD) \quad [1].$$

As throughout the present Book II documentation, the notation $\text{Rect}(-, -)$ and $\text{Sq}(-)$ is documentary shorthand for geometric figures. It is not a definition of a numerical product or numerical square.

6.2.2 The classical diagram

Euclid constructs the square $CEFD$ on CD , joins the diagonal DE , draws through B a parallel to CE (or DF), then through the resulting intersection H a line parallel to the baseline, and finally through A a parallel to the vertical sides. In the annotated diagram we also mark N as the midpoint of BD , so that the traditional name of the gnomon NOP has a literal set of boundary labels on the figure [1]. The construction is shown in Figure 6.1.




Figure 6.1: The classical mechanism of Euclid II.6. The baseline has order $A - C - B - D$ and C bisects AB . The square $CEFD$ is built on CD . The point N is the midpoint of BD . The dashed square LG is equal to the square on BC ; the shaded remainder is Euclid's gnomon NOP , and the auxiliary boundary labels N, O, P are shown explicitly on the figure. The rectangle AM has adjacent side magnitudes AD and DB .

The same picture exposes the central geometry. Since $AC = CB$, the parallelograms AL and CH have equal content by I.36. The diagonal construction makes CH and HF complementary parallelograms, so I.43 gives their equality. Thus $AL = HF$. Adding the same rectangle CM identifies the whole rectangle AM with the gnomon NOP .

6.3 Classical Proof

The classical proof can be organized into four content steps.

First, the midpoint condition gives

$$AC = CB.$$

The parallelograms AL and CH stand on these equal bases and lie between the same parallels, so I.36 yields

$$AL = CH.$$

Second, CH and HF are complements about the diagonal DE of the relevant parallelogram configuration, hence I.43 gives

$$CH = HF.$$

Therefore

$$AL = HF.$$

Third, Euclid adds the same rectangle CM to both sides. The whole figure AM is therefore equal to the gnomon NOP . The construction gives $DM = DB$, so AM is the rectangle contained by AD and DB . Hence

$$\text{Rect}(AD, DB) = \text{gnomon } NOP.$$

Finally, the figure LG is the square on BC . Adding it to both sides gives

$$\text{Rect}(AD, DB) + \text{Sq}(BC) = \text{gnomon } NOP + \text{Sq}(BC).$$

The right-hand side is the whole square $CEFD$ on CD , proving II.6 [1].

The classical local DAG is therefore

$$\begin{array}{c} \text{I.36: equal parallelograms on equal bases} \quad + \quad \text{I.43: equal complements} \\ \Downarrow \\ \text{Rect}(AD, DB) = \text{gnomon } NOP \\ \Downarrow \\ \text{add the square on } BC \\ \Downarrow \\ \text{Rect}(AD, DB) + \text{Sq}(BC) = \text{Sq}(CD). \end{array}$$

This is not the Lean DAG. The production proof does not construct a gnomon object and does not call I.36 or I.43 locally.

6.4 The Geometric Identity in the Current Formalization

The public theorem receives

```
(A B C D : Geo.Point)
(hMidC : HilbertIsMidpoint Geo C A B)
(hABD : Geo.Between A B D)
```

so C is the midpoint of AB and B lies between A and D . The midpoint hypothesis gives

$$A - C - B \quad \text{and} \quad AC \cong CB.$$

Combining $A - C - B$ with $A - B - D$ by Hilbert betweenness transitivity yields

$$C - B - D \quad \text{and} \quad A - C - D.$$

Thus the full order

$$A - C - B - D$$

is derived, not read from a diagram.

At the exact-scissors level the public target is

$$\text{Sq}(BC) + \text{Rect}(AD, BD) \simeq_{\text{sc}} \text{Sq}(CD).$$

The order of the two summands on the left is the one present in the actual Lean conclusion. The theorem proves `HilbertScissorsEq`; it does not merely prove the weaker general content relation and it does not assert an equality of real numbers.

A second formal distinction is important. The square on BC in the public conclusion is the concrete square BCS_0S_1 . The rectangle contained by AD, BD is represented by the supplied quadrilateral $P_0P_1P_2P_3$. The statement is therefore about concrete representatives whose side magnitudes satisfy the required contained-by predicates.

6.5 Reusable Book II Infrastructure Used

6.5.1 Proposition II.3: extend the unequal-parts rectangle

The first major exact dissection is obtained by applying `euclid_proposition_2_3` to AD cut at B , with BD as the fixed second side:

$$\text{Rect}(AD, BD) \simeq_{\text{sc}} \text{Rect}(AB, BD) + \text{Sq}(BD).$$

This is the external-section analogue of introducing a square term in the earlier Book II identities.

6.5.2 Proposition II.1: split the old segment at its midpoint

The binary theorem `euclid_proposition_2_1_two` is then applied to AB cut at C , again with BD as the fixed side:

$$\text{Rect}(AB, BD) \simeq_{\text{sc}} \text{Rect}(AC, BD) + \text{Rect}(CB, BD).$$

Composing with the II.3 decomposition gives

$$\text{Rect}(AD, BD) \simeq_{\text{sc}} (\text{Rect}(AC, BD) + \text{Rect}(CB, BD)) + \text{Sq}(BD).$$

6.5.3 Midpoint transport

The midpoint congruence $AC \cong CB$ is converted into equality of rectangle content by `rectangle_transport_scissorsEq`:

$$\text{Rect}(AC, BD) \simeq_{\text{sc}} \text{Rect}(CB, BD).$$

Consequently

$$\text{Rect}(AD, BD) \simeq_{\text{sc}} (\text{Rect}(CB, BD) + \text{Rect}(CB, BD)) + \text{Sq}(BD).$$

No numerical factor 2 is introduced. The phrase “twice the rectangle” means two scissors terms with the same geometric content.

6.5.4 Proposition II.4 on the produced segment

The decisive second decomposition calls `euclid_proposition_2_4` directly on CD cut at B . The formal substitution is

$$A_{\text{II.4}} := C, \quad B_{\text{II.4}} := D, \quad C_{\text{II.4}} := B.$$

The required order hypothesis is exactly the derived betweenness $C - B - D$. The result is

$$\text{Sq}(CD) \simeq_{\text{sc}} (\text{Rect}(CB, BD) + \text{Sq}(BC)) + (\text{Rect}(CB, BD) + \text{Sq}(BD)).$$

This use of II.4 is one of the sharpest differences between II.5 and II.6 in the current formal library. II.5 imported the II.4 module but did not call `euclid_proposition_2_4`; II.6 calls it explicitly.

6.6 Proof Architecture of the Reconstruction

The production file preserves the names of the private incremental lemmas used during proof development. Their `testXX` names are implementation history only; because the declarations are `private`, they do not form part of the public API.

6.6.1 Step 1: recover the full order

`proposition2_6_test01_order` derives

$$C - B - D, \quad A - C - D, \quad AC \cong CB, \quad CA \cong CB.$$

The only input is the midpoint structure together with $A - B - D$.

6.6.2 Step 2: decompose the rectangle on AD, BD

`proposition2_6_test02_ad_decomposition` calls II.3 and proves

$$\text{Rect}(AD, BD) \simeq_{\text{sc}} \text{Rect}(AB, BD) + \text{Sq}(BD).$$

6.6.3 Step 3: split AB at C

`proposition2_6_test03_ab_decomposition` calls the binary II.1 theorem:

$$\text{Rect}(AB, BD) \simeq_{\text{sc}} \text{Rect}(AC, BD) + \text{Rect}(CB, BD).$$

6.6.4 Step 4: compose the two decompositions

`proposition2_6_test04_compose_decompositions` uses the same concrete representative of $\text{Rect}(AB, BD)$ in both preceding lemmas. Therefore no representative-uniqueness transport is needed at this junction. The result is

$$\text{Rect}(AD, BD) \simeq_{\text{sc}} (\text{Rect}(AC, BD) + \text{Rect}(CB, BD)) + \text{Sq}(BD).$$

6.6.5 Step 5: use the midpoint congruence

`proposition2_6_test05_midpoint_rectangle_transport` proves

$$\text{Rect}(AC, BD) \simeq_{\text{sc}} \text{Rect}(CB, BD)$$

by transporting the first adjacent side along $AC \cong CB$ while the second side remains congruent to BD .

6.6.6 Step 6: normalize the left decomposition

`proposition2_6_test06_double_cross` inserts the midpoint transport into Step 4:

$$\boxed{\text{Rect}(AD, BD) \simeq_{\text{sc}} (\text{Rect}(CB, BD) + \text{Rect}(CB, BD)) + \text{Sq}(BD).}$$

The proof uses only addition, reflexivity, and transitivity of exact scissors equality.

6.6.7 Step 7: expand the square on CD by II.4

`proposition2_6_test07_square_cd_decomposition` applies II.4 to the segment CD cut at B :

$$\boxed{\text{Sq}(CD) \simeq_{\text{sc}} (\text{Rect}(CB, BD) + \text{Sq}(BC)) + (\text{Rect}(CB, BD) + \text{Sq}(BD)).}$$

A useful implementation decision is visible here. The final assembly supplies the *same* rectangle representative $Z_0Z_1Z_2Z_3$ to both occurrences of $\text{Rect}(CB, BD)$, and it reuses the same supplied square on BD as in Step 6. Hence the following algebraic bridge can match the two decompositions literally.

6.6.8 Step 8: the abstract scissors bridge

`proposition2_6_test08_scissors_bridge` is purely algebraic at the level of formal scissors sums. For terms P, Q, R, S, T , it assumes

$$P \simeq (R + R) + T, \quad Q \simeq (R + S) + (R + T),$$

and proves

$$S + P \simeq Q.$$

The only rearrangement is associativity and commutativity of the formal sum, implemented by `ac_rfl`. The scissors equalities themselves are combined by `HilbertScissorsEq.add`, reflexivity, symmetry, and transitivity.

There is no cancellation theorem here. In particular, the proof does not infer equality by subtracting the common copies of R or T . It simply adds and rearranges exact scissors equalities.

6.6.9 Step 9: assemble II.6

`proposition2_6_test09_assembly` makes the literal substitution

$$\begin{aligned} P &:= \text{Rect}(AD, BD), \\ Q &:= \text{Sq}(CD), \\ R &:= \text{Rect}(CB, BD), \\ S &:= \text{Sq}(BC), \\ T &:= \text{Sq}(BD), \end{aligned}$$

into the abstract bridge. Steps 6 and 7 are exactly its two hypotheses, so the conclusion is the proposition:

$$\text{Sq}(BC) + \text{Rect}(AD, BD) \simeq_{\text{sc}} \text{Sq}(CD).$$

6.7 Lean Formalization

6.7.1 The public theorem signature

The following is the current compiling public signature, copied from the production source. It is intentionally explicit: the theorem receives the concrete cut diagrams and rectangle/square representatives required by the reused Book II APIs.

```

theorem euclid_proposition_2_6
  [HilbertIncidence Geo]
  [HilbertEuclideanPlane Geo]

  (A B C D : Geo.Point)
  (hMidC : HilbertIsMidpoint Geo C A B)
  (hABD : Geo.Between A B D)

  -----
  -- Data for test 06.
  -----

  (E2 H2 L2 X2 : Geo.Point)
  (hRect2 : IsRectangle Geo A D H2 E2)
  (hDH_BD2 : Geo.Congruent D H2 B D)
  (hELE2 : Geo.Between E2 L2 H2)
  (hEXD2 : Geo.Between E2 X2 D)
  (hBXL2 : Geo.Between B X2 L2)
  (hLeftPar2 : IsParallelogram Geo L2 E2 A B)
  (hRightPar2 : IsParallelogram Geo D H2 L2 B)

  -- Square on BD, shared by tests 06 and 07.
  (F G : Geo.Point)
  (hSquareBD : IsSquare Geo B D F G)

  -- Rect(AD,BD).
  (P0 P1 P2 P3 : Geo.Point)
  (hAD_BD :
    IsRectangleContainedBy Geo
      P0 P1 P2 P3 A D B D)

  -- Rect(AB,BD).
  (V0 V1 V2 V3 : Geo.Point)
  (hAB_BD :
    IsRectangleContainedBy Geo
      V0 V1 V2 V3 A B B D)

  (E3 H3 L3 X3 : Geo.Point)
  (hRect3 : IsRectangle Geo A B H3 E3)
  (hBH_BD3 : Geo.Congruent B H3 B D)
  (hELE3 : Geo.Between E3 L3 H3)
  (hEXB3 : Geo.Between E3 X3 B)
  (hCXL3 : Geo.Between C X3 L3)

```

```
(hLeftPar3 : IsParallelogram Geo L3 E3 A C)
(hRightPar3 : IsParallelogram Geo B H3 L3 C)
```

```
-- Rect(AC,BD).
```

```
(Y0 Y1 Y2 Y3 : Geo.Point)
```

```
(hAC_BD :
```

```
  IsRectangleContainedBy Geo
    Y0 Y1 Y2 Y3 A C B D)
```

```
-- The single representative R = Rect(CB,BD).
```

```
(Z0 Z1 Z2 Z3 : Geo.Point)
```

```
(hCB_BD :
```

```
  IsRectangleContainedBy Geo
    Z0 Z1 Z2 Z3 C B B D)
```

```
-----
-- Data for test 07: II.4 on CD cut at B.
-----
```

```
(D0 E0 L0 X0 : Geo.Point)
```

```
(hSquareCD : IsSquare Geo C D E0 D0)
```

```
(hD0L0E0 : Geo.Between D0 L0 E0)
```

```
(hD0X0D : Geo.Between D0 X0 D)
```

```
(hBX0L0 : Geo.Between B X0 L0)
```

```
(hII4LeftPar :
```

```
  IsParallelogram Geo L0 D0 C B)
```

```
(hII4RightPar :
```

```
  IsParallelogram Geo D E0 L0 B)
```

```
-- Representatives required internally by II.4.
```

```
(J0 J1 J2 J3 : Geo.Point)
```

```
(K0 K1 K2 K3 : Geo.Point)
```

```
(hCD_CB :
```

```
  IsRectangleContainedBy Geo
    J0 J1 J2 J3 C D C B)
```

```
(hCD_BD :
```

```
  IsRectangleContainedBy Geo
    K0 K1 K2 K3 C D B D)
```

```
-----
-- Left II.3 decomposition inside II.4.
-----
```

```
(D1 E1 L1 X1 : Geo.Point)
```

```
(hRectLeft :
```

```
  IsRectangle Geo D C E1 D1)
```

```
(hCE1_BC :
```

```
  Geo.Congruent C E1 B C)
```

```
(hD1L1E1 : Geo.Between D1 L1 E1)
```

```
(hD1X1C : Geo.Between D1 X1 C)
```

```

(hBX1L1 : Geo.Between B X1 L1)

(hLeftLeftPar :
  IsParallelogram Geo L1 D1 D B)

(hLeftRightPar :
  IsParallelogram Geo C E1 L1 B)

-- Square on BC.
(S0 S1 : Geo.Point)
(hSquareBC : IsSquare Geo B C S0 S1)

-----

-- Right II.3 decomposition inside II.4.
-----

(D2 E4 L4 X4 : Geo.Point)

(hRectRight :
  IsRectangle Geo C D E4 D2)

(hDE4_BD :
  Geo.Congruent D E4 B D)

(hD2L4E4 : Geo.Between D2 L4 E4)
(hD2X4D : Geo.Between D2 X4 D)
(hBX4L4 : Geo.Between B X4 L4)

(hRightLeftPar :
  IsParallelogram Geo L4 D2 C B)

(hRightRightPar :
  IsParallelogram Geo D E4 L4 B) :

HilbertScissorsEq Geo
  (hilbertParallelogramTerm Geo B C S0 S1 +
   hilbertParallelogramTerm Geo P0 P1 P2 P3)
  (hilbertParallelogramTerm Geo C D E0 D0)

```

The large signature falls into three geometric groups.

1. The points A, B, C, D , midpoint data, and $A - B - D$ encode the baseline configuration.
2. The first block supplies the two exact decompositions used to reach Step 6: II.3 on AD cut at B , II.1 on AB cut at C , a square on BD , and representatives of the three required rectangles.
3. The second block supplies the II.4 diagram on CD cut at B , including its two internal II.3 configurations, the square on BC , and the intermediate rectangle representatives required by II.4.

The conclusion itself is much smaller than the witness data:

```

HilbertScissorsEq Geo
  (hilbertParallelogramTerm Geo B C S0 S1 +

```



```

hilbertParallelogramTerm Geo P0 P1 P2 P3)
(hilbertParallelogramTerm Geo C D E0 D0)

```

and is exactly the formal version of

$$\text{Sq}(BC) + \text{Rect}(AD, BD) = \text{Sq}(CD).$$

6.7.2 The final bridge call

After the two large decompositions have been established, the conceptual end of the proof is only the following substitution into the abstract bridge:

```

exact
  proposition2_6_test08_scissors_bridge
    Geo
      (hilbertParallelogramTerm Geo P0 P1 P2 P3)
      (hilbertParallelogramTerm Geo C D E0 D0)
      (hilbertParallelogramTerm Geo Z0 Z1 Z2 Z3)
      (hilbertParallelogramTerm Geo B C S0 S1)
      (hilbertParallelogramTerm Geo B D F G)
    h06
    h07

```

This makes the proof architecture unusually transparent: geometry produces two normal forms, and the last step is an exact-scissors rearrangement of those normal forms.

6.8 Relationship with Earlier Book II Propositions

6.8.1 Proposition II.1

II.1 is called directly through `euclid_proposition_2_1_two`. Its role is to split $\text{Rect}(AB, BD)$ at the midpoint C . Thus the basic rectangle-distribution theorem remains the additive engine of Book II.

6.8.2 Proposition II.3

II.3 is also called directly. Applied to AD cut at B , it introduces the square on the added segment BD :

$$\text{Rect}(AD, BD) = \text{Rect}(AB, BD) + \text{Sq}(BD)$$

at the exact-scissors level.

6.8.3 Proposition II.4

II.4 is a direct theorem dependency and supplies the expansion of the square on CD after B has been proved to lie between C and D . Since the current II.4 theorem itself composes earlier Book II infrastructure, II.6 inherits that established internal structure rather than rebuilding it.

6.8.4 Proposition II.5

The production file imports `CGJteamLab.Proposition2_5`, so the II.5 module lies immediately below II.6 in the module graph. Nevertheless `euclid_proposition_2_5` is not called by any of the II.6 private helpers or by the public theorem.

Mathematically the two propositions are companions: II.5 concerns an unequal cut inside a bisected segment, whereas II.6 concerns an extension beyond the endpoint. Formally their proof DAGs differ. The current II.5 proof does not call II.4, while II.6 uses II.4 directly as one of its two main normal-form decompositions.

6.9 Relationship with Book I Infrastructure

The classical proof uses I.36 and I.43 explicitly. The Lean proof does not replay those steps proposition-locally. Their geometric effect has already been absorbed into the rectangle, parallelogram, square, and scissors infrastructure on which II.1–II.4 are built.

Book I geometry still remains foundational below the current theorem: right-angle propagation, squares, parallelograms, parallel lines, and the finite-scissors representation all depend on the established Hilbert/Book I layers. The gain is architectural: II.6 can be expressed as a composition of stable Book II content interfaces rather than as a second formalization of Euclid’s gnomon diagram.

6.10 Formalization Notes and Hidden Geometric Data

6.10.1 The extension order is proved

The diagram suggests $A - C - B - D$, but the public theorem assumes only midpoint data for C together with $A - B - D$. The private order lemma explicitly proves $C - B - D$ and $A - C - D$. This matters because II.4 can be applied to CD cut at B only after $C - B - D$ is available as a theorem.

6.10.2 Midpoint data have two independent roles

`HilbertIsMidpoint` is not a numerical assertion that C has coordinate one half. Its betweenness component supplies $A - C - B$, while its congruence component supplies $AC \cong CB$. The former feeds the order DAG; the latter feeds rectangle transport.

6.10.3 Representative reuse is deliberate

In the final II.4 call the same Z representative is used twice for $\text{Rect}(CB, BD)$, and the same square $BDFG$ is shared with the Step 6 decomposition. This is not a mathematical restriction on the content identity. It is a deliberate proof-engineering choice that lets the final abstract scissors bridge identify the repeated terms literally and avoids unnecessary representative transports.

6.10.4 The square terms remain geometric representatives

The terms called $Sq(BC)$, $Sq(BD)$, and $Sq(CD)$ are concrete squares supplied through `IsSquare`. They are inserted into the scissors calculus as parallelogram terms. No operation sends a segment to a real number or squares a numerical length.

6.10.5 No gnomon primitive

As in II.5, the formal library does not introduce a special gnomon type. Euclid’s shaded remainder is represented extensionally by a formal sum of rectangle and square terms. In II.6 the combination of II.4 with the abstract scissors bridge replaces the gnomon bookkeeping completely.

6.10.6 No cancellation or subtraction of content

The identity strongly resembles a difference-of-squares calculation, but the proof does not subtract a square or cancel equal rectangles. The abstract bridge is additive: it uses only exact scissors equality, addition, associativity/commutativity of the formal sum, symmetry, and transitivity.

6.11 Axiomatic and Semantic Status

The public theorem assumes

```
[HilbertIncidence Geo]
[HilbertEuclideanPlane Geo]
```

and the production file introduces no new proposition-specific `axiom`, `sorry`, or `admit`. Both the module build and the full project build were reported clean.

The user-verified transitive axiom audit is

```
#print axioms Geometry.euclid_proposition_2_6


'Geometry.euclid_proposition_2_6' depends on axioms: [propext,
Classical.choice,
Geometry.bookZero_nullSegment1,
Quot.sound]
```

The precise status is therefore not “axiom-free.” Rather, II.6 introduces no new proposition-specific axiom and no `sorry`; its reported transitive dependency list is exactly the four declarations above and contains no `sorryAx`.

Semantically, the theorem lies in the exact finite-scissors layer. It proves `HilbertScissorsEq` for the supplied configurations. It does not invoke a numerical area function, a segment multiplication operation, or a general cancellation principle for equal content.

6.12 Dependency Summary

The actual theorem-call DAG can be displayed as



The module chain is slightly different:

Proposition2_5 module $\longrightarrow$ Proposition2_6 module,

but `euclid_proposition_2_5` itself is not called.

The classical local chain is different again:

I.36 + I.43 $\longrightarrow$ gnomon identification $\longrightarrow$ add square on BC $\longrightarrow$ II.6.

The formal reconstruction therefore proves the same geometric-content identity through a different DAG rather than translating Euclid line by line.

New geometric axiom in II.6:	no
New content axiom in II.6:	no
New proposition-local helper theorem:	yes, nine private helpers
New reusable public helper theorem:	no
New <code>HilbertRectangle</code> theorem:	no
New Interface / Book Zero / Scissors theorem:	no
Directly reuses earlier Book II theorem:	yes, II.1, II.3, II.4
Imports II.5 module:	yes
Calls <code>euclid_proposition_2_5</code> :	no
Uses exact <code>HilbertScissorsEq</code> :	yes
Uses general equicomplementability:	no
Uses numerical area:	no
Uses segment multiplication:	no
Uses content cancellation:	no
Public theorem compiles:	yes
Module build:	clean (user verified)
Full project build:	clean (user verified)
Proposition-local <code>sorry/admit</code> :	no
Transitive <code>sorryAx</code> :	absent in reported audit

6.13 Conclusion

Proposition II.6 is a strong checkpoint for the reuse of the Book II rectangle calculus. Classically, Euclid proves the identity by constructing a gnomon inside the square on CD and comparing its

pieces through I.36 and I.43. The current Lean proof reaches the same content equality without a gnomon primitive and without numerical algebra.

Its architecture is instead a composition of three already established Book II mechanisms: II.3 decomposes the rectangle on AD, BD , II.1 splits the old segment AB at its midpoint, and II.4 expands the square on CD at the point B . Midpoint congruence normalizes the repeated cross rectangles, and a small abstract scissors lemma performs the final associative-commutative rearrangement. The result is an exact **HilbertScissorsEq** proof with no new axiom, no **sorry**, no segment multiplication, and no numerical area.

Appendix A

Greenberg: Foundational and Semantic Control

.1 Euclid I.1–I.20 in Greenberg’s Hilbert-Style Development

.1.1 Purpose of the comparison

This appendix records the status of Euclid’s first twenty propositions in Marvin J. Greenberg’s Hilbert-style development of elementary geometry. The comparison is deliberately stricter than a list of mathematically related results. For each proposition I.1–I.20, the question is whether Greenberg provides a corresponding result and, if so, in what form: a complete synthetic proof, an axiom, a proposition whose proof is left as an exercise, a proof outline or hint, or a proof that already uses numerical measurement.

This distinction is important for the present formalization. The fact that the mathematical content of a Euclidean proposition occurs in a modern axiomatic treatment does not imply that a complete synthetic proof, suitable for direct reconstruction in Lean, is actually given there.

.1.2 Comparison table

Euclid	Greenberg counterpart	Status in Greenberg	Relation to the present formalization
I.1	Equilateral triangle construction; explicitly discussed as Euclid’s Proposition 1.	Euclid’s construction is discussed together with its continuity gap. Greenberg repairs the existence argument by means of circular continuity.	The proposition is not simply a consequence of incidence, order, and congruence alone in Greenberg’s presentation. It therefore has a different foundational status from many later propositions.
I.2	Segment laying-off; Congruence Axiom C-1.	The essential construction is incorporated into the congruence axioms rather than repoved as Euclid’s proposition.	In the Hilbert layer this operation belongs naturally to the primitive segment-construction infrastructure.
I.3	Cutting off a segment equal to a given shorter segment.	Its content is distributed between segment construction and segment ordering; no separate reconstruction of Euclid I.3 is required in Greenberg’s organization.	The formal proof is assembled from lower-level construction and order lemmas rather than imported as a named Euclidean theorem.
I.4	SAS triangle congruence.	SAS is Congruence Axiom C-6. Greenberg explicitly discusses why Euclid’s superposition argument is not accepted as an axiomatic proof.	In a Hilbert-style formalization I.4 is therefore foundational rather than a theorem derived from the preceding Euclidean propositions.
I.5	Proposition 3.10: base angles of an isosceles triangle.	A complete synthetic proof is given from SAS.	This is a direct and useful Hilbert-style counterpart of Euclid I.5.
I.6	Proposition 3.18: converse of the isosceles base-angle theorem.	The proposition is stated, but its proof is left as an exercise.	Greenberg confirms the theorem and its place in the theory, but does not supply a complete proof for direct transcription into Lean.
I.7	Uniqueness phenomenon for triangles constructed on a fixed base.	No direct proposition-by-proposition reconstruction of Euclid I.7 was identified. Related uniqueness information is built into the congruence and triangle-laying-off machinery.	The formalization must expose the exact uniqueness statement required by the Euclidean proof rather than relying on a named Greenberg counterpart.
I.8	Proposition 3.22: SSS criterion for congruence.	SSS is stated as a proposition. Its proof is assigned as an exercise, with a substantial hint based on the corollary to SAS and a case analysis.	Greenberg supplies the mathematical route but not a complete printed synthetic proof.
I.9	Angle bisector construction; Proposition 4.4.	The result occurs in the neutral-geometry development, but the proof is assigned to the reader.	The existence of the construction is standard in the theory, while the formal reconstruction still requires the missing synthetic details.
I.10	Midpoint construction; Proposition 4.3 and associated exercise.	Greenberg supplies a detailed step-by-step proof scheme in the exercises, with the individual steps to be justified.	This is close to a proof blueprint, but it is not presented as a complete finished proof in the main text.
I.11–I.12	Proposition 3.16: existence of a perpendicular through a point.	A complete synthetic proof is given, treating separately the cases in which the point lies on the line and does not lie on the line.	The single Hilbert-style proposition covers the geometric content split by Euclid into I.11 and I.12.
I.13–I.14	Supplementary angles, rays, angle addition/subtraction, and related order results.	The required content is distributed through the congruence and betweenness theory rather than organized as two direct Euclidean counterparts.	These propositions illustrate why a proposition-number correspondence is insufficient: the modern proof infrastructure reorganizes the underlying elementary geometry.

.1.3 Result of the query

The comparison shows that Greenberg does *not* contain a continuous sequence of twenty complete synthetic proofs corresponding to Euclid I.1–I.20.

Instead, the first twenty propositions are distributed among several different layers of the Hilbert-style theory. Some Euclidean propositions become axioms or immediate consequences of axioms; some receive complete synthetic proofs; some are stated but assigned as exercises; some are represented by more general modern propositions; and some are treated only after numerical measurement has been introduced.

For the present project, this is a significant distinction. Greenberg is an important reference for the mathematical correctness and dependency structure of the Hilbert reconstruction, but it is not a source from which the Lean proofs of I.1–I.20 can simply be transcribed.

.1.4 Consequences for the formal reconstruction

The comparison also clarifies the role of the intermediate Hilbert library developed in the project. A textbook proof can suppress operations that become explicit obligations in Lean: permutations of collinearity, orientations of betweenness, transport along rays, normalization of unoriented angles, transport of segment and angle order under congruence, and repeated proofs of noncollinearity.

Thus the formal reconstruction has two distinct source levels:

1. the classical mathematical argument, for which Greenberg and related literature provide reference results and proof ideas;
2. the explicit synthetic proof infrastructure required to make those arguments executable in Lean.

The second level is represented by the project’s Book Zero and Hilbert interface. It should not be confused with a claim that the underlying Euclidean theorems are new. Its purpose is to expose and organize the elementary proof operations that conventional mathematical prose routinely leaves implicit.

.1.5 Current checkpoint

At the present checkpoint, Euclid Book I, Propositions I.1–I.20 have been reconstructed in the project. The Greenberg comparison should therefore be read as a source audit for this completed initial block, not as a claim about the status of the remaining propositions of Book I. Further sources can be added to the comparison independently as the literature review proceeds.

Appendix A

Hartshorne: Equal Content and the Algebra of Areas

A.1 Purpose and scope

This appendix records the Hartshorne-side source audit for Euclid Book II at the current checkpoint, Propositions II.1–II.6. Its purpose is not to create a second proposition-by-proposition commentary. Instead, it records what Hartshorne contributes to the semantic interpretation of Euclid’s rectangle and square equalities, which propositions he singles out, and which source-level distinctions should remain stable as Book II grows.

Hartshorne is especially important here because he separates Euclid’s equality of plane figures from numerical area. In Section 22 he defines equal content synthetically and states that the propositions of Book II are valid when Euclid’s equality of figures is read in that sense [5].

A.2 Hartshorne’s organization of Book II

Hartshorne’s treatment places Book II inside the theory of equal content rather than inside a prior arithmetic of numerical lengths. A rectangle described by two segments can therefore play a product-like geometric role without those segments first becoming numbers. This is the background of the familiar interpretation of Book II as an “algebra of areas” [5].

The phrase is structural, not a claim that Euclid has introduced a field of segment lengths. Modern formulas such as

$$a(b + c) = ab + ac$$

are useful mnemonics for the pattern of a geometric decomposition, but the objects compared by Euclid remain plane figures.

Hartshorne’s brief Euclid summary explicitly lists II.1 and II.4, but does not list II.2 or II.3 among its selected Book II entries. In particular, II.4 is recorded as the statement that the square on the whole line equals the squares on its two segments plus twice the rectangle on the two segments [5]. The omission of II.2 and II.3 should not be read as a negative judgment about them: his general area discussion states that the results of Book II are valid for equal content.

A.3 Comparison table

Table A.1: Euclid II.1–II.6 in Hartshorne’s equal-content framework

Euclid	Hartshorne counterpart	Status in Hartshorne	Source-audit remarks and relevance to formalization
II.1	The rectangle contained by two lines is the sum of the rectangles contained by one of them and the segments of the other.	Hartshorne lists II.1 explicitly in his brief Euclid summary and, more generally, places Book II inside the theory of equal content.	This is the cleanest source match for the formal rectangle-distribution API. The Lean reconstruction strengthens this direct-dissection instance to exact <code>HilbertScissorsEq</code> , but that strengthening is a project-level fact and should not be attributed to Hartshorne’s general equal-content relation.
II.2	Square on the whole line decomposed into the two rectangles pairing the whole with the two parts.	II.2 is not separately listed in Hartshorne’s selected brief Book II summary; it is covered by his general statement that the Book II equalities are valid for equal content.	The source supports the synthetic content reading, while the Lean proof exposes an additional representation issue: the two orders of the containing segments are reconciled geometrically by rectangle rotation rather than by numerical commutativity.
II.3	Rectangle on the whole and one part decomposed into the rectangle on the two parts together with the square on that part.	II.3 is likewise not isolated in the selected brief summary; its semantic status is supplied by Hartshorne’s general Book II equal-content result.	The formal theorem is an exact-scissors specialization of the binary II.1 API. The square is used as a representative of the rectangle contained by the same segment twice. Unlike II.2, no side-order transport is required.
II.4	Square on the whole decomposed into the two squares on the parts together with twice the rectangle contained by the parts.	Hartshorne explicitly lists II.4 in his brief Euclid summary, and Section 22 supplies the equal-content semantics for the equality of the figures.	The Lean theorem strengthens the configured case to exact <code>HilbertScissorsEq</code> . It composes II.2 with two II.3 instances and treats “twice the rectangle” as two independent representatives of the same rectangle content, without numerical area or segment multiplication.
II.5	The rectangle on the unequal parts of an internally divided bisected segment, together with the square on the displacement from the midpoint, equals the square on the half.	Hartshorne explicitly singles out II.5, together with II.6 and II.11, in his discussion of the area identities used later in the construction of the regular pentagon. The proposition remains within the synthetic equal-content theory rather than numerical area.	For the formal reconstruction the significant point is that the midpoint condition and the rectangle identity still live below segment arithmetic. The current Lean proof reaches exact <code>HilbertScissorsEq</code> through earlier Book II rectangle decompositions and geometric transport; the modern difference-of-squares formula is only an interpretive shadow.

Continued on next page

Table A.1 – continued from previous page

Euclid	Hartshorne counterpart	Status in Hartshorne	Source-audit remarks and relevance to formalization
II.6	The rectangle contained by the whole extended line and the added segment, together with the square on the half, equals the square on the half plus the added segment.	Hartshorne explicitly mentions II.6 in the same geometrical-algebra block as II.5 and II.11. He also notes that Proposition III.36 is proved by several applications of II.6 together with I.47.	This gives II.6 a source role stronger than a generic Book II example: it is a reusable equal-content identity in Hartshorne’s later Euclidean development. The Lean theorem again strengthens the configured statement to exact <code>HilbertScissorsEq</code> , but does so by II.1, II.3, II.4, midpoint transport, and formal scissors addition rather than by numerical algebra.

A.4 Semantic observations retained from Hartshorne

Several points should remain stable as later propositions are added.

- Euclid’s equality of plane figures in the area block should not be silently replaced by equality of real-valued areas.
- The phrase “rectangle contained by” two segments names a geometric figure through its adjacent side magnitudes; it is not itself a numerical multiplication operation.
- Hartshorne’s equal content is the appropriate general semantic control for Book II. Exact finite equidecomposability is a stronger local conclusion and must be justified proposition by proposition.
- Modern algebraic identities are useful documentary shadows of the synthetic configurations, not substitutes for the geometric proof.
- Hartshorne’s discussion of property (Z) belongs to the converse direction in the theory of content. Nothing in II.1–II.4 requires recovering metric data from a hypothesis of equal content.

A.5 II.1–II.6: from distribution to the two midpoint identities

The first six propositions now display two stages of Book II structure.

II.1–II.4 establish and compose the basic rectangle calculus. II.1 supplies the general additive mechanism, II.2 exposes geometric side-order symmetry, II.3 gives a direct rectangle-plus-square specialization, and II.4 composes those results into the square-of-a-sum pattern.

II.5 and II.6 then add midpoint/order information. They are companion difference-of-squares identities: II.5 treats an unequal cut inside a bisected segment, whereas II.6 treats a segment added beyond the endpoint. Hartshorne explicitly singles out both propositions in his later discussion of geometrical algebra. The formal project keeps the two results in the same synthetic content layer while allowing their Lean DAGs to differ.

For II.6 the modern mnemonic

$$a^2 + (2a + x)x = (a + x)^2$$

should therefore be read exactly as a mnemonic. In the current Lean theorem the two repeated cross rectangles are concrete scissors terms, the square terms are concrete square representatives, and the final equality is obtained by exact-scissors addition and rearrangement.


A.6 Result of the Hartshorne audit

At the II.6 checkpoint Hartshorne supplies the main semantic control for the opening of Book II. His contribution is not a separate proof script for each of II.1–II.6, but a framework in which their equalities are understood as equalities of content between plane figures. II.5 and II.6 are additionally singled out by Hartshorne for their later use in geometrical constructions.

The Lean development refines this framework in two ways that are specific to the current implementation. First, the configured II.1–II.6 theorems land at exact `HilbertScissorsEq`. Second, the formal predicate `IsRectangleContainedBy` makes representative independence and side order explicit. These refinements should remain distinguished from claims made by Hartshorne himself.

A.7 Current checkpoint

This appendix now concerns Euclid Book II, Propositions II.1–II.6. The structural picture is



Within the Lean rectangle calculus, II.1 is the reusable distributive core, II.2 tests geometric side-order symmetry, II.3 returns to direct specialization of the binary II.1 theorem, and II.4 demonstrates composition. II.5 and II.6 add midpoint/order geometry on the internal and external sides of the same classical pattern. Hartshorne’s explicit later use of II.6 also confirms that these identities function as reusable synthetic content lemmas, not merely as isolated exercises in symbolic-looking algebra.

Appendix B

Hilbert: Content and Segment Arithmetic

B.1 Purpose and scope

This appendix records the Hilbert-side source audit for Euclid Book II at the current checkpoint, Propositions II.1–II.6. Hilbert is relevant in two distinct ways: Chapter IV supplies the synthetic theory of plane content, whereas Section 15 later develops an arithmetic of segments. The documentary purpose is to keep those layers separate and to identify which one is actually used by the current formal reconstruction.

B.2 Hilbert’s organization of the relevant theory

Hilbert’s Chapter IV distinguishes finite equidecomposability from the more general relation of equicomplementability. The theory of plane area begins on the basis of Groups I–IV and does not need to be introduced as a numerical area function [2].

For the present project the useful hierarchy is

exact finite dissection $\implies$ equicomplementability $\implies$ later numerical area semantics,

with no converse inserted merely for convenience.

Section 15 is a different layer. Hilbert there defines segment addition and a geometric product, using a fixed unit segment and parallel construction, and then proves the familiar algebraic laws, including commutativity and distributivity [2]. That later arithmetic is a valuable control source precisely because it shows what would be required if Book II were to be formalized through segment multiplication. The current Lean development does not take that route.

B.3 Comparison table

Table B.1: Euclid II.1–II.6 against Hilbert’s content and segment-arithmetic layers

Euclid	Hilbert counterpart	Status in the Grundlagen	Source-audit remarks and relevance to formalization
II.1	Chapter IV finite decomposition and content theory; later Section 15 distributivity of segment multiplication is a structural analogue.	No one-to-one numbered Hilbert theorem reproducing Euclid II.1 was located in the current audit. The relevant material is organized by content theory and, later, by segment arithmetic rather than by Euclid’s Book II numbering.	The formal proof belongs to the earlier content/dissection layer. Additivity is obtained from an actual rectangle cut and representative transport, not by invoking the later distributive law for a segment product.
II.2	Chapter IV content plus the geometric symmetry of a rectangle; Section 15 commutativity is only a later arithmetic analogue.	Again the source correspondence is structural rather than proposition-level. Hilbert’s later commutative segment multiplication must not be read backward into the scissors proof.	The Lean theorem makes the distinction concrete: the required exchange of the two containing sides is implemented by cyclic rotation of a rectangle representative, preserving exact scissors content.
II.3	Chapter IV content/dissection; later Section 15 distributivity gives the arithmetic pattern only after segment multiplication has been defined.	No separate proposition-level Hilbert counterpart was located. The source supports the distinction between geometric content operations and the later arithmetic of segments.	II.3 is a direct specialization of the binary II.1 cut with the chosen segment used as the fixed side. The square is packaged as a rectangle contained by the same segment twice. No segment product and no side-order transport are needed.
II.4	Chapter IV finite content composition; later Section 15 distributivity and commutativity give an arithmetic analogue of the square expansion.	No one-to-one Hilbert proposition corresponding to Euclid II.4 was located in the current audit. Hilbert’s source role remains architectural.	The formal theorem composes exact II.2 with two exact II.3 instances and a geometric rectangle rotation. Despite the modern shadow $(a + b)^2 = a^2 + b^2 + 2ab$, no segment product, unit segment, or numerical area is used.
II.5	Chapter IV content/dissection; Section 15 segment multiplication supplies only a later arithmetic analogue of the identity.	No one-to-one Hilbert proposition corresponding to Euclid II.5 was located in the current audit. The source role remains the separation of synthetic content from the later arithmetic of segments.	The formal theorem combines midpoint and order information with exact II.1 and II.3 rectangle decompositions, representative transport, and scissors addition. The modern shadow $(a - x)(a + x) + x^2 = a^2$ is not used as a proof object; no segment product, unit segment, numerical area, or cancellation axiom is introduced.

Continued on next page

Table B.1 – continued from previous page

Euclid	Hilbert counter-part	Status in the Grundlagen	Source-audit remarks and relevance to formalization
II.6	Chapter IV content/dissection; Section 15 segment multiplication supplies only a later arithmetic analogue of the external-section identity.	No one-to-one Hilbert proposition corresponding to Euclid II.6 was located in the current audit. The relevant source role remains the separation between synthetic content and later segment arithmetic.	The formal theorem combines order from the midpoint/extension configuration with exact II.1, II.3, and II.4 decompositions, rectangle transport, and a purely additive scissors bridge. The modern shadow $a^2 + (2a + x)x = (a + x)^2$ is not a proof object; no segment product, unit segment, numerical area, subtraction, or cancellation axiom is introduced.

B.4 Structural observations retained from Hilbert


The following distinctions should be preserved as Book II grows.

- Plane content and segment arithmetic are separate theories in the Grundlagen and should remain separate in the documentation.
- Exact finite dissection is stronger than the general content relation; when Lean proves `HilbertScissorsEq`, that fact comes from the local proof architecture rather than from a blanket identification of all equal-content figures with equidecomposable ones.
- Segment multiplication is not primitive. It requires a unit segment, proportion geometry, and parallel constructions before algebraic laws such as commutativity and distributivity can even be stated.
- Consequently, a distributive-looking Euclidean proposition can be formalized below the arithmetic layer when its geometry is literally a finite cut-and-paste argument.

B.5 II.1–II.6: exact dissection below segment arithmetic

The opening six propositions give a useful stress test of the separation. II.1 establishes a reusable exact rectangle split. II.2 shows that the apparent exchange of factors can still be implemented geometrically by rotating a rectangle presentation. II.3 then shows that even this extra symmetry operation is unnecessary when the ordered side data already match the binary II.1 interface. II.4 is the first compositional step: it combines II.2 with two II.3 expansions and uses only exact scissors addition and representation transport. II.5 then adds midpoint and order data and composes the same rectangle/scissors mechanisms into the internal unequal-parts identity. II.6 moves the second cut outside the original segment and, in the current Lean DAG, uses II.4 directly to normalize the square on the produced segment. Both remain entirely below segment arithmetic.

The resulting formal chain is therefore



not

segment field $\longrightarrow$ multiplication $\longrightarrow$ algebraic simplification.

B.6 Result of the Grundlagen audit

At the II.6 checkpoint Hilbert's main contribution remains architectural. Chapter IV validates the use of a synthetic content layer independently of numerical area, while Section 15 marks a later arithmetic boundary. The current Book II formalization through II.6 remains on the earlier side of that boundary.

The rectangle API therefore has a clear foundational interpretation: congruent segments and right angles determine representative rectangles, and the scissors layer compares their content. The algebra-like behavior observed in II.1–II.3 is derived from that geometry.

B.7 Current checkpoint

This appendix now concerns Euclid Book II, Propositions II.1–II.6. At this stage the durable Hilbert-side picture is

Groups I–IV geometry $\longrightarrow$ plane-content/dissection layer $\longrightarrow$ rectangle calculus for II.1–II.6,

while Section 15 segment arithmetic remains a later comparison layer rather than a dependency of the proofs.

Appendix C


Borsuk–Szmielw: Axiomatic Layers and Representation

C.1 Why this source matters

Karol Borsuk and Wanda Szmielw organize elementary geometry in a way that is unusually close to the architectural questions arising in a formal reconstruction.

Their primitive geometric language consists of points, lines, planes, betweenness, and equidistance. The axioms are divided into five groups: incidence, order, congruence, continuity, and finally either the Euclidean axiom E or the Bolyai–Lobachevskian axiom. The first four groups constitute absolute geometry.

More important for the present purpose, the authors explicitly state that one should study not only which theorems are true, but also which means are necessary to prove them. Their exposition is therefore deliberately layered:



This makes the book especially useful for deciding where a theorem belongs in a formal library. A result corresponding to a Euclidean proposition may turn out to be:

- part of the incidence/order language;
- a consequence of congruence alone;

- available before continuity;
- available only after continuity;
- genuinely Euclidean, requiring E ;
- or absent as a named theorem because Borsuk–Szmielew replace it by a stronger structural result.

The last column of the table below therefore asks a practical question: what does the Borsuk–Szmielew placement suggest for the Hilbert interface and the Book Zero layer?

C.2 Layer notation

For compactness the table uses the following descriptions.

Layer	Meaning
Inc/Ord	Chapter I: incidence and order, including betweenness, half-lines, half-planes, angles as ordered ray configurations, triangles, quadrangles, Pasch-type results, polygons, and triangulation.
Congr-no-Co	Chapter II: congruence added, but the Axiom of Continuity is not used.
Absolute	Chapter III: the Axiom of Continuity is added; the full first four groups of axioms are now available.
Euclidean E	Chapter V: absolute geometry plus the Axiom of Euclid E , equivalent in the book to Euclid’s Fifth Postulate.

C.3 Euclid in the Borsuk–Szmielew development

Table C.1: Euclidean propositions in the axiomatic layers of Borsuk–Szmielew

Euclid	Borsuk–Szmielew counterpart	Layer	Structural significance for the reconstruction
I.1	No one-to-one construction theorem was located. Equilateral triangles are defined in Chapter II, but the audited text does not reproduce Euclid’s two-circle construction as a named theorem.	No direct counterpart	This is a useful absence. Borsuk–Szmielew do not organize the congruence theory by reconstructing Euclid’s first circle construction. The formal project should therefore keep the classical construction DAG separate from the axiomatic layer in which an equilateral-point existence theorem is eventually supplied.
I.2	Axiom C5: on every half-line with origin a , for every segment pq , there exists exactly one point b such that $ab = pq$.	Congr-no-Co	Euclid’s segment transport has become primitive and is strengthened to existence plus uniqueness on a prescribed half-line. This is precisely interface-level segment layoff, not a proposition that later proofs should reconstruct from circles.

Continued on next page

Table C.1 – continued from previous page

Euclid	Borsuk–Szmielew counterpart	Layer	Structural significance for the reconstruction
I.3	Axiom C5 together with segment order and betweenness.	Congr-no-Co	Cutting a prescribed length from a longer segment is absorbed into ray layoff plus order. The natural formal location is a small Book Zero lemma over segment layoff and betweenness.
I.4	Theorem 22(I): two sides and the included angle determine a congruent triangle. The text immediately records this as the SAS congruence criterion.	Congr-no-Co	SAS is developed before continuity. This is strong evidence for keeping it in the neutral congruence interface. Euclid’s superposition argument is not part of the Borsuk–Szmielew proof architecture.
I.5	Theorem 23: in a triangle, congruent sides are opposite congruent angles; in particular, the base angles of an isosceles triangle are congruent.	Congr-no-Co	The isosceles theorem is packaged together with its converse in one biconditional result. The reusable API should reflect this stronger relationship instead of treating I.5 and I.6 as unrelated endpoints.
I.6	The converse direction of Theorem 23: congruent opposite angles imply congruent opposite sides.	Congr-no-Co	Borsuk–Szmielew eliminate Euclid’s separate reductio construction. Once segment and angle congruence are organized properly, I.6 is simply the reverse implication of the same structural theorem as I.5.
I.7	No separate theorem corresponding exactly to Euclid I.7 was located. Its uniqueness content is absorbed into the order theory, unique angle construction (Theorem 13), and the strong congruence axioms.	Congr-no-Co	This is an example of a Euclidean proposition disappearing under a stronger interface. The formal question is not how to imitate I.7, but which general uniqueness and side-of-line lemmas make its role unnecessary.
I.8	Theorem 21: if the three corresponding sides of two triangles are congruent, then the three corresponding angles are congruent.	Congr-no-Co	This is the SSS content in the Borsuk–Szmielew organization, and it is available without continuity and without the Euclidean axiom. For the formal project this strongly supports treating SSS as derived neutral congruence infrastructure rather than Euclidean input.
I.9	Theorems 33–34: a bisector lies between the sides of the angle, and every angle has exactly one bisector.	Congr-no-Co	The source gives a stronger result than Euclid’s bare construction: existence and uniqueness are packaged together. Angle-bisector existence therefore belongs naturally in the reusable interface.
I.10	Theorems 31–32: an equidistant collinear point is between the endpoints, and every segment has exactly one midpoint.	Congr-no-Co	This is especially valuable formally. Borsuk–Szmielew do not take the diagrammatic position of the midpoint for granted; they first prove the betweenness theorem and only then existence and uniqueness. This is exactly the separation needed in Lean.
I.11	Theorem 76: through a point lying on a line there exists exactly one line in the plane perpendicular to the given line.	Congr-no-Co	Perpendicular erection is already a neutral interface theorem. Later propositions should consume it directly rather than replay an auxiliary Euclidean triangle construction.

Continued on next page

Table C.1 – continued from previous page

Euclid	Borsuk–Szmielew counterpart	Layer	Structural significance for the reconstruction
I.12	Theorem 77: through a point outside a given line there exists exactly one perpendicular to that line.	Congr-no-Co	Dropping a perpendicular is likewise available before continuity. The theorem is stronger and cleaner than Euclid’s construction-oriented formulation and belongs naturally in the Hilbert interface.
I.13	Sections 21–24 of Chapter II: addition of free angles, adjacent angles, right angles, and the fact that an angle adjacent to a right angle is right (Theorems 63, 69–71).	Congr-no-Co	Borsuk–Szmielew replace informal angle arithmetic by an explicit calculus of free angles. This is a strong source precedent for using synthetic supplement/addition relations rather than numerical degrees in the formalization.
I.14	Theorem 13 gives unique construction of a congruent angle on a specified side; Theorems 14–15 control adjacent angles and their uniqueness.	Congr-no-Co	The converse straight-line criterion is absorbed into a more general uniqueness theory for angle placement. The formal analogue should be a small collinearity/adjacency consequence of that interface.
I.15	Theorem 16: vertical angles are congruent.	Congr-no-Co	A direct reusable theorem. Its low position in the congruence chapter confirms that vertical-angle congruence should be generic infrastructure for later triangle and parallel arguments.
I.16	Theorem 35: an exterior angle of a triangle is greater than each nonadjacent interior angle.	Congr-no-Co	This is one of the central neutral theorems. The proof explicitly uses midpoint, vertical-angle congruence, SAS, and angle order. It is also the engine for constructing nonintersecting lines before the Euclidean axiom enters.
I.17	A short consequence of Theorem 35 together with the developed theory of adjacent and right angles. It is not necessary to use the later Saccheri–Legendre theorem for this Euclidean proposition.	Congr-no-Co	This distinction prevents an important misclassification. Borsuk–Szmielew later prove stronger angle-sum results after continuity, but Euclid I.17 itself already belongs to the earlier neutral congruence layer.
I.18	Theorem 40, one direction: in a triangle the greater side lies opposite the greater angle.	Congr-no-Co	I.18 and I.19 are unified. Borsuk–Szmielew package strict side and angle comparison into a biconditional theorem, exactly the sort of strong reusable comparison lemma useful in Book Zero.
I.19	Theorem 40, converse direction: the greater angle lies opposite the greater side.	Congr-no-Co	The converse is not a separate construction. It is the reverse direction of the same strict comparison equivalence used for I.18.
I.20	Theorem 50: for distinct points a, b, c , $[ab] + [bc] > [ac]$, with equality exactly when $B(a, b, c)$.	Congr-no-Co	This is substantially stronger than the triangle-only formulation. It combines triangle inequality with the collinear equality case and uses the explicit algebra of free segments. It is an excellent model for a robust segment-order interface.

Continued on next page

Table C.1 – continued from previous page

Euclid	Borsuk–Szmielw counterpart	Layer	Structural significance for the reconstruction
I.21	Theorem 52 generalizes the side-sum part to a convex polygonal line inside a triangle; the angle part is controlled by the Exterior Angle Theorem 35 and the order theory of rays.	Congr-no-Co	The source replaces the one-off proposition by a more general polygonal inequality. For formal work the useful target is therefore the reusable generalized lemma rather than the exact Euclidean diagram.
I.22	No one-to-one prescribed-three-sides construction theorem was located in the audited elementary sequence. The book has the triangle inequality and a strong congruence construction axiom, but does not replay Euclid’s circle-intersection construction here.	No direct counterpart	This should be kept as an explicit gap in the source map. Existence of a triangle with three prescribed side lengths is logically different from SSS uniqueness/congruence and should not be silently identified with Theorem 21.
I.23	Theorem 13: for a given ray, chosen half-plane, and given angle, there exists exactly one ray on that side making a congruent angle.	Congr-no-Co	Euclid’s angle-copy construction becomes a primitive-style transport theorem with uniqueness. This is one of the strongest direct precedents for the angle-construction API already used in the Hilbert route.
I.24	Theorem 41: with two corresponding sides fixed, comparison of the included angles is equivalent to comparison of the third sides.	Congr-no-Co	Borsuk–Szmielw combine the Euclidean comparison theorem and its converse into one biconditional. This is a natural interface theorem for replacing configuration-heavy local proofs.
I.25	The converse direction of Theorem 41.	Congr-no-Co	As with I.18/I.19, the source suggests exposing one symmetric comparison theorem rather than maintaining two large proposition-sized proofs.
I.26	Theorem 22(II): one side and the two angles adjacent to it determine a congruent triangle (ASA).	Congr-no-Co	The ASA part of I.26 is explicit and neutral. The audited text does not package Euclid’s second AAS configuration under a separate theorem number here, so the appendix does not silently identify the two formulations.
I.27	Theorems 37–39. Equal appropriate transversal angles give nonintersection; Theorem 39 concludes that through an external point there is at least one line not intersecting the given line.	Congr-no-Co	This is one of the most important source placements in the book. The angle-to-parallel direction and existence of a nonintersecting line are established before continuity and before the Euclidean axiom.
I.28	A consequence of the same Theorems 37–38 together with the earlier adjacent/vertical-angle normalization theorems.	Congr-no-Co	The two Euclidean angle criteria are wrappers over the neutral nonintersection theorem. The conclusion “parallel” does not by itself make the result Euclidean.

Continued on next page

Table C.1 – continued from previous page

Euclid	Borsuk–Szmielew counterpart	Layer	Structural significance for the reconstruction
I.29	After E , Chapter V develops the Euclidean characterization of parallel lines and Theorem 10: angles whose corresponding sides are parallel are congruent.	Euclidean E	This is the reversal of I.27. Before E , appropriate angle data produce nonintersection; after E , parallelism can be converted back into angle congruence. The source therefore marks the same neutral-to-Euclidean boundary that the formal reconstruction uses.
I.30	Chapter V, Theorem 7: parallelism of lines is transitive.	Euclidean E	Parallel transitivity is explicitly downstream of the Euclidean axiom. It should therefore be classified as Euclidean interface infrastructure, not as a neutral convenience lemma.
I.31	Chapter II, Theorem 39 gives existence of a non-intersecting line. Axiom E plus that theorem gives Chapter V, Theorem 1, uniqueness; Theorem 6 states the corresponding unique parallel through a point.	Existence: Congr-no-Co; uniqueness: Euclidean E	Borsuk–Szmielew make the existence/uniqueness split exceptionally clean. This is a direct architectural model for keeping the two layers separate in Lean.
I.32	Chapter V, Theorem 2: the sum of the angles of every triangle is π .	Euclidean E	The equality form of the triangle angle-sum theorem is placed immediately after the Euclidean axiom. A synthetic Lean statement need not use numerical measure, but its dependency must remain Euclidean.
I.33	Chapter V, Theorem 12, sufficiency direction: a plane quadrangle is a parallelogram when one pair of corresponding directed sides is parallel and the corresponding segments are congruent.	Euclidean E	This theorem is stronger and more useful than a literal copy of I.33. It characterizes a parallelogram from exactly the sort of parallel-plus-congruence data that arise in I.33. It is a strong candidate for reusable interface infrastructure.
I.34	Chapter V, Theorem 12, necessity direction and proof: in a parallelogram the diagonal triangles are congruent, yielding equality of opposite sides and the corresponding angle data.	Euclidean E	The proof architecture is almost tailor-made for the formal I.34 interface: parallel half-lines give angle congruences, then triangle congruence supplies opposite-side data. I.34 should expose this package for immediate downstream use.
I.35	No Euclidean equal-content or equicomplementability theorem corresponding directly to I.35 was located. Chapter I already has a substantial theory of polygons and triangulation, while Chapter V has the parallelogram theory needed immediately before I.35.	No direct counterpart	The absence is informative. Borsuk–Szmielew supply rigorous figure/triangulation infrastructure and Euclidean parallelogram infrastructure, but they do not insert a Hilbert/Hartshorne-style equal-content layer at this point. Thus they are highly useful for the components feeding I.35, but not the primary source for the semantics of its conclusion.

C.4 The strongest architectural correspondences

The most useful feature of the Borsuk–Szmielw source is that several Euclidean propositions are replaced by stronger interface theorems.

C.4.1 Transport is primitive-strength infrastructure

Axiom C5 is stronger than the bare construction in I.2: it gives exactly one point on a chosen half-line at a prescribed distance. Theorem 13 is the angular analogue: on a chosen side of a ray, an angle congruent to a given angle can be constructed uniquely.

Thus two operations that appear relatively late or require elaborate construction in Euclid become basic transport mechanisms:

segment layoff and angle layoff.

This is directly consonant with the Hilbert-style formal interface.

C.4.2 Pairs of Euclidean propositions become biconditionals

Theorem 23 unifies I.5 and I.6:

equal sides $\iff$ equal opposite angles.

Theorem 40 unifies I.18 and I.19:

greater side $\iff$ greater opposite angle.

Theorem 41 similarly unifies I.24 and I.25 when two corresponding sides are fixed:

greater included angle $\iff$ greater third side.

For library design, these are more informative than the proposition numbers. They identify the natural reusable theorem boundaries.

C.4.3 Midpoints and bisectors include positional information

The midpoint discussion is especially close to the needs of formal geometry. Borsuk–Szmielw first prove that a collinear point equidistant from the endpoints of a segment must lie between them. Only then do they prove existence and uniqueness of the midpoint.

The angle-bisector discussion has the same pattern: first the bisector is shown to lie between the sides of the angle, then existence and uniqueness are established.

The lesson for formalization is precise. “Midpoint” and “bisector” should not be treated merely as metric equalities; their order/ray placement is part of the reusable theorem package.

C.5 The neutral parallel theory is already large

The source is particularly valuable around Euclid I.27–I.31.

From the Exterior Angle Theorem and the congruence/order theory, Borsuk–Szmielew prove that an appropriate equality of transversal angles forces two lines to be disjoint. They then obtain Theorem 39:

through an external point there exists at least one nonintersecting line.

This theorem belongs to Chapter II, before continuity and before the Euclidean axiom.

Only much later, in Chapter V, is Axiom *E* added:

through an external point there is at most one nonintersecting line.

Together the two statements give uniqueness. Chapter V then derives the Euclidean parallel theory: characterization of disjoint coplanar lines, uniqueness of the parallel, transitivity, angles with parallel sides, and parallelograms.

This produces a clean formal boundary:

angle criterion $\Rightarrow$ nonintersection	before <i>E</i> ,
existence of a nonintersecting line	

while

uniqueness of the parallel	after <i>E</i> .
parallelism $\Rightarrow$ angle congruence	
parallel transitivity	
triangle angle sum	

For the present reconstruction this is one of the strongest independent checks on the placement of I.27, I.29, I.30, I.31, and I.32.

C.6 Theorem 12 of Chapter V as an interface theorem


The parallelogram theorem deserves special attention.

Borsuk–Szmielew define a parallelogram in the expected way, by two pairs of opposite sides lying on parallel lines. Their Theorem 12 then proves a necessary and sufficient condition using only one pair of directed parallel sides and equality of the corresponding segments.

In the necessity direction, the proof starts from a parallelogram, transports angles using parallel half-lines, and obtains congruence of the two triangles cut by a diagonal. Opposite-side congruence follows.

In the sufficiency direction, one parallel pair together with one segment congruence forces the remaining pair to be parallel.

This theorem simultaneously explains why I.33 and I.34 should sit close to the reusable Euclidean parallelogram interface:



It is therefore not merely a historical counterpart. It is a plausible API boundary for subsequent formal work.

C.7 What the book contributes at I.35


The transition at I.35 is revealing because the source provides two sides of the problem but not the final semantic relation.

Chapter I develops polygons, intersections of polygons with half-planes, and triangulation. This is a rigorous combinatorial theory of plane figures built very low in the axiomatic hierarchy.

Chapter V develops the Euclidean parallelogram theory and supplies the congruence infrastructure immediately preceding I.35.

What is not developed there is a Euclidean theory of equal content or equicomplementability comparable to Hilbert Chapter IV or Hartshorne Section 22. Later references to area in the Bolyai–Lobachevskian part do not supply such a replacement; the authors explicitly say that they will not develop a general theory of area there.

For the formal reconstruction the source therefore has a sharply defined role:



but

Hilbert/Hartshorne remain the control sources for equicomplementability/equal content.

This is not a weakness of the audit. It tells us exactly where one source stops and another must take over.

C.8 Consequences for the formal library

The Borsuk–Szmielw audit suggests a practical classification of reusable results.

C.8.1 Hilbert interface candidates

The following belong naturally to a general synthetic interface rather than to individual Euclidean proposition files:

- unique segment layoff on a ray;
- unique angle layoff on a chosen side;
- SAS and SSS-style triangle congruence;
- isosceles theorem and converse;
- midpoint existence, uniqueness, and betweenness;
- angle-bisector existence, uniqueness, and ray order;
- vertical angles;
- Exterior Angle Theorem;
- strict side/angle comparison;
- perpendicular existence and uniqueness;
- neutral angle criteria for nonintersection.

C.8.2 Book Zero candidates

The developed calculus of free segments and free angles also points to a natural macro-language:

- segment less-than and trichotomy;
- transport of order under congruence;
- segment addition and subtraction;
- triangle inequality with the collinear equality case;
- angle less-than and trichotomy;
- angle addition and adjacency;
- supplement/right-angle normalization.

These are exactly the sorts of operations that prevent later proposition files from expanding into repeated low-level incidence and order proofs.

C.8.3 Euclidean interface candidates

After Axiom *E*, the natural reusable layer contains:

- unique parallel through a point;
- characterization of Euclidean parallels;
- parallel transitivity;
- congruence of angles with respectively parallel sides;
- Euclidean triangle angle sum;
- the Chapter V parallelogram characterization.

The source thus gives not just another list of theorem counterparts, but a plausible decomposition of the formal library itself.

C.9 Conclusion

Borsuk and Szmielew are particularly valuable for the reconstruction because their book asks almost the same structural question as the formal project: not only whether a theorem is true, but which axiomatic resources are actually needed to prove it.

Their treatment makes several boundaries unusually visible. Much of the elementary order and congruence theory, including the Exterior Angle Theorem, midpoints, bisectors, perpendiculars, triangle comparison, and existence of nonintersecting lines, is developed before continuity. The Axiom of Euclid then contributes the uniqueness needed to obtain the full Euclidean theory of parallels. Their parallelogram theorem packages the material around I.33–I.34 in a form that is especially attractive as formal interface infrastructure.

At I.35 the source becomes useful in a different way. Its early theory of polygons and triangulation and its later parallelogram theory provide important components of the proof, while the semantic theory of equal content is left to other sources.

The resulting appendix should therefore be read as an axiomatic-layer map and an API-design aid, not merely as another historical comparison with Euclid.

Bibliography

- [1] Euclid, *The Thirteen Books of Euclid's Elements*, edited and translated by Sir Thomas L. Heath, 3 vols., Dover Publications, New York, 1956. (Reprint of the second edition, Cambridge University Press, 1926.)
- [2] David Hilbert, *Grundlagen der Geometrie*, B. G. Teubner, Leipzig, 1899.
- [3] Karol Borsuk and Wanda Szmielew, *Foundations of Geometry: Euclidean and Bolyai–Lobachevskian Geometry, Projective Geometry*, North-Holland Publishing Company, Amsterdam, 1960.
- [4] Marvin Jay Greenberg, *Euclidean and Non-Euclidean Geometries: Development and History*, 4th ed., W. H. Freeman and Company, New York, 2008.
- [5] Robin Hartshorne, *Geometry: Euclid and Beyond*, Springer, New York, 2000.